

Elementary Particles I

3 – Strong Interaction

Resonances, Isospin, Strangeness, Unitary Symmetries

Strong Interaction

Originally pictured as an attractive force between nucleons, required to overcome repulsive Coulomb interaction in nuclei

Main features:

- *Strength*
- *Short range*
- *Charge independence*

For a long time, difficult to understand: lot of guesswork, many models

Today, believed to be a *residual force* between 'color neutral' particles (hadrons), a remnant of color interaction between quarks and gluons

Somewhat similar to Van der Waals/Covalent bond between 'neutral' molecules, coming from electromagnetic interaction between charged electrons and nuclei

Yukawa Theory

First attempt to model strong interaction after the electromagnetic:
Exchange of mediator particles → Prediction of *pion*

Mass > 0

Limited range

Spin ≠ 1

Vector particle would yield
repulsive forces between identical particle

Charged, Neutral

Same force for pp , nn , pn

Electromagnetism

$$\frac{\partial^2 \varphi}{\partial t^2} - \nabla^2 \varphi = -\rho \quad \text{Wave equation}$$

Scalar potential

$$\nabla^2 \varphi = \rho \quad \text{Static case}$$

$\rho_G(\mathbf{r}) = e\delta(\mathbf{r})$ Point source
at the origin

$$\rightarrow \varphi_G(\mathbf{r}) = \frac{e}{r} \quad \text{Green's function}$$

≡ Coulomb potential

Yukawa

$$\frac{\partial^2 \varphi}{\partial t^2} - \nabla^2 \varphi - m^2 = -\rho \quad \text{Wave equation}$$

Pion field

$$\nabla^2 \varphi + m^2 = \rho \quad \text{Static case}$$

$\rho_G(\mathbf{r}) = g\delta(\mathbf{r})$ Point source
at the origin

$$\rightarrow \varphi_G(\mathbf{r}) = \frac{g e^{-mr}}{r} \quad \text{Green's function}$$

≡ Yukawa potential

Pions

Discovered after the II World War (Cosmic Rays, Accelerators)
Properties

Mass	$\begin{cases} 135 \text{ MeV} & \text{Neutral} \\ 139 \text{ MeV} & \text{Charged} \end{cases}$
Spin	0
Parity	-
Charge parity	+
Lifetime	$\begin{cases} 25 \cdot 10^{-9} \text{ s} & \text{Charged} \\ 10^{-16} \text{ s} & \text{Neutral} \end{cases}$
Decay modes (Dominant)	$\begin{cases} \mu\nu & \text{Charged} \\ \gamma\gamma & \text{Neutral} \end{cases}$

Stable vs. strong decays, as the *lightest hadron*
Copiously produced at first accelerators (synchrocyclotrons)
Charged pions easily focused into collimated, high energy beams



Scattering

As for electromagnetic, strong interaction can be investigated by scattering experiments

Perform experiments like

$$p + p, \quad p + n, \quad \pi^\pm + p, \quad \pi^\pm + n$$

Pion: Spinless \rightarrow Understanding πN scattering easier than NN



Total cross section plots - Observe lot of structure

Propagators in the s-channel - I


Take first a QED example: Bhabha scattering at $\sqrt{s} \ll M_{Z^0}$

$$e^- + e^+ \rightarrow e^- + e^+$$

Two one-photon diagrams

$$\text{Virtual photon propagator} = \frac{1}{q^2}$$

t-channel: Virtual photon has $q^2 < 0$ space-like
s-channel: Virtual photon has $q^2 > 0$ time-like




t-channel s-channel

Taking radiative corrections to one loop

$$\text{Virtual photon propagator} = \frac{1}{q^2 \left(1 - \bar{\Pi}_\gamma^{(2)}(q^2)\right)}$$

Correction resulting from fermion e.m. currents circulating in the loop, after renormalization



Propagators in the s-channel - II

Among all fermion currents circulating in the loop, take a muon pair.

Now, a $\mu^+\mu^-$ pair has bound states, like a hydrogen atom.

For these, total energy is $< 2m_\mu$: Binding energy < 0

When $\sqrt{s} = E_{CM} \sim M_{bound} \equiv M$

$$q^2 = s = E_{CM}^2$$

$$\rightarrow \frac{1}{q^2 \left(1 - \bar{\Pi}_\gamma^{(2)}(q^2)\right)} \approx \frac{1}{q^2 - M(M - i\Gamma)} = \frac{1}{E_{CM}^2 - M^2 + iM\Gamma}$$

$$\rightarrow \frac{1}{q^2 \left(1 - \bar{\Pi}_\gamma^{(2)}(q^2)\right)} \approx \frac{1}{(E_{CM} - M) \underbrace{(E_{CM} + M)}_{\approx 2M} + iM\Gamma} \approx \frac{1}{2M} \frac{1}{(E_{CM} - M) + i\Gamma/2} \quad \text{Breit-Wigner form}$$

This term tied to bound state being *unstable*: $\Gamma > 0 \rightarrow \tau = 1/\Gamma < \infty$
Unlike the H atom, muonic atom annihilates into various channels

The existence of bound states for the current coupled to the photon is reflected into resonant behavior of the s-channel scattering amplitude

NB Resonant peaks in total, elastic $e^+ e^-$ cross-section not observed because of their exceedingly small width

Propagators in the s-channel - III

General rule:

Every time the intermediate state can couple to an unstable state (excited bound state, genuine elementary particle coupled to decay channels, ...), the s-channel propagator shows resonant behavior when the total energy is close to the mass of the unstable state

Δ -Resonance: Formation

First observed by Fermi and collaborators in πN scattering (1951)

With some caveats, can be considered as a kind of excited nucleon state

$$\pi^+ + p \rightarrow \Delta^{++} \rightarrow \pi^+ + p$$



Different spin,
quark content

Also observed in other charge states Δ^+ , Δ^- , Δ^0 and in many different processes (strong, e.m. and weak)

Some analogy with photon excitation of atomic levels


$$\gamma + A \rightarrow B \rightarrow \gamma + A, \quad A \text{ ground state, } B \text{ excited level}$$

Good indication that the nucleon is a *composite* object






Discovery of Δ - 1951


Chicago synchrocyclotron




Collect first data on 2-body reactions:



Plastic scintillators



Δ^{++} Resonance



Propagators in the t -channel - I

The same propagator describes the t -channel amplitude, $t=q^2<0$:

$$\frac{1}{q^2 \left(1 - \bar{\Pi}_{\gamma}^{(2)}(q^2)\right)} \approx \frac{1}{q^2 - M(M - i\Gamma)} \underset{\Gamma \ll M}{\approx} \frac{1}{q^2 - M^2} \quad \text{'Pole' amplitude}$$

In this case, there is *no* resonant behavior: $q^2 - M^2 < 0$ strictly

Rather, the amplitude can be seen as an extension of the virtual photon idea, corresponding to the exchange of a *virtual particle*, with mass M and width Γ , or lifetime $1/\Gamma$. In the previous example, it would be a *virtual muonic atom*.

As for the virtual photon, the virtual particle exchanged is said to be *off mass-shell*:

$$q^2 \neq M^2$$

Propagators in the t -channel - II

Besides being very appealing as a qualitative visualization of processes, this interpretation also appears to be superficially consistent with perturbation theory. But...

...It is unfortunately not very useful as a tool for quantitative work in strong interactions physics, just because perturbative expansion cannot be maintained for strong coupling constant.

Most simply, diagrams with more than one particle exchanged correspond to amplitudes *larger* than diagrams with just one...

One π Exchange \leftrightarrow Yukawa Potential

Nevertheless, just as an interesting exercise:

Take NN scattering at small q^2 as due to *one pion exchange*: This can be maintained, to some extent (or so one believes).

Then


$$A \propto \frac{1}{q^2 - m_\pi^2}$$

In the static potential limit

$$E_C \approx E_A$$

$$q^2 = (E_C - E_A)^2 - (\mathbf{p}_C - \mathbf{p}_A)^2 \approx -(\mathbf{p}_C - \mathbf{p}_A)^2 = -|\mathbf{q}|^2$$

$$\rightarrow \frac{1}{q^2 - m_\pi^2} \approx \frac{1}{-|\mathbf{q}|^2 - m_\pi^2} = -\frac{1}{|\mathbf{q}|^2 + m_\pi^2}$$



Assuming Born approximation as valid here

$$V(r) \propto \int e^{i\mathbf{q}\cdot\mathbf{r}} \left(-\frac{1}{|\mathbf{q}|^2 + m_\pi^2} \right) d^3\mathbf{q} \propto -\frac{e^{-m_\pi r}}{r} \quad \text{Yukawa potential}$$

Potential Scattering

Formalism of potential scattering:

Not a proper tool to describe relativistic regime (particle creation/destruction) → Go for Field Theory

Nevertheless:

Believed to be somewhat useful to get insight into simplest (2-body) reactions, like elastic scattering, even at high energy

Phase shifts analysis:

Try to reconstruct the interaction structure from scattering data

Past: Lot of work spent in the attempt of modeling 'simplest' reactions (e.g. Mandelstam representation, Regge poles, ...)

Now: The 'simplest' reactions finally understood to be quite complicated, much more than anticipated (← Non perturbative interaction regime)

Phase Shifts and Resonances - I

Partial waves expansion

$$d\sigma = v \frac{|f|^2}{v} d\Omega = |f|^2 d\Omega \rightarrow \frac{d\sigma}{d\Omega} = |f|^2$$

Scattering amplitude:

$$f(\theta) = \frac{1}{2ik} \sum_{l=0}^{\infty} (2l+1)(e^{2i\delta_l} - 1) P_l(\cos \theta) = \frac{1}{k} \sum_{l=0}^{\infty} (2l+1) \frac{e^{2i\delta_l} - 1}{2i} P_l(\cos \theta)$$

$$f_l = \frac{e^{2i\delta_l} - 1}{2i} = e^{i\delta_l} \frac{e^{i\delta_l} - e^{-i\delta_l}}{2i} = e^{i\delta_l} \sin \delta_l$$

$$\rightarrow \frac{1}{f_l} = \frac{1}{\sin \delta_l} e^{-i\delta_l} = \frac{1}{\sin \delta_l} (\cos \delta_l - i \sin \delta_l) = \cot \delta_l - i$$

$$\rightarrow f_l = \frac{1}{\cot \delta_l - i}$$

$$\cot \delta_l \Big|_{\delta_l = \frac{\pi}{2}} = 0 - \frac{1}{\sin^2 \delta_l} \Bigg|_{\delta_l = \frac{\pi}{2}} \left(\delta_l - \frac{\pi}{2} \right) + \dots \approx - \left(\delta_l - \frac{\pi}{2} \right)$$

Phase Shifts and Resonances - II

For E_R such that $\delta_l(E_R) = \frac{\pi}{2}$, expand into power series around E_R :

$$\delta_l(E) = \delta_l(E_R) + \left. \frac{d\delta_l}{dE} \right|_{E=E_R} (E - E_R) + \dots, \quad \frac{2}{\Gamma} \equiv \left. \frac{d\delta_l}{dE} \right|_{E=E_R} \rightarrow \delta_l \approx \frac{\pi}{2} + \frac{E - E_R}{\Gamma/2}$$

$$\rightarrow \cot \delta_l \underset{E \sim E_R}{\approx} - \left(\delta_l - \frac{\pi}{2} \right) = - \left(\frac{\pi}{2} + \frac{E - E_R}{\Gamma/2} - \frac{\pi}{2} \right) \approx - \frac{E - E_R}{\Gamma/2} = \frac{E_R - E}{\Gamma/2}$$

$$\rightarrow f_l \approx \frac{1}{\frac{(E_R - E)}{\Gamma/2} - i} = \frac{\Gamma/2}{E - E_R + i\Gamma/2} \quad \text{Breit-Wigner resonant amplitude}$$

E_R : characteristic energy of the system

$1/\Gamma$: Phase variation at $E_R \rightarrow [1/\Gamma] = \text{Time}$

Phase Shifts and Resonances - III

Partial cross-section for l wave:


$$\rightarrow |f_l|^2 = \sin^2 \delta_l = \frac{\Gamma^2/4}{(E - E_R)^2 + \Gamma^2/4},$$

Total cross-section = Sum of partial wave cross-sections

Often dominated by a resonance in one partial wave

Resonance 'symptoms':

- a) Fast increasing phase shift, going through $\pi/2$ at maximum rate
- b) $|f_l|^2$ strongly peaked
- c) Wave function large
- d) $d\delta/dk$, and delay, strongly peaked



Resonances - I

Generalize concept of stationary state:

$$\psi(\mathbf{r}, t) = \varphi(\mathbf{r}) e^{-iE_0 t} \rightarrow \int_{-\infty}^{+\infty} e^{-iE_0 t} e^{iEt} dt = \delta(E - E_0)$$

(Amplitude to find energy E when system is prepared in the state ψ)

to a kind of non-stationary, decaying state

$$e^{-iEt} = e^{-i(E_0 - i\Gamma)t} = e^{-iE_0 t} e^{-\Gamma t}, \quad t > 0$$

Complex E : Just meaning
"System is unstable"

$$\int_0^{+\infty} e^{-i(E_0 - i\Gamma)t} e^{iEt} dt = \int_0^{+\infty} e^{-i(E_0 - E - i\Gamma)t} dt = -\frac{1}{i(E_0 - E - i\Gamma)} e^{-i(E_0 - E - i\Gamma)t} \Big|_0^{+\infty} = \frac{i}{(E - E_0 + i\Gamma)}$$

(Breit-Wigner:

Amplitude to find energy E when system prepared in the state ψ)


$$|\psi|^2 \propto \left| \frac{i}{E - E_0 + i\Gamma} \right|^2 = \left| \frac{E - E_0 - i\Gamma}{(E - E_0)^2 + \Gamma^2} \right|^2 = \frac{(E - E_0 - i\Gamma)(E - E_0 + i\Gamma)}{\left[(E - E_0)^2 + \Gamma^2 \right]^2} = \frac{\left((E - E_0)^2 + \Gamma^2 \right)}{\left[(E - E_0)^2 + \Gamma^2 \right]^2} = \frac{1}{(E - E_0)^2 + \Gamma^2}$$

Resonances - II

Non-stationary levels may result from a particular shape of the effective potential

Non stationary, scattering state
But: *Almost* stationary...

Long lifetime, sharp quantum numbers: Like a *stable* state



$$\left. \begin{aligned} \Gamma &= 1/\text{time constant of decaying state} \approx \text{time uncertainty} \\ &\quad \text{Half width at half maximum} \approx \text{energy uncertainty} \end{aligned} \right\} \rightarrow \Delta E \Delta t \sim \Gamma \frac{1}{\Gamma} = 1$$

Δ Resonance Formation - I

Take πp scattering at low energy: use phase shift analysis
Some complication arising from spin 1/2

$k \sim m, r \leq R = \frac{1}{m} \rightarrow l = kr \leq 1$ Limited range, low energy: just 2 waves S and P

$$J = 1/2 \oplus 0 \oplus l = 1/2 \oplus l = \begin{cases} 1/2 & S-wave \\ 1/2, 3/2 & P-wave \end{cases}$$

Expand first incident wave:

$$e^{ikz} \underbrace{\chi_{1/2}^{+1/2}}_{\text{spin eigenstate}} = \frac{1}{2ikr} \sum_{l=0}^1 (2l+1) \left(e^{ikr} - (-1)^l e^{-ikr} \right) P_l(\cos \theta) \chi_{1/2}^{+1/2}$$

$$e^{ikz} \chi_{1/2}^{+1/2} = \frac{1}{2ikr} \sum_{l=0}^1 \sqrt{4\pi(2l+1)} \left(e^{ikr} - (-1)^l e^{-ikr} \right) Y_l^0(\cos \theta) \chi_{1/2}^{+1/2}$$

$$Y_l^0 \chi_{1/2}^{+1/2} = \sqrt{\frac{l+1}{2l+1}} y_{l+1/2}^{+1/2} - \sqrt{\frac{l}{2l+1}} y_{l-1/2}^{+1/2}$$

Spin spherical harmonics

$$y_{l+1/2}^{+1/2} = \sqrt{\frac{l+1}{2l+1}} Y_l^0 \chi_{1/2}^{+1/2} + \sqrt{\frac{l}{2l+1}} Y_l^1 \chi_{1/2}^{-1/2}, \quad y_{l-1/2}^{+1/2} = \sqrt{\frac{l+1}{2l+1}} Y_l^1 \chi_{1/2}^{-1/2} - \sqrt{\frac{l}{2l+1}} Y_l^0 \chi_{1/2}^{+1/2}$$

Δ Resonance Formation - II

$$\begin{aligned} & \frac{1}{2ikr} \sum_{l=0}^1 \sqrt{4\pi(2l+1)} \left(e^{ikr} - (-1)^l e^{-ikr} \right) Y_l^0(\cos \theta) \chi_{1/2}^{+1/2} \\ &= \frac{1}{2ikr} \sum_{l=0}^1 \sqrt{4\pi(2l+1)} \left(e^{ikr} - (-1)^l e^{-ikr} \right) \left(\sqrt{\frac{l+1}{2l+1}} y_{l+1/2}^{+1/2} - \sqrt{\frac{l}{2l+1}} y_{l-1/2}^{+1/2} \right) \\ &= \frac{1}{2ikr} \sum_{l=0}^1 \sqrt{4\pi} \left(e^{ikr} - (-1)^l e^{-ikr} \right) \left(\sqrt{l+1} y_{l+1/2}^{+1/2} - \sqrt{l} y_{l-1/2}^{+1/2} \right) \end{aligned}$$

Scattering amplitude: Phase shifts only modify *outgoing* spherical wave

$$\begin{aligned} f(\theta) &= \frac{1}{2ik} \sum_{l=0}^{\infty} (2l+1)(a_l - 1) P_l(\cos \theta) \\ \rightarrow f(\theta) &= \frac{\sqrt{4\pi}}{2ik} \sum_{l=0}^{\infty} \left(\sqrt{l+1} y_{l+1/2}^{+1/2} (a_l^+ - 1) - \sqrt{l} y_{l-1/2}^{+1/2} (a_l^- - 1) \right) \\ a_l^{\pm} &= e^{2i\delta_l^{\pm}} - 1 \end{aligned}$$

△ Resonance Formation - III

Re-arrange scattering amplitude:

$$f(\theta) = \frac{1}{2ik} \sum_{l=0}^1 \left[(l+1)(a_l^+ - 1) + l(a_l^- - 1) \right] P_l^0(\cos \theta) \chi_{1/2}^{+1/2} + (a_l^+ - a_l^-) P_l^{+1}(\cos \theta) e^{i\varphi} \chi_{1/2}^{-1/2}$$

$$= \frac{1}{2ik} \sum_{l=0}^1 \left[(l+1)(a_l^+ - 1) + l(a_l^- - 1) \right] P_l^0(\cos \theta) \chi_{1/2}^{+1/2} + \frac{1}{2ik} \underbrace{\sum_{l=0}^1 (a_l^+ - a_l^-) P_l^{+1}(\cos \theta) e^{i\varphi} \chi_{1/2}^{-1/2}}_{h(\theta)}$$

Spin non-flip amplitude Spin flip amplitude

Differential cross-section:

$$\frac{d\sigma}{d\Omega} = \frac{1}{k^2} |g(\theta)|^2 + |h(\theta)|^2 \quad g, h \text{ spin eigenfunctions orthogonal}$$

$P_0^0 = 1, P_1^0 = \cos \theta, P_1^{+1} = -\sin \theta$ Associate Legendre functions

$$\frac{d\sigma}{d\Omega} = \frac{1}{k^2} \left| \left(a_0^+ - 1 \right) + \left[2 \left(a_1^+ - 1 \right) + \left(a_1^- - 1 \right) \right] \cos \theta \right|^2 + \left| \left(a_1^+ - a_1^- \right) (-\sin \theta) \right|^2$$

$$\rightarrow \frac{d\sigma}{d\Omega} = \frac{1}{k^2} (A_0 + A_1 \cos \theta + A_2 \cos^2 \theta), \quad A_0, A_1, A_2 \text{ Energy dependent coefficients}$$

Δ Resonance Formation - IV

Around $\sqrt{s} \sim 1230$ MeV find $\frac{d\sigma}{d\Omega} = \frac{1}{k^2} (1 + 3\cos^2 \theta)$
consistent with the decay of a $J=3/2$ state

Indeed, taking for example $J_z = +1/2$:

$$|3/2, +1/2\rangle = \sqrt{\frac{1}{3}} |1/2, -1/2\rangle Y_1^{+1} + \sqrt{\frac{2}{3}} |1/2, +1/2\rangle Y_1^0$$

$$\frac{dN}{d\Omega} \propto \frac{1}{3} |Y_1^{+1}|^2 + \frac{2}{3} |Y_1^0|^2 = \frac{1}{3} \frac{1}{2} \sin^2 \theta + \frac{2}{3} \cos^2 \theta = \frac{1}{6} + \frac{3}{6} \cos^2 \theta \propto 1 + 3 \cos^2 \theta$$

Width:

ΔE = Breit-Wigner full width at half maximum ~ 100 MeV

$$\Delta t \sim 1/\Delta E = 1/100 \text{ MeV}^{-1}$$

$$\rightarrow \Delta t = 10^{-2} \cdot \hbar c \cdot 1/c = 10^{-2} \cdot 197 \text{ MeV fm} \cdot 1/(3 \times 10^{23} \text{ fm s}) \sim 0.7 \cdot 10^{-23} \text{ s}$$


Parity $\eta_\Delta = \eta_p \eta_\pi \eta_{orb} = (+1)(-1)(-1)^{l=1} = +1$

DNA Markers : Δ Angular Distributions


Experimental data nicely fit a simple picture where around $T_\pi = 200$ MeV the dominant amplitude is $J=3/2$, namely:

The large peak observed in the total cross-section can be traced back to a resonant amplitude in the $L=1$, $J=3/2$ partial wave

Several attempts to recover phase shifts from data in this energy range (Fermi, ...):
Messy game, lots of ambiguities



Δ^{++} : More Fingerprints



Cross-section

Phase

Production Resonances

With higher energy beams available,
new processes become possible.


Use *virtual pions* to excite nucleon
levels

$$p + p \rightarrow n + \Delta^{++} \rightarrow n + p + \pi^+$$



Resonance is *produced* in the *t*-channel, rather than *formed* in the *s*-channel.

Not directly observed in the cross-section vs. energy plot.

But: Resonance mass and quantum numbers are *invariant* properties, like the corresponding quantities of a stable particle





The Bubble Chamber





CERN 2m Bubble Chamber

Bubble Chamber Events - I



Bubble Chamber Events - II



Knock-on electron
Beam + Atomic Electron
 \rightarrow *Beam + Free Electron*
 Usually modest energy

$\pi\mu e$ kinematics

π^+ only: π^- is usually captured to a π -mesic atom
 π decays after stopping: 'long' lifetime..
 μ Energy, momentum:

$$E_\mu = \frac{1}{2m_\pi} (m_\pi^2 + m_\mu^2 - 0) \sim 109.9 \text{ MeV} \rightarrow p_\mu = \sqrt{109.9^2 - 106^2} \sim 29.1 \text{ MeV}$$

$$\rightarrow \beta_\mu = \frac{p_\mu}{E_\mu} \sim \frac{29.1}{109.9} \sim 0.265, \gamma_\mu \sim 1.04 \text{ when created}$$


Would expect typical path length $\sim \beta_\mu \gamma_\mu c \tau_\mu \sim 182 \text{ m}$

But: μ quickly slows down by $\frac{dE}{dx} \rightarrow$ Total path length \sim few cm

Positron spiralling down: Energy loss by $\begin{cases} \text{ionization} \\ \text{radiation} \end{cases}$

Magnetic Analysis & Accuracy

Motion of a charged particle in a magnetic field: Cylindrical helix coaxial to \mathbf{B}



$$r = \frac{p_{\perp}}{0.3B} \quad r: m, p_{\perp}: GeV, \quad B: T$$

Get p from s

$$\sin \frac{\theta}{2} = \frac{L}{2r} \xrightarrow{L \ll 2r} \frac{\theta}{2} \approx \frac{L}{2r} \rightarrow \theta \approx \frac{0.3BL}{p_{\perp}}$$

$$s = r - r \cos \frac{\theta}{2} \approx r \left[1 - \left(1 - \frac{\theta^2}{4} \right) \right] = r \frac{\theta^2}{8} \approx \frac{0.3BL^2}{8p_{\perp}}$$

$$\rightarrow p_{\perp} \approx \frac{0.3BL^2}{8s}$$

Take 3 measured points, with single point accuracy σ

Then:


$$s = x_B - \frac{x_A + x_B}{2} \rightarrow \sigma_s^2 = \sigma^2 + \frac{1}{2}\sigma^2 = \frac{3}{2}\sigma^2$$

$$\frac{\sigma_{p_{\perp}}}{p_{\perp}} = \frac{\sigma_s}{s} = \sqrt{\frac{3}{2}} \frac{\sigma}{s} = \sqrt{\frac{3}{2}} \frac{\sigma 8p_{\perp}}{0.3BL^2} = \sqrt{\frac{300 \cdot 64}{18}} \frac{\sigma p_{\perp}}{BL^2} \approx 32.7 \frac{\sigma p_{\perp}}{BL^2}$$

$N \geq 10$, uniformly spaced points:

$$\frac{\sigma_{p_{\perp}}}{p_{\perp}} \approx 28.3 \frac{\sigma p_{\perp}}{BL^2 \sqrt{N+4}}$$

Bubble Chamber Reconstruction



Particle	\mathbf{p}_x	\mathbf{p}_y	\mathbf{p}_z	E
K-	8213.4	-248.3	15.2	8232
p	0	0	0	938.3
Sum	8213.4	-248.3	15.2	9170.3
K-	1481.8	27.8	224	1578.1
p-	149.7	-11.3	38.8	208.6
p+	37.9	-122.2	-22.7	190.7
p	1508.6	128.5	-70.5	1782.6
K0	3545.6	-162.9	-245	3592.4
Sum	6723.6	-140.1	-75.4	7352.4
Difference	-1489.8	108.2	-90.6	-1817.9


mass 1032.153

This mass doesn't correspond to a known particle - so there must be at least two neutral particles from the collision leaving the bubble chamber undetected.

Δ -Resonance: Production

Observe Δ^{++} resonance production as a peak in the invariant (p, π^+) mass distribution


Take reaction



$$m_{p\pi_1}^2 = (p_p + p_{\pi_1})^2 = (E_p + E_{\pi_1})^2 - (\mathbf{p}_p + \mathbf{p}_{\pi_1})^2$$

$$m_{p\pi_2}^2 = (p_p + p_{\pi_2})^2 = (E_p + E_{\pi_2})^2 - (\mathbf{p}_p + \mathbf{p}_{\pi_2})^2$$

2 entries per reconstructed event:
Count everything



Meson Resonances - I

Expect resonant behavior also for mesonic systems, e.g. $\pi\pi$:
Virtual and real pion coupled at the strong vertex

Observation of meson resonances possible in production experiments

Remark:

Taking baryon resonances only, possible isospin:


Minimum coupling is between nucleon and pion

→ Expect $1 \oplus 1/2 = 1/2, 3/2$ as observed

Take meson resonances:


Minimum coupling is between pion and pion

→ Expect $1 \oplus 1 = 0, 1, 2$ I=2 mesons not observed



Meson Resonances - II

Take reaction



Observe strong enhancements for

$$m_{\pi\pi} \sim 760, 1260, 1550 \text{ MeV}$$

$$m_{\pi n} \sim 1230 - 1550 \text{ MeV}$$

Interpretation:

Meson	Baryon	Resonances
$\rho(760)$		
$f_0(1250)$	$\rightarrow \pi^\pm \pi^\mp$	$\Delta^{+-}(1232) \rightarrow n \pi^\pm$
$g(1550)$		




FIG. 2. Scatter plot of $M(\pi^+ \pi^-)$ versus $M(\pi^+ n)$ with the projections on both axes.

Spin-parity of the ρ Meson - I

Use angular distributions to investigate ρ spin, parity

$$S_\pi = 0 \rightarrow J_\rho = L_{\pi\pi} \equiv L$$

$$\rightarrow \psi_{final} \propto Y_l^m(\theta, \varphi)$$

$$\eta_P^{(\rho)} = \eta_P^{(\pi)} \eta_P^{(\pi)} (-1)^l = (-1)^l$$

Suppose the produced ρ mesons uniformly populate the $2l+1$ J_3 substates: Then, by a property of spherical harmonics

$$\frac{dP}{d\Omega} = \frac{1}{2J+1} \sum_{m=-l}^{+l} Y_l^m(\theta, \varphi) Y_l^{*m}(\theta, \varphi); \quad \sum_{m=-l}^{+l} Y_l^m Y_l^{*m} = \frac{2l+1}{4\pi}$$
$$\rightarrow \frac{dP}{d\Omega} = \frac{1}{2J+1} \frac{2J+1}{4\pi} = \frac{1}{4\pi} \text{ Uniform distribution}$$

So a non-uniform angular distribution indicates some *polarization* of the decaying state, useful to perform spin-parity analysis

Spin-parity of the ρ Meson - II

Observe CM angular distribution
for different $\pi\pi$ mass 'slices'


In the ρ resonance mass region
(about 700-800 MeV)

$$\frac{dP}{d\Omega} \propto \cos^2 \theta \propto |Y_1^0(\cos \theta)|^2 \rightarrow l = 1$$

→ The ρ is a *vector* particle

Interestingly, in the f_0 mass region
(about 1250-1350 MeV) observe
some indication of spin 2

$$\frac{dP}{d\Omega} \propto (3\cos^2 \theta - 1)^2 \propto |Y_2^0|^2 \rightarrow l = 2$$



Isospin - I

Charge independence leads to a new classification scheme:

All hadrons cast into *isospin multiplets*

Strong interaction identical for all members of each multiplet

proton p
neutron n

$\left. \begin{array}{l} p \\ n \end{array} \right\}$ 2 states of the *nucleon* $N = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ 2 states system - isospinor

Base $\begin{pmatrix} 1 \\ 0 \end{pmatrix} \equiv p, \quad \begin{pmatrix} 0 \\ 1 \end{pmatrix} \equiv n$

Base states: *doublet*

π^+
 π^0
 π^-

$\left. \begin{array}{l} \pi^+ \\ \pi^0 \\ \pi^- \end{array} \right\}$ 3 states of the *pion* $\pi = \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}$ 3 state system - isovector

Base $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \equiv \pi^+, \quad \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \equiv \pi^0, \quad \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \equiv \pi^-$

Base states: *triplet*

Isospin - II

Isospins add up as angular momenta (Astonished? More on this later...)

For πN system obtain:

$$\left. \begin{array}{l} \pi : I = 1 \\ N : I = 1/2 \end{array} \right\} \rightarrow \pi N : I = 1 \oplus 1/2 = \begin{cases} 1/2 & \text{doublet} \\ 3/2 & \text{quadruplet} \end{cases}$$

By using Clebsch-Gordan coefficients, expand physical (particle) states into total isospin eigenstates

Single particle: Base states

$$I_N = 1/2 ; |p\rangle = |1/2, +1/2\rangle , |n\rangle = |1/2, -1/2\rangle$$

$$I_\pi = 1 ; |\pi^+\rangle = |1, +1\rangle , |\pi^0\rangle = |1, 0\rangle , |\pi^-\rangle = |1, -1\rangle$$

Isospin - III

Expand physical, 2 particle states into total isospin eigenstates:

$$|\pi^- p\rangle = |1, -1, 1/2, +1/2\rangle = \sqrt{\frac{1}{3}} |3/2, -1/2\rangle - \sqrt{\frac{2}{3}} |1/2, -1/2\rangle$$

$$|\pi^+ n\rangle = |1, +1, 1/2, -1/2\rangle = \sqrt{\frac{1}{3}} |3/2, +1/2\rangle + \sqrt{\frac{2}{3}} |1/2, +1/2\rangle$$

$$|\pi^+ p\rangle = |1, +1, 1/2, +1/2\rangle = |3/2, +3/2\rangle$$

$$|\pi^- n\rangle = |1, -1, 1/2, -1/2\rangle = |3/2, -3/2\rangle$$

$$|\pi^0 p\rangle = |1, 0, 1/2, +1/2\rangle = \sqrt{\frac{2}{3}} |3/2, +1/2\rangle - \sqrt{\frac{1}{3}} |1/2, +1/2\rangle$$

$$|\pi^0 n\rangle = |1, 0, 1/2, -1/2\rangle = \sqrt{\frac{2}{3}} |3/2, -1/2\rangle + \sqrt{\frac{1}{3}} |1/2, -1/2\rangle$$

Isospin - IV

Guess isospin is a new *symmetry* for hadrons: connect to some *invariance* property (like angular momentum).

Non-trivial conservation rule follows:

Total isospin conserved by all strong processes

Interesting predictions for πN scattering and reactions:

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- n \rightarrow \pi^- n \end{cases} \rightarrow A_A = A_B = A_{3/2} \quad \text{pure } I = 3/2$$

$$\begin{cases} (A) \pi^+ n \rightarrow \pi^+ n \\ (B) \pi^- n \rightarrow \pi^- n \end{cases} \rightarrow A_A = \frac{1}{3} A_{3/2} + \frac{2}{3} A_{1/2}, A_B = A_{3/2}$$

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- p \rightarrow \pi^- p \end{cases} \rightarrow A_A = A_{3/2}, A_B = \frac{1}{3} A_{3/2} - \frac{2}{3} A_{1/2}$$

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- p \rightarrow \pi^0 n \end{cases} \rightarrow A_A = A_{3/2}, A_B = \sqrt{\frac{2}{9}} A_{3/2} - \sqrt{\frac{2}{9}} A_{1/2}$$

Isospin - V

If $A_{3/2} \gg A_{1/2}$

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- n \rightarrow \pi^- n \end{cases} \rightarrow \sigma_A = \sigma_B$$

$$\begin{cases} (A) \pi^+ n \rightarrow \pi^+ n \\ (B) \pi^- n \rightarrow \pi^- n \end{cases} \rightarrow \sigma_A \simeq \frac{1}{9} \sigma_B$$

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- p \rightarrow \pi^- p \end{cases} \rightarrow \sigma_A \simeq 9\sigma_B$$

$$\begin{cases} (A) \pi^+ p \rightarrow \pi^+ p \\ (B) \pi^- p \rightarrow \pi^0 n \end{cases} \rightarrow \sigma_A \simeq \frac{9}{2} \sigma_B$$

Still lacking: *What exactly is isospin?*

What is Spin? - I

For any physical system with $m>0$, we are allowed to choose CM as a reference frame.

When the system is rotationally invariant, states are observed to group into multiplets of size n , $n=1,2,3,\dots$ (*size n = number of states*)

States of a multiplet: *Same energy*

States belonging to different multiplets must be distinguished by some internal quantum number: Provisionally call the corresponding observable the particle spin

States of any given multiplet must be identified by some internal quantum number: Provisionally call the corresponding observable the 3rd component of the particle spin

What is Spin? - II

Question: *What is the observable we have called spin?*

Answer: *Get some insight from conservation laws.*

Discover spin is just another kind of (non-orbital) angular momentum

$$\mathbf{J} = \mathbf{L} + \mathbf{S} \quad \text{Total angular momentum}$$

For any system: Extend to S known properties of \mathbf{L}

- (S_x, S_y, S_z) analogue to (L_x, L_y, L_z) : Hermitian operators, infinitesimal generators of rotations around x, y, z
- Commutators: $[S_x, S_y] = iS_z$ + Cyclical permutations
- By assuming rotational invariance, in the CM H and S^2 commute
 $\rightarrow S^2, S_3$ are conserved

What is Spin? - III

- Besides other quantum numbers, in the CM reference frame all possible stationary states are then labeled by S^2, S_z according to angular momentum algebra:

S^2 Eigenvalues: $s(s+1), s = 0, 1/2, 1, 3/2, 2, \dots$

Sequence of multiplets

S_z Eigenvalues: $\underbrace{-s \dots + s}_{2j+1}, 2s+1 \equiv$ Multiplet size $1, 2, 3, 4, 5, \dots$

- Each multiplet understood to realize an *irreducible representation* of some (unknown) symmetry group in the Hilbert space

NB Multiplets with even multiplicity are observed $\rightarrow 2j+1 = 2, 4, \dots$
Implies j can be *integer* or *half-integer*

What is Spin? - IV

Representation:

A set of matrices acting on some kind of 'vectors', labeled by the integer $2s+1$

- Must have 3 independent matrices ($= S_x, S_y, S_z$) for each rep.
- Must have $2j+1$ independent 'vectors' ($= \text{base states}$) for each rep

Size of matrices: $(2s+1) \times (2s+1)$

Each matrix correspond to a *specific rotation*

- Must depend on 3 parameters ($= \text{rotation angles}$)

What is Spin? - V

Integer s : Like l

- L eigenvalues are integer only $0, 1, 2, \dots \rightarrow 2l+1 = 1, 3, 5, \dots$ odd integer
- l identifies an *irreducible representation* of the rotation group $SO(3)$
- (L_x, L_y, L_z) : 3 matrices of size $1x1, 3x3, 5x5, \dots$ operating on different objects of size $1, 3, 5, \dots$: *Spherical Tensors* (e.g. Spherical Harmonics)

Half-integer s : Minimum size is for $s=1/2 \rightarrow 2 \times 2$

- 2-component 'vectors' acted upon by 2×2 matrices called *spinors*
Not really like ordinary vectors..

From the algebraic properties of \mathbf{S} :

Spin symmetry group must be a close relative of $SO(3)$

Just including extra values for s as compared to l

Matrix Fun - I

Take $j=1/2$:

Must represent rotations of 2-component spinors by 2×2 matrices

1) Naive attempt: Try with orthogonal matrices

$$M \equiv \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ orthogonal}$$

$$\rightarrow MM^T = 1$$

$$\rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\rightarrow \begin{cases} a^2 + b^2 = 1 \\ c^2 + d^2 = 1 \quad \& a, b, c, d \text{ real} \\ ac + bd = 0 \end{cases}$$

\rightarrow 1 free parameter

\rightarrow KO to represent a 3D rotation

Matrix Fun - II

2) Better approach: Unitary matrices

$$U \equiv \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ unitary} \rightarrow UU^\dagger = 1 \rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \bar{a} & \bar{c} \\ \bar{b} & \bar{d} \end{pmatrix} = \begin{pmatrix} a\bar{a} + b\bar{b} & a\bar{c} + b\bar{d} \\ c\bar{a} + d\bar{b} & c\bar{c} + d\bar{d} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\rightarrow \begin{cases} a\bar{a} + b\bar{b} = 1 \\ c\bar{c} + d\bar{d} = 1 \quad \& a, b, c, d \text{ complex} \rightarrow 4 \text{ free parameters} \\ a\bar{c} + b\bar{d} = 0 \\ c\bar{a} + d\bar{b} = 0 \end{cases}$$

Require extra condition:

$\det M = 1 \rightarrow ad - bc = 1 \rightarrow 3 \text{ free parameters}$

\rightarrow OK to represent a 3D rotation

Possible because absolute phase of states is irrelevant

Matrix Fun - III

Set of all 2×2 matrices satisfying the 4 conditions above:

A group, called the *Special Unitary group of dimension 2*, or $SU(2)$.

$SU(2)$ vs $SO(3)$:

3 parameters \rightarrow 3 generators

Commutators identical \rightarrow They share the same *algebra*

The moral:

$O(3)$ and $SU(2)$ are *more or less* the same group

\rightarrow All the irr.reps of $SO(3)$ are also good for $SU(2)$

$SU(2)$ - I

Instead of starting from rotations, just start from $SU(2)$ defined as the set of all the 2×2 , unitary matrices (with $\det=1$)

Not bound to understand this transformation of states as induced by a rotation of axis in the physical, 3D space.

Free to interpret any $SU(2)$ matrix as representing a unitary, unimodular transformation in the Hilbert space of any two-state, degenerate system.

Do not need to specify what is the physical system whose two independent states we take as base vectors in the Hilbert space.

$SU(2)$ - II

Some matrix fun:

4 complex parameters \rightarrow 8 real parameters

4 unitarity conditions: $\left. \begin{array}{l} UU^\dagger = 1 \\ (U^\dagger)_{ij} = U_{ji}^* \end{array} \right\} \rightarrow \sum_{j=1}^2 a_{ij} a_{jk}^* = \delta_{ik}, \quad i, k = 1, 2$

1 unimodularity condition: $\det U = 1$

$\rightarrow 8 - 5 = 3$ free parameters

One diagonal generator, σ_3

\rightarrow Rank 1 group

\rightarrow One invariant function of generators

Quadratic: $\sigma^2 = \sigma_1^2 + \sigma_2^2 + \sigma_3^2$

$SU(2)$ - III

Some insight into $SU(2)$ generators:

U unitary $\rightarrow U = e^{iH}$, H Hermitian

$$\det U = 1 \rightarrow \det e^{iH} = 1 \rightarrow e^{itr(H)} = 1 \rightarrow tr(H) = 0$$

3 free parameters \rightarrow 3 generators

3 Hermitian, traceless 2×2 matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Any $SU(2)$ matrix can be written as a linear combination of the 3 generators, the *Pauli matrices*

What is Isospin? - I

When looking at strongly interacting particles, observe particle states similarly grouping themselves into multiplets of size 1,2,3,4

States of a multiplet \simeq Same mass

→ States belonging to different multiplets must be distinguished by some internal quantum number: By analogy, call the corresponding observable the particle *isospin*

→ States of any given multiplet must be identified by some *internal* quantum number: Call the corresponding observable the *3rd component of the particle isospin*

What is Isospin? - II

Notice: Isospin symmetry is not exact (broken), still is quite good
Indeed, looking at symmetry breaking mass splittings:

$$\frac{m_n - m_p}{m_n} \simeq \frac{939.57 - 938.27}{939.57} \simeq 0.0014 \text{ Nucleon doublet}$$

$$\frac{m_{\pi^\pm} - m_{\pi^0}}{m_{\pi^\pm}} \simeq \frac{139.6 - 135.0}{139.6} \simeq 0.011 \text{ Pion triplet}$$

For a long time: Breaking entirely blamed on electromagnetic effects, which is only partially true (e.g. neutral and charged members indeed have quite different e.m. interactions contributing to their mass).

Today: Isospin taken as an 'accidental' symmetry, not due to some fundamental property of hadron constituents or strong interaction

What is Isospin? - III

Question: What is the observable we have called *isospin*?

Answer: *There is no classical analogy!*

Simply, as we observe the neutron and proton to be almost degenerate in mass, we can state they are just two states of the same physical system, the *nucleon*.

We guess the two nucleon states are the ‘vectors’ spanning the fundamental representation of a symmetry group, which we identify with $SU(2)$.

What is Isospin? - IV

Guess: $SU(2)$ is a symmetry of all the strongly interacting particles.
Therefore:

All strong interacting particles should fill some $SU(2)$ representation

This is actually true, after neglecting small symmetry breaking effects within each multiplet (see later)

As for any other symmetry, expect the invariance property to yield a conservation law

What is Isospin? - V

What is conserved in this case?

Since there is no classical analogy, stick to our algebraic skills to get insight

$SU(2)$ algebra is just the same as $O(3)$, so we can expect the same conserved observables for a closed system of strongly interacting particles:

$$\mathbf{J}^2, J_3 \leftrightarrow \mathbf{I}^2, I_3$$

This is the origin of the common wisdom '*Isospin is like Angular Momentum*'

$SU(2)$ Multiplet Graphics

Within any given $SU(2)$ multiplet, states can be represented as points on a straight line

Reason is the group structure of $SU(2)$:


3 parameters $\rightarrow 3$ generators

*Just 1 invariant function of generators:
 $I^2 \rightarrow$ Multiplets identified just by I*

*Generators do not commute with each other
 \rightarrow States in any multiplet identified just by I_3*

Define 2 ladder operators: $I_{\pm} = I_1 \pm iI_2$

Action: Shift states right or left on the multiplet line, i.e. increment/decrement I_3 by 1



Observe:
 I_3 eigenvalues symmetric wrt 0

Conjugate Representation - I

More fun with matrices...

D : Any representation

$$\psi' = D(\alpha)\psi$$

$\rightarrow D(\alpha) = e^{i\alpha F}$, F hermitian \leftarrow True because D is unitary

Take complex conjugate of equations

$$\psi^* = D^*\psi$$

Get another representation

$$D^* = e^{-i\alpha(F)^*} = e^{i\alpha[-(F)^*]} \equiv e^{i\alpha\tilde{F}}$$

Relation bewteen new and old generators

$$\rightarrow \tilde{F} = -(F^*)$$

Conjugate Representation - II

Take D of $SU(2)$ fundamental representation:

F Hermitian $\rightarrow \tilde{F}$ Hermitian

\rightarrow Real eigenvalues for both F, \tilde{F} , and $f_i = -f_i^*$

\rightarrow Since f_i are symmetric wrt 0, so are f_i^*

$\rightarrow \{f_i\} \equiv \{f_i^*\}$ \tilde{F} eigenvalues are just a re-labeling of F 's

Direct and conjugate representations are said to be *equivalent*

True for $SU(2)$, generally false

Product of Representations - I

Take a system made of 2 nucleons: *What is the total isospin?*
 $SU(2)$ is equivalent to $O(3) \rightarrow$ Can use Clebsch-Gordan coefficients

But: Can also re-formulate the problem in a different way
Each nucleon spans the fundamental representation of $SU(2)$, **2**

Then a 2 nucleon system span the *direct product rep.* **2** \otimes **2**

Question:

What are the irreducible representations of $SU(2)$ contained in any state of 2 nucleons?

Need to decompose **2** \otimes **2** into a *direct sum* of irr.rep.


Product of Representations - II

Answer (After a little group theory):

$$\mathbf{2} \otimes \mathbf{2} = \mathbf{1} \oplus \mathbf{3}$$

Answer (Graphical):

*Center the segment carrying the 2 states of representation **2** (1st nucleon) over the 2 states of representation **2** (2nd nucleon)*
→ Get a set of 4 states, decomposing into 2 sets of 1 and 3 states



I -Spin Multiplets: The Nonstrange Zoo

Amazingly large number of resonant states

p, n	P_{11}	****	$\Delta(1232)$	P_{33}	****
$N(1440)$	P_{11}	****	$\Delta(1600)$	P_{33}	***
$N(1520)$	D_{13}	****	$\Delta(1620)$	S_{31}	****
$N(1535)$	S_{11}	****	$\Delta(1700)$	D_{33}	****
$N(1650)$	S_{11}	****	$\Delta(1750)$	P_{31}	*
$N(1675)$	D_{15}	****	$\Delta(1900)$	S_{31}	**
$N(1680)$	F_{15}	****	$\Delta(1905)$	F_{35}	****
$N(1700)$	D_{13}	***	$\Delta(1910)$	P_{31}	****
$N(1710)$	P_{11}	***	$\Delta(1920)$	P_{33}	***
$N(1720)$	P_{13}	****	$\Delta(1930)$	D_{35}	***
$N(1900)$	P_{13}	**	$\Delta(1940)$	D_{33}	*
$N(1990)$	F_{17}	**	$\Delta(1950)$	F_{37}	****
$N(2000)$	F_{15}	**	$\Delta(2000)$	F_{35}	**
$N(2080)$	D_{13}	**	$\Delta(2150)$	S_{31}	*
$N(2090)$	S_{11}	*	$\Delta(2200)$	G_{37}	*
$N(2100)$	P_{11}	*	$\Delta(2300)$	H_{39}	**
$N(2190)$	G_{17}	****	$\Delta(2350)$	D_{35}	*
$N(2200)$	D_{15}	**	$\Delta(2390)$	F_{37}	*
$N(2220)$	H_{19}	****	$\Delta(2400)$	G_{39}	**
$N(2250)$	G_{19}	****	$\Delta(2420)$	$H_{3,11}$	****
$N(2600)$	$I_{1,11}$	***	$\Delta(2750)$	$I_{3,13}$	**
$N(2700)$	$K_{1,13}$	**	$\Delta(2950)$	$K_{3,15}$	**

Baryons
 $I=1/2$ $I=3/2$

$L_{2J+1,2I+1}$ $L = S, P, D, \dots$

LIGHT FLAVORED $\mathcal{J}^P = B = 0$	
$J^P(J^{PC})$	$J^P(J^{PC})$
• π^\pm	$1^-(0^-)$
• π^0	$1^-(0^-+)$
• η	$0^+(0^-+)$
• $f_0(400-1200)$	$0^+(0^++)$
• $\rho(770)$	$1^+(1^-)$
• $\omega(782)$	$0^-(1^-)$
• $\eta'(958)$	$0^+(0^-+)$
• $f_0(980)$	$0^+(0^++)$
• $a_0(980)$	$1^-(0^++)$
• $\phi(1020)$	$0^-(1^-)$
• $b_1(1170)$	$0^-(1^-+)$
• $b_1(1235)$	$1^+(1^-+)$
• $a_1(1260)$	$1^-(1^-+)$
• $f_2(1270)$	$0^+(2^++)$
• $f_1(1285)$	$0^+(1^-+)$
• $\eta(1295)$	$0^+(0^-+)$
• $\pi(1300)$	$1^-(0^-+)$
• $a_2(1320)$	$1^-(2^++)$
• $f_0(1370)$	$0^+(0^-+)$
• $b_1(1380)$	$?^-(1^-+)$
• $\pi_1(1400)$	$1^-(1^-+)$
• $f_1(1420)$	$0^+(1^-+)$
• $\omega(1420)$	$0^-(1^-)$
• $f_2(1430)$	$0^+(2^++)$
• $\eta(1440)$	$0^+(0^-+)$
• $a_0(1450)$	$1^-(0^-+)$
• $\rho(1450)$	$1^+(1^-)$
• $f_0(1500)$	$0^+(0^-+)$
• $f_1(1510)$	$0^+(1^-+)$
• $f_2'(1525)$	$0^+(2^++)$
• $f_2(1565)$	$0^+(2^++)$
• $\pi_1(1600)$	$1^-(1^-+)$
• $X(1600)$	$2^+(?^-)$
• $a_1(1640)$	$1^+(1^-+)$
• $f_2(1640)$	$0^+(2^++)$
• $\eta_2(1645)$	$0^+(2^-+)$
• $\omega(1650)$	$0^-(1^-)$
• $X(1650)$	$0^-(?^-)$
• $a_2(1660)$	$1^-(2^++)$
• $\omega_3(1670)$	$0^-(3^-)$


I=2 ???

Mesons
 $I=0,1$

Baryon Resonances Systematics

Two families of nucleon excited states: First, lightest states


N^* isospin = 1/2




Δ isospin = 3/2

Many sub-families for each one (increasing J , parity + or -)


Non-strange Baryons – $I = 1/2$




Non-strange Baryons – $I=3/2$



Non-Strange Mesons – $I=0$



Non-Strange Mesons – $I=1$



Gell-Mann – Nishijima Rule

B = Baryon number

Q = Charge in e units

I_3 = Isospin 3rd component

Empirical relationship for pions:

$$Q = I_3$$

Linking electromagnetic and strong properties of pions:

Electric charge as *3rd component* of isospin vector

Extend to nucleons:

$$Q = I_3 + B/2 \quad \text{Gell-Mann - Nishijima relation}$$

More complicated properties:

Electric charge as both *scalar* and *3rd component*

Strangeness - I

Strange particles discovered in cosmic rays at the end of the '40s, and then quickly observed at the first GeV accelerators
Why strange?

Large production cross section → Like ordinary hadrons
Long lifetime → Like weak decays

Understood as carriers of a new quantum number: *Strangeness*

Ordinary hadrons $S = 0$

Strange particles $S \neq 0$

Strangeness conserved by strong, e.m. processes, violated by weak

Explain funny behavior, also predicting *associated production* to guarantee S conservation in strong & EM processes:

Strange particles always produced in pairs

Strangeness - II

For strong processes, S similar to electric charge and to baryon or lepton numbers

But:

S not absolutely conserved
 S not the source of a physical field

Large variety of strange particles, both baryons and mesons, including many strange resonances

Generalize Gell-Mann Nishijima relation to

$$Q = I_3 + \frac{B+S}{2} = I_3 + \frac{Y}{2}$$

Hypercharge

The Lightest Strange Particles

I_3	$S=+1$	$S=-1$
+1/2	K^+	K^0
-1/2	\bar{K}^0	K^-

Spin 0

I_3	$S=+1$	$S=-1$
+1/2	K^{*+}	\bar{K}^{*0}
-1/2	K^{*0}	K^{*-}

Spin 1

I_3	S	name
0	-1	Λ^0
+1,0,-1	-1	$\Sigma^+, \Sigma^-, \Sigma^0$
+1/2,-1/2	-2	Ξ^0, Ξ^-
0	-3	Ω^-

Baryons

I_3	S	name
0	+1	$\bar{\Lambda}^0$
+1,0,-1	+1	$\bar{\Sigma}^+, \bar{\Sigma}^0, \bar{\Sigma}^-$
+1/2,-1/2	+2	$\bar{\Xi}^0, \bar{\Xi}^-$
0	+3	$\bar{\Omega}^-$

Antibaryons

Isospin of Strange Particles

Isospin conservation in

$$\pi^- + p \rightarrow \pi^- + p$$

leads in a natural way to extend to virtual states like

$$\pi^- + p \rightarrow (K^0 + \Lambda^0)^* \rightarrow \pi^- + p$$

Therefore strange particles should group into I-spin multiplets.


Λ^0 only observed as a neutral state \rightarrow Singlet , $I = 0$

Observe 3 charge states for K: Triplet?

$$\pi^- + p : I = 1/2, 3/2 \rightarrow K \text{ must be } I = 1/2, 3/2$$

Quartets not observed \rightarrow 2 Doublets! Predict two neutral K states, with opposite S

Would imply charge +2



Must be different particles!

Bubble Chambers & Particle Zoology

Example: Historical Picture

$$K^- + p \rightarrow K^0 + K^+ + \Omega^-$$

$$K^0 \rightarrow \pi^+ + \pi^-$$

$$K^+ \rightarrow \pi^+ + \pi^0 \text{ (unseen)}$$


$$\Omega^- \rightarrow \Lambda^0 + K^-$$

$$\Lambda^0 \rightarrow p + \pi^-$$

$$K^- \rightarrow \pi^- + \pi^0 \text{ (unseen)}$$

Beam momentum 4.2 GeV

Magnetic field 2 T



Old Hyperon Beam & Spectrometer

FNAL - '70s Beam & Detector of Hyperon Experiment




Fig. 1



Figure 1
The Hyperon Magnet under construction

Hyperon Gymnastics

Old Hyperon Beam & Spectrometer


Reconstruct decays: $\Sigma^- \rightarrow n + \pi^-$, $\Xi^- \rightarrow \Lambda^0 + \pi^-$

π : Identification (Threshold Cherenkov) + Magnetic Analysis

n : Calorimeter

p : Identification (Cherenkov π Veto) + Magnetic Analysis + Calorimeter

$\Lambda^0 \rightarrow p + \pi$: Identification + Magnetic Analysis



Particle Identification: Cherenkov - I

Fast, charged particle passing through a dielectric medium

Cherenkov radiation emitted for $\beta > \frac{1}{n}$, n refractive index

Main features:

Emission angle:

$$\cos \theta_c = \frac{1}{\beta n} \quad \text{Cherenkov angle}$$

For ultrarelativistic particles:

$$\lim_{\beta \rightarrow 1} (\cos \theta_c) = \frac{1}{n} \quad \text{Asymptotic angle}$$

Spectrum:

Medium	n	θ_{min} deg	$P_{thresh}(\pi)$ GeV	N_{ph} eV $^{-1}$ cm $^{-1}$
Air	1.00028	1.36	5.9	0.21
Isobutane	1.00217	3.77	2.12	0.94
Aerogel	1.0065	6.51	1.3	4.7
Water	1.33	41.2	0.16	160.8
Quartz	1.46	46.7	0.13	196.4

1/ λ^2 spectrum: Blue/Near UV very important...

$$\frac{d^2N}{dx d\lambda} = 2\pi\alpha z^2 \frac{1}{\lambda^2} \sin^2 \theta_c \quad \text{photons/cm}^2, \quad z \text{ particle charge in } e \text{ units}$$

$$\frac{d^2N}{dx dE} = \frac{\alpha}{\hbar c} z^2 \sin^2 \theta_c \approx 365 z^2 \sin^2 \theta_c \quad \text{photons}/(cm \cdot eV)$$

Number of photons/cm small...

Particle Identification: Cherenkov - II

Translate light signal into an electric charge: *Photomultiplier*, or similar
Typical result with a PM:

$$N_{pe} \approx 365 \int_{E_{\min}}^{E_{\max}} \varepsilon_{coll}(E) \varepsilon_{det}(E) \sin^2 \theta_c(E) dE$$

N. of photoelectrons/cm obtained

Collection efficiency

Conversion efficiency

Cherenkov angle depending on E :

$$\cos \theta_c = \frac{1}{\beta n(\lambda)} = \frac{1}{\beta n(E)}$$

Dispersion of refractive index

Typically:

$$N_{pe} \leq 100 \sin^2 \theta_c \quad \text{Photoelectrons/cm}$$

Threshold counter

$$\begin{aligned} \beta > \frac{1}{n} \rightarrow \frac{p}{E} > \frac{1}{n} \rightarrow \frac{p}{\sqrt{p^2 + m^2}} > \frac{1}{n} \rightarrow p^2 > \frac{1}{n^2} (p^2 + m^2) \\ \rightarrow p^2 \left(1 - \frac{1}{n^2}\right) > \frac{m^2}{n^2} \rightarrow p^2 > \frac{m^2}{n^2 - 1} \rightarrow p > \frac{m}{\sqrt{n^2 - 1}} \end{aligned}$$

Threshold momentum

Can discriminate among different masses with the same momentum

The Strange Zoo

		Λ	P_{01}	****	Ξ^0	P_{11}	****	Σ^+	P_{11}	****			$\bullet K^\pm$	1/2(0 ⁻)	
$\Lambda(1405)$		S_{01}	****	Ξ^-		P_{11}	****	Σ^0	P_{11}	****			$\bullet K^0$	1/2(0 ⁻)	
$\Lambda(1520)$		D_{03}	****	$\Xi(1530)$		P_{13}	****	Σ^-	P_{11}	****			$\bullet K_S^0$	1/2(0 ⁻)	
$\Lambda(1600)$		P_{01}	***	$\Xi(1620)$			*	$\Sigma(1385)$	P_{13}	****			$\bullet K_L^0$	1/2(0 ⁻)	
$\Lambda(1670)$		S_{01}	****	$\Xi(1690)$			***	$\Sigma(1480)$		*			$K_0^*(800)$	1/2(0 ⁺)	
$\Lambda(1690)$		D_{03}	****	$\Xi(1820)$	D_{13}		***	$\Sigma(1560)$		**			$\bullet K^*(892)$	1/2(1 ⁻)	
$\Lambda(1800)$		S_{01}	***	$\Xi(1950)$			***	$\Sigma(1580)$	D_{13}	*			$\bullet K_1(1270)$	1/2(1 ⁺)	
$\Lambda(1810)$		P_{01}	***	$\Xi(2030)$			***	$\Sigma(1620)$	S_{11}	**			$\bullet K_1(1400)$	1/2(1 ⁺)	
$\Lambda(1820)$		F_{05}	****	$\Xi(2120)$			*	$\Sigma(1660)$	P_{11}	***			$\bullet K^*(1410)$	1/2(1 ⁻)	
$\Lambda(1830)$		D_{05}	****	$\Xi(2250)$			**	$\Sigma(1670)$	D_{13}	****			$\bullet K_0^*(1430)$	1/2(0 ⁺)	
$\Lambda(1890)$		P_{03}	****	$\Xi(2370)$			**	$\Sigma(1690)$		**			$\bullet K_2^*(1430)$	1/2(2 ⁺)	
$\Lambda(2000)$			*	$\Xi(2500)$			*	$\Sigma(1750)$	S_{11}	***			$K(1460)$	1/2(0 ⁻)	
$\Lambda(2020)$		F_{07}	*					$\Sigma(1770)$	P_{11}	*			$K_2(1580)$	1/2(2 ⁻)	
$\Lambda(2100)$		G_{07}	****					$\Sigma(1775)$	D_{15}	****			$K(1630)$	1/2(?)	
$\Lambda(2110)$								$\Sigma(1840)$	P_{13}	*			$K_1(1650)$	1/2(1 ⁺)	
$\Lambda(2325)$		F_{05}	***					$\Sigma(1880)$	P_{11}	**			$\bullet K^*(1680)$	1/2(1 ⁻)	
$\Lambda(2350)$		D_{03}	*					$\Sigma(1915)$	F_{15}	****			$\bullet K_2(1770)$	1/2(2 ⁻)	
$\Lambda(2585)$		H_{09}	***					$\Sigma(1940)$	D_{13}	***			$\bullet K_3^*(1780)$	1/2(3 ⁻)	
			**					$\Sigma(2000)$	S_{11}	*			$\bullet K_2(1820)$	1/2(2 ⁻)	
								$\Sigma(2030)$	F_{17}	****			$K(1830)$	1/2(0 ⁻)	
								$\Sigma(2070)$	F_{15}	*			$K_0^*(1950)$	1/2(0 ⁺)	
Ω^-			****					$\Sigma(2080)$	P_{13}	**			$K_2^*(1980)$	1/2(2 ⁺)	
$\Omega(2250)^-$			***					$\Sigma(2100)$	G_{17}	*			$\bullet K_4^*(2045)$	1/2(4 ⁺)	
$\Omega(2380)^-$			**					$\Sigma(2250)$		***			$K_2(2250)$	1/2(2 ⁻)	
$\Omega(2470)^-$			**					$\Sigma(2455)$		**			$K_3(2320)$	1/2(3 ⁺)	
								$\Sigma(2620)$		**			$K_5^*(2380)$	1/2(5 ⁻)	
Baryons, $S=-1,-2,-3$ (Antibaryons not shown)												Mesons, $S=\pm 1$			
								$\Sigma(3000)$		*			$K_4(2500)$	1/2(4 ⁻)	
								$\Sigma(3170)$		*			$K(3100)$? (? ??)	

Higher Symmetry

Experimental evidence for several 'multiplets of multiplets'

$J^P=0^-$

I	S=+1	S=0	S=-1
0		$\eta, \bar{\eta}$	
1/2	K		\bar{K}
1		π	

$J^P=1^-$

I	S=+1	S=0	S=-1
0		ω, φ	
1/2	K^*		\bar{K}^*
1		ρ	

$J^P=2^+$

I	S=+1	S=0	S=-1
0		f_0, f_1	
1/2	K^{**}		\bar{K}^{**}
1		a_2	

Mesons

$J^P=1/2^+$

I	S=-2	S=-1	S=0
0			Λ^0
1/2		Ξ	N
1			Σ

$J^P=3/2^+$

I	S=-3	S=-2	S=-1	S=0
0	Ω^-			
1/2			Ξ^*	
1				Σ^*
3/2				Δ

Baryons

Remember:

*Each square is a I-spin multiplet, with size $2I+1$
 Total of 45 particle states in this page!*

$SU(3)$ - I

Try to find a larger group to encompass both strangeness and isospin into a unified symmetry scheme.

Requirements:

2 commuting generators, since both S and I_3 are defined within any observed supermultiplet

($SU(2)$ has just one, I_3)

Multiplet structure matching experimental data

$SU(3)$ - II

Take $SU(3)$ as candidate to extend $SU(2)$:

Group of unitary, unimodular 3×3 matrices

9 complex parameters \rightarrow 18 real parameters

9 unitarity conditions:
$$\left. \begin{array}{l} UU^\dagger = 1 \\ (U^\dagger)_{ij} = U_{ji}^* \end{array} \right\} \rightarrow \sum_{j=1}^3 a_{ij} a_{jk}^* = \delta_{ik}, \quad i, k = 1, \dots, 3$$

1 unimodularity condition: $\det U = 1$

$\rightarrow 18 - 10 = 8$ free, real parameters

$SU(3)$ - III

As usual, for any unitary matrix

$$U = e^{iH}, \quad H \text{ Hermitian}$$

$$\det U = 1 \rightarrow \det e^{iH} = 1 \rightarrow e^{itr(H)} = 1 \rightarrow tr(H) = 0$$

8 parameters \rightarrow 8 generators

Generalize Pauli matrices to *Gell-Mann matrices*

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

$$\lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}$$

$SU(3)$ - IV

Commutators:

$$[\lambda_i, \lambda_j] = f_{ijk} \lambda_k, \quad f_{ijk} \text{ structure constants}$$

Two diagonal generators, λ_3 and λ_8
 → Rank 2 group
 → 2 invariant functions of generators

Quadratic: $C^{(2)} = \sum_{i,j=1}^8 \delta_{ij} \lambda_i \lambda_j$

Cubic: $C^{(3)} = \sum_{i,j,k=1}^8 f_{ijk} \lambda_i \lambda_j \lambda_k$

$F_i \equiv \frac{\lambda_i}{2}$ Definition

Identify: $\begin{cases} I_3 = F_3 & \text{Isospin 3rd component} \\ Y = \frac{2}{\sqrt{3}} F_8 & \text{Hypercharge} \end{cases}$

Compare to $SU(2)$:

$$[\sigma_i, \sigma_j] = i \varepsilon_{ijk} \sigma_k$$

One diagonal generator, σ_3
 → Rank 1 group

→ 1 invariant function of generators


Quadratic: $C^{(2)} = \sum_{i,j=1}^3 \delta_{ij} \sigma_i \sigma_j$

$SU(3)$ Surprises

Fundamental representation (3×3 matrices): **3**

Find eigenvalues & eigenvectors for 3:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} I_3 = \frac{1}{2} \\ Y = \frac{1}{3} \end{cases}; \quad \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \rightarrow \begin{cases} I_3 = -\frac{1}{2} \\ Y = \frac{1}{3} \end{cases}; \quad \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \rightarrow \begin{cases} I_3 = 0 \\ Y = -\frac{2}{3} \end{cases}$$



→ 3 independent base states

→ I_3, Y eigenvalues not symmetrical wrt origin

→ Conjugate representation: 3^* different from 3

→ For both $3, 3^*$ hypercharge eigenvalues fractionary $Y = B + S$

→ $Q = I_3 + Y/2$ fractionary!!!

$SU(3)$ Multiplets - I


States identified by Y, I_3 eigenvalues
→ Points in a plane

Hexagonal/Triangular symmetry

Specified by 2 integers (p, q)

Multiplicity (i.e. size)


$$n = \frac{1}{2}(p+1)(q+1)(p+q+2)$$



Multiplet $(1,4)$
Frequently indicated by $n=35$

$SU(3)$ Multiplets - II

Products and decomposition into irr.rep.:
Proceed graphically as for $SU(2)$





Hadrons and $SU(3)$: The Eightfold Way


- All the hadronic multiplets nicely fit some $SU(3)$ representation
- No hadron found which does not fit

Baryons $J^P = 1/2^+, 3/2^+$

Mesons $J^{PC} = 0^{-+}$



8 10



The Hard Facts: $SU(3)$ Breaking

$J^P=0^-$

I	$S=-1$	$S=0$	$S=+1$
0		$\eta(547), \eta'(958)$	
1/2	$\bar{K}(496)$		$K(496)$
1		$\pi(137)$	

$J^P=1/2^+$

I	$S=-2$	$S=-1$	$S=0$
0		$\Lambda^0(1116)$	
1/2	$\Xi(1317)$		$N(938)$
1		$\Sigma(1192)$	

$J^P=1^-$

I	$S=-1$	$S=0$	$S=+1$
0		$\omega(782), \varphi(1020)$	
1/2	$\bar{K}^*(892)$		$K^*(892)$
1		$\rho(770)$	

$J^P=3/2^+$

I	$S=-3$	$S=-2$	$S=-1$	$S=0$
0	$\Omega^-(1672)$			
1/2		$\Xi^*(1530)$		
1			$\Sigma^*(1385)$	
3/2				$\Delta(1232)$

As before, but including masses:
 $SU(3)$ is not an exact symmetry

Mass differences within a multiplet are large, typ. $\Delta m/m \sim 10\text{-}20\%$

$SU(3)$ Breaking: Mass Formulas - I

Since $SU(3)$ is a broken symmetry, try to find a sensible breaking scheme

Take an effective Hamiltonian:

Part $SU(3)$ -Invariant + Part non $SU(3)$ -Invariant

$$m_{hadron} \simeq \langle hadron | H_s | hadron \rangle, \quad H_s = H_0 + H'$$

$$\langle a | H_s | a \rangle \rightarrow \underbrace{\langle a | U^{-1}}_{\substack{SU(3)-transformed \\ state}} H \underbrace{| a \rangle}_{\substack{SU(3)-transformed \\ state}}$$

$$\rightarrow \langle a | U^{-1}(H_0 + H')U | a \rangle = \langle a | U^{-1}H_0U | a \rangle + \langle a | U^{-1}H'U | a \rangle$$

$$H_0: \text{ invariant} \quad \rightarrow U^{-1}H_0U = H_0$$

$$H': \text{ non invariant} \rightarrow U^{-1}H'U \neq H'$$

$$\rightarrow \langle a | H | a \rangle = \langle a | U^{-1}H_0U | a \rangle + \langle a | U^{-1}H'U | a \rangle = \langle a | H_0 | a \rangle + \langle a | U^{-1}H'U | a \rangle$$

Must guess $SU(3)$ properties of H'

$SU(3)$ Breaking: Mass Formulas - II

Must guess $SU(3)$ properties of H'
Since the largest breaking concerns strange particles, suppose

$$\rightarrow H' \propto F_8 \propto Y$$

Reminder: $I_3 = F_3$, $Y = \frac{2}{\sqrt{3}} F_8$

According to $SU(3)$ algebra:

Gell-Mann Okubo mass formula

$$\langle a | H' | a \rangle \propto \langle a | F_8 | a \rangle \propto A + BY + C [Y^2/4 - I(I+1)]$$

A, B, C :
constants, rep. dependent

$$m(Y, I) = m_0 + bY + C [Y^2/4 - I(I+1)]$$

$SU(3)$ Breaking: Mass Formulas - III

$S = -3$ decuplet member not observed.

What is the mass?

Take mass differences between decuplet members:


$$\Delta m_{ij} = m_i - m_j = b(\Delta Y)_{ij} + C \left[\left(Y_i^2 - Y_j^2 \right) / 4 - \left(I_i(I_i+1) - I_j(I_j+1) \right) \right]$$

From $\Delta(1232)$, $\Sigma^*(1385)$, $\Xi^*(1530)$:

$$m_\Sigma - m_\Delta \approx m_\Xi - m_\Sigma \approx 150 \text{ MeV}$$

→ Predict missing $S = -3$, $J = 3/2$ decuplet baryon

Named Ω^- , predicted mass $m_\Omega \simeq 1672 \text{ MeV}$



The Ω^- Discovery at BNL

