

Elementary Particles II

1 – QCD

Color, Gauge Fields, Gluons,
Asymptotic Freedom, Confinement,
Perturbative QCD, Quarkonium

Re-examining the Evidence

2

Experiments probing the EM structure, like DIS:

Scaling of the structure functions

Evidence for point-like constituents, funny behavior:

Like free particles when interacting with EM currents at high Q^2

Never observed outside hadrons → Tightly bound?

Experiments probing the strong interaction:

Large particle zoo

Evidence for highly symmetrical grouping and ordering

Strong suggestion of a substructure: Quarks

Funny, ad-hoc rules driving the observed symmetry

Constituents?

3

Besides the general, puzzling behavior observed, finally identify a number of serious problems, cast into a few issues:

*Baryons and the Pauli Principle
R Ratio
 π^0 Decay Rate
 τ Lepton Branching Ratios*

From all these questions a common conclusion:

Our picture of the quark model is not complete

Pauli Principle

4

Quark model:

Besides its many, remarkable successes, a central point is at issue:

The baryon wave function (space \times spin \times flavor) is symmetric

Pauli Principle seems to be lost, which is very bad news:


The Spin-Statistics Theorem is a consequence of very general principles of relativistic QFT, not easily dismissed

R Ratio - I

5

Assume the process $e^+e^- \rightarrow \text{hadrons}$ to proceed at the lowest order through

$$e^+e^- \rightarrow q\bar{q} \rightarrow \text{hadrons}$$



As for DIS:

Don't care about quark *hadronization*, assume the time scales for hard and soft sub-processes to be wildly different

$$\sigma(e^+e^- \rightarrow \mu^+\mu^-) = \frac{4\pi\alpha^2}{3s}$$

$$\sigma(e^+e^- \rightarrow q\bar{q}) = \frac{4\pi\alpha^2 Q_q^2}{3s}, \quad Q_q = \text{quark charge in } e \text{ units}$$

$$R(E_{CM}) = \frac{\sigma(e^+e^- \rightarrow \text{adroni})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = \frac{\sum_q \sigma(e^+e^- \rightarrow q\bar{q})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = \sum_q Q_q^2$$

Sum extended to all accessible quark flavors $\rightarrow 2m_q < E_{CM}$

R Ratio - II

6

R counts the number of different quark species created at any given E_{CM} . Expect:

$$u,d \rightarrow R = \left(\frac{2}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 = \frac{5}{9} \quad \text{Low energy}$$

$$u,d,s \rightarrow R = \left(\frac{2}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 = \frac{6}{9} \quad E > 1-1.5 \text{ GeV}$$

$$u,d,s,c \rightarrow R = \left(\frac{2}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(\frac{2}{3}\right)^2 = \frac{10}{9} \quad E > 3 \text{ GeV}$$

By taking 3 quark species
of any flavor:

$$u,d \rightarrow R = \frac{15}{9}$$


$$u,d,s \rightarrow R = \frac{18}{9}$$

$$u,d,s,c \rightarrow R = \frac{30}{9}$$

R Ratio - III

7


$$R = \sigma(\text{hadrons})/\sigma(\mu^+\mu^-)$$



π^0 Decay Rate - I

8

Originally calculated by taking p, \bar{p} in the triangle loop (Steinberger 1949)



Steinberger's calculation:

Yukawa model to account for πpp vertex

Point-like nucleons \rightarrow QED couplings to photons

Nucleon current in the loop: 4-vector J^μ

[Actually *axial* vector, to match pion -ve parity]

π^0 spinless: Only 4-vector available p_μ

\rightarrow Decay amplitude $\sim p_\mu J^\mu$

π^0 Decay Rate - II

9

With a proton loop rate OK (!)

By naively replacing the proton loop by a quark loop:

$$J^\mu \approx \sum_{i=u}^d q_i \bar{\psi}_i \gamma^\mu \gamma^5 \psi_i = e \left(\frac{2}{3} \bar{u} \gamma^\mu \gamma^5 u - \frac{1}{3} \bar{d} \gamma^\mu \gamma^5 d \right)$$

Amplitude: 2 vertexes

Each vertex $\propto \sqrt{\alpha} = e \rightarrow$ Amplitude $\propto e^2$

Sum over light quarks u, d :

$$\sum_{i=u,d} a_i Q_i^2 = e^2 \left[1 \cdot \left(\frac{2}{3} \right)^2 - 1 \cdot \left(-\frac{1}{3} \right)^2 \right] = e^2 \left[\frac{4}{9} - \frac{1}{9} \right] = e^2 \frac{1}{3}$$

$$\Gamma_{quark} (\pi^0 \rightarrow \gamma\gamma) = \frac{1}{9} \Gamma_{proton} (\pi^0 \rightarrow \gamma\gamma) \quad ???$$

\rightarrow Wrong by a factor 9!

Bad news for the quark model

π^0 Decay Rate - III

10

Upon looking more carefully to the problem, things look actually even worse:

By taking seriously the quark model, one cannot escape consequences of approximate *chiral symmetry* of light quarks

Then simple guess on approximate symmetry of the initial state would lead to conclude *the neutral pion is stable!* Another quark model puzzle..

Explanation of this paradox led to discovery of the first *anomaly* in QFT (Adler, Bell & Jackiw)


Advanced topic, quite relevant to the Standard Model:
Quantum field theories must be *anomaly free* in order to be renormalizable

Interesting conditions for SM to be anomaly free, including *charge quantization*

π^0 Decay Rate - IV

11

Direct method:




π^0 produced in a first thin foil, when not decayed do not contribute to e^+ yield from γ conversion in a second thin foil

$$Y(d) = N \{ A + B [1 - \exp(-d/\lambda)] \}$$

$\lambda = \beta \gamma c \tau \simeq \gamma c \tau$ Energy dependent


Use known energy spectra for pions



@TBA

$$\tau = 0.897 \pm 0.021 \cdot 10^{-16} \text{ s}$$


$$\Gamma = 7.34 \pm 0.18 \pm 0.11 \text{ eV}$$



π^0 Decay Rate - V

12

Primakoff effect



Very simple idea:
Get a high energy photon beam + high Z target
Pick-up a virtual photon from the nuclear Coulomb field
2-photon coupling will (sometimes) create a π^0

$$\frac{d\sigma_{\text{Prim}}}{d\Omega_{\text{LAB}}} \simeq \Gamma_{\pi^0 \rightarrow \gamma\gamma} Z^2 \frac{8\alpha E^4}{m_{\pi^0}^2} \frac{\left| F(q^2) \right|^2}{q^4} \sin^2 \theta_{\pi^0}$$



Strongly forward peaked
Quickly increasing with energy
Strongly Z dependent: Coherence

$\Gamma = 1/\tau$ extracted by measuring the differential cross-section
Nuclear form factor required

π^0 Decay Rate - VI

13

Recent experiment: PrimEx at Jefferson Lab (Virginia)




τ Lepton Decays

14

τ : Heavy brother of e and μ

$m_\tau = 1776$ MeV

Weak decays:



$q, \bar{q} : u, d, s$

$BR(e) \sim 18\%$

$BR(\mu) \sim 17\%$

$BR(q\bar{q}) \sim 65\%$

In the absence of color, weak interaction universality would lead to predict:

$$BR(e) \sim BR(\mu) \sim BR(q\bar{q}) \sim 33\%$$

With color:

$$\Gamma(q\bar{q}) \sim 3 \Gamma(l\bar{l}) \rightarrow BR(q\bar{q}) \sim \frac{3 \Gamma(l\bar{l})}{3 \Gamma(l\bar{l}) + 2 \Gamma(l\bar{l})} \sim 60 \% \text{ OK}$$

Color - I

15

New hypothesis:

There is a new degree of freedom for quarks: Color

Each quark can be found in one of 3 different states
Internal space (mathematically identical to flavor):

States = 3-component complex vectors

Base states:

$$R(ed) = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, G(reen) = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, B(lue) = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Needless to say, nothing to do with our old, beloved color concept (nature, art, politics..):

Just a *name* for another, non-classical property of hadron constituents

Color - II

16

Provided one can build a color wave function for 3 fermions which is antisymmetric, the Fermi statistics problem is solved

Total wave function of a baryon

$$\psi = \psi_{color} \underbrace{\psi_{orbital} \psi_{spin} \psi_{flavor}}_{Symmetric} \rightarrow \psi_{color}: \text{Antisymmetric}$$

To account for 3 different color states, the R ratio must be multiplied by 3 \rightarrow OK with experimental data

Just the same conclusion for hadronic τ decays: Multiply rate by 3

The correct π^0 rate is obtained by inserting a factor 9

Color - III

17

Observe:

When computing R , τ decay rates we add the *rates* for different colors
→Factor $\times 3$

We deal with quarks as with real, on-shell particles: Ignore fragmentation

When computing π^0 decay rate, we add the *amplitudes*
→Factor $\times 9$

Quarks in the loop are virtual particles: *Amplitudes interfere*

Color - IV

18

Must be possible to build hadron states as color *singlets*

Do not expect hadrons to fill larger irr.reps.:

Would imply large degeneracies for hadron states, not observed

In other words:

Color is fine, but we do not observe any colored hadron

Therefore we assume the color charge is *confined*:

Never observed directly, as it is the electric charge

Why? For the moment, nobody really knows

Color - V

19

How colored hadrons would show up?

Just as an example:

Should the nucleon fill the **3** of $SU(3)_C$, there would be 3 different species of protons and neutrons.

Then each nucleon level in any nucleus could accommodate 3 particles instead of one:

The nuclear level scheme would be far different from the observed one

Color - VI

20

Guess $SU(3)$ as the color group

Take the two fundamental decompositions:

$$\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$$

Baryons

$$\mathbf{3} \otimes \mathbf{3}^* = \mathbf{1} \oplus \mathbf{8}$$

Mesons

Both feature a singlet in the direct sum: OK

No singlets in $\mathbf{3} \otimes \mathbf{3}$: OK

Can't say the same for other groups...

Take $SU(2)$ as an example:

Say the quarks live in the adjoint $SU(2)$ representation, $\mathbf{3}$

Then for qq


$$\mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{5}$$

Observe: This is $\mathbf{3}$ of $SU(2)$, which is quite different from $\mathbf{3}$ of $SU(3)$

Diquarks can be in color singlet

→ Should find diquarks as commonly as baryons or mesons..

Colored Quarks



21

$$I_3^c = \frac{\lambda_3}{2}, \quad Y^c = \frac{\lambda_8}{\sqrt{3}}$$

	I_3^c	Y^c		I_3^c	Y^c
R	+1/2	+1/3	\bar{R}	-1/2	-1/3
G	-1/2	+1/3	\bar{G}	+1/2	-1/3
B	0	-2/3	\bar{B}	0	+2/3

$SU(3)_C$ is an *exact symmetry*:

$$m_u^{(R)} = m_u^{(G)} = m_u^{(B)}, \dots$$

Beware: $SU(3)_C$ has nothing to do with $SU(3)_F$:
Quark quantum numbers are independent from their color state

They are left unchanged by QCD transitions

Uncolored Hadrons

22

According to our fundamental hypothesis:

$$\text{Mesons: } \mathbf{3} \otimes \mathbf{3}^* = \mathbf{1} \oplus \mathbf{8}$$

$$\rightarrow \psi_c = \frac{1}{\sqrt{3}} (R\bar{R} + G\bar{G} + B\bar{B})$$

$$\text{Baryons: } \mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$$

$$\rightarrow \psi_c = \frac{1}{\sqrt{6}} (RGB - RBG + GBR - GRB + BRG - BGR)$$

In both cases, pick singlet

Mesons: *No particular exchange symmetry*
(2 non identical particles)

Baryons: *Fully antisymmetrical color wave function*
(3 identical particles)

Color Interaction: QCD

23

Color: A new degree of freedom for quarks
Compare to other quantum numbers:

Baryonic/Leptonic numbers
Conserved, *not originating interactions*

Electric charge
Conserved, *origin of the electromagnetic field*

A deep question:

What is the true origin of the electromagnetic interaction?

We have freely used the interaction term $j^\mu A_\mu$, only based on the classical analogy:
But supposedly quantum mechanics is more general than classical mechanics/electromagnetism..

Is there any deeper origin for it?

QED as a Gauge Theory - I

24

Symmetry:

Absolute phase not defined for a wave function.

Expect invariance as per our old acquaintance, Noether's Theorem

$$L_0 = \bar{\psi}(x) (i\gamma^\mu \partial_\mu - m) \psi(x) \quad \text{Free Dirac Lagrangian}$$

Global gauge (=Phase) transformation:

$$G : \psi(x) \rightarrow \psi'(x) = U_\theta \psi(x) = e^{-iq\theta} \psi(x) \quad q\theta : \text{New phase} \propto \text{Charge}$$

$\rightarrow L_0$ invariant wrt $G \rightarrow$ Charge conservation

Just meaning:

Take *all* particle states, re-phase each state proportionally to its charge

QED as a Gauge Theory - II

25

Generalize to local phase transformation:

$$G_L : \psi(x) \rightarrow \psi'(x) = U_\theta \psi(x) = e^{-iq\theta(x)} \psi(x) \text{ Local gauge transformation}$$

$\rightarrow L_0$ not invariant wrt G_L : Derivative term troublesome

$$L_0 = \bar{\psi}(x) (i\gamma^\mu \partial_\mu - m) \psi(x) \rightarrow L_0' = i\bar{\psi}(x) e^{+iq\theta(x)} \gamma^\mu \partial_\mu (e^{-iq\theta(x)} \psi(x)) - m\bar{\psi}(x) \psi(x)$$

$$L_0' = i[\bar{\psi}(x) \gamma^\mu \partial_\mu \psi(x) - iq\partial_\mu [\theta(x)] \psi(x)] - m\bar{\psi}(x) \psi(x)$$

$$L_0' = \{i\bar{\psi}(x) \gamma^\mu \partial_\mu \psi(x) + q\partial_\mu [\theta(x)] \psi(x) - m\bar{\psi}(x) \psi(x)\}$$

$$L_0' = [i\bar{\psi}(x) \gamma^\mu \partial_\mu \psi(x) - m\bar{\psi}(x) \psi(x)] + q\partial_\mu [\theta(x)] \psi(x) \neq L_0$$

\rightarrow Local gauge invariance cannot hold in a world of free particles

Symmetry requires interaction

QED as a Gauge Theory - III

26

New transformation rule:

$$\begin{cases} \psi(x) \rightarrow \psi'(x) = U_\theta \psi(x) = e^{-iq\theta(x)} \psi(x) & \text{As before} \\ A_\mu(x) \rightarrow A_\mu(x) + q \partial_\mu \theta(x) & \text{New character in the comedy} \end{cases}$$

Equivalent to re-define derivative for ψ :


$$\partial_\mu \rightarrow \partial_\mu + iqA_\mu \quad \text{Vector field}$$

Add a new term to Lagrangian:

$$L_i = -\underbrace{q\bar{\psi}(x)\gamma^\mu\psi(x)}_{j^\mu} A_\mu \quad \text{Interaction term}$$

Same as classical electrodynamics

$$L_0 = \bar{\psi}(x)(i\gamma^\mu\partial_\mu - m)\psi(x) \rightarrow L_0 + L_i = \bar{\psi}(x)(i\gamma^\mu\partial_\mu - m)\psi(x) - q\bar{\psi}(x)\gamma^\mu\psi(x)A_\mu$$



Sum is invariant

QED as a Gauge Theory - IV

27

...And another one:

$$-\frac{1}{4}F^{\mu\nu}F_{\mu\nu}, \quad F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu \quad \text{Field energy}$$

Must be there because the field carries energy+momentum+angular momentum

Reminder:

$F^{\mu\nu}$ is the EM field

$$F^{\mu\nu} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix}$$

Field must be massless to have L gauge invariant

$$\frac{1}{2}m^2A_\mu^2 \rightarrow \frac{1}{2}m^2(A_\mu(x) + q\partial_\mu\theta(x))^2 \neq \frac{1}{2}m^2A_\mu^2 \quad \text{if } m \neq 0$$

QED as a Gauge Theory - V

28

Consider all the phase transformations as defined before:

$$\psi(x) \rightarrow \psi'(x) = U_\theta \psi(x) = e^{-iq\theta(x)} \psi(x)$$

The full set is a group: $U(1)$ Unitary, 1-dimensional

$$e^{-iq\theta_1(x)} e^{-iq\theta_2(x)} \psi(x) = e^{-iq[\theta_1(x)+\theta_2(x)]} \in U(1)$$

1 parameter: $\theta(x)$

$$\text{Abelian: } e^{-iq\theta_1(x)} e^{-iq\theta_2(x)} \psi(x) = e^{-iq\theta_2(x)} e^{-iq\theta_1(x)} \psi(x)$$

$U(1)$ is the (Abelian) *gauge group* of QED

Equivalent to $SO(2)$, group of 2D rotations

QCD as a Gauge Theory - I

29

Extend gauge transformations to a 3-component wave function:

$$\Psi \equiv \begin{pmatrix} \psi_R \\ \psi_G \\ \psi_B \end{pmatrix}$$

Global gauge transformation: Phase change for individual components

→ Phase change will mix color components

$$G_G^C : \Psi(x) \rightarrow \Psi'(x) = \mathbf{U}_G \cdot \Psi(x) = e^{-ig\mathbf{M}} \cdot \Psi(x) \quad \mathbf{U}_G \text{ unitary} \rightarrow \mathbf{M} \text{ Hermitian}$$

$$e^{-ig\mathbf{M}} = \mathbf{1} - ig\mathbf{M} + \frac{(-ig\mathbf{M})^2}{2!} + \dots \quad \mathbf{M}: 3 \times 3 \text{ Hermitian matrix}$$

\mathbf{M} acting on the 3 color components of the quark state

Since the color symmetry group is $SU(3)_c$:

$$\mathbf{M} = \sum_{i=1}^8 \lambda_i \theta_i \equiv \vec{\lambda} \cdot \vec{\theta}$$

$\vec{\lambda}$: Vector of 8 3×3 Gell-Mann matrices; $\vec{\theta}$: Vector of 8 parameters

QCD as a Gauge Theory - II

30

As for QED, extend to *local* gauge transformations

As before, in order to guarantee invariance of L :

→ Re-define derivative adding new vector fields:

$$\partial_\mu \rightarrow \partial_\mu \mathbf{1} + ig \mathbf{C}_\mu$$

$$\mathbf{C}_\mu : \begin{cases} \text{4-vector field} & \text{Lorentz structure} \\ \text{Matrix } \in SU(3)_c & \text{Color space} \end{cases}$$

We know how to express any Hermitian matrix $\in SU(3)_c$:

Use $SU(3)_c$ generators → Gell-Mann matrices

$$\rightarrow \mathbf{C}_\mu = \frac{1}{2} \sum_{i=1}^8 \mathbf{G}_\mu^a \boldsymbol{\lambda}_a \equiv \vec{G}_\mu \cdot \vec{\lambda} \quad 8 \text{ fields required: Gluons}$$

So gluons are a bit like 8 different “photons”, exchanged between color charges

But: *They are non Abelian*

QCD as a Gauge Theory - III

31

Local gauge transformation for $SU(3)_C$:

$$\begin{cases} \Psi(x) \rightarrow \Psi'(x) = U_\theta \Psi(x) = e^{-ig\vec{\lambda} \cdot \vec{\theta}(x)} \Psi(x) \\ G_\mu^a(x) \rightarrow G_\mu^a(x)' = G_\mu^a(x) + \partial_\mu \theta^a + g \sum_{b,c=1}^8 f^{abc} G_\mu^b(x) \theta^c(x) \quad a = 1, \dots, 8 \end{cases}$$

Very important: New term, coming from $SU(3)$
being non Abelian

Reminder:

$$\left[\frac{\lambda_a}{2}, \frac{\lambda_b}{2} \right] = if^{abc} \frac{\lambda_c}{2} \quad f^{abc}: SU(3) \text{ structure constants}$$

$$L_0 = \bar{\Psi}(x) (i\gamma^\mu \partial_\mu - m) \Psi(x) \rightarrow L_0 + L_i$$

$$L_i = -g \underbrace{\left[\bar{\Psi}(x) \gamma^\mu \left(\frac{\vec{\lambda}}{2} \right) \Psi(x) \right]}_{\vec{j}_\mu} \cdot \vec{G}_\mu \quad \text{Interaction term}$$

$$-\frac{1}{4} \vec{G}_{\mu\nu} \cdot \vec{G}^{\mu\nu} = -\frac{1}{4} \sum_{a=1}^8 G_{\mu\nu}^a \cdot G^{a\mu\nu} \quad \text{Field energy term}$$

QCD as a Gauge Theory - IV

32

Take the expression of fields in terms of potentials:

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a - g \sum_{b,c=1}^8 f_{abc} G_\mu^b G_\nu^c$$

Very important: *Absent in QED (f=0)*
New term, coming from $SU(3)$ being non Abelian

$\rightarrow G_{\mu\nu}^a G_{\mu\nu}^a$ contains terms with $\underbrace{\partial_\mu G_\nu^a \cdot G_\mu^b G_\nu^c}_{3 \text{ gluons}}, \underbrace{G_\mu^b G_\nu^c \cdot G_\mu^b G_\nu^c}_{4 \text{ gluons}}$

When translated into Feynman rules/diagrams, these pieces of L correspond to 3 and 4 gluons vertices

So:

The form of QCD Lagrangian leads to predict the existence of a new kind of *gluon-gluon color interaction*

QCD as a Gauge Theory - V

33

Since color interaction is tied to color charge, we are saying *the gluons carry their own color charge*

Sounds unfamiliar? Well, that's all after playing with a non-Abelian gauge group.

Unlike the electric charge, color charge can manifest itself in more than one way.


Indeed, gluons carry a type of color charge different from quarks/antiquarks:

Color + Anticolor



QCD as a Gauge Theory - VI

34


QCD Vertices



$$ig \frac{\lambda_a}{2} \gamma_\mu \text{ Similar to QED:}$$



$$\propto g \quad (\text{Lorentz structure not shown})$$



$$\propto g^2 \quad (\text{Lorentz structure not shown})$$

Colored Gluons - I

35

Compare to mesons in $SU(3)_F$: *Flavor + Antiflavor*

But: *Gluons are not bound states of Color+Anticlor!*

Still, they share the same math:

Gluons live in the adjoint (8) irr.rep. of $SU(3)_C$

$$|1\rangle = \frac{1}{\sqrt{2}}(r\bar{b} + b\bar{r}), |2\rangle = -\frac{i}{\sqrt{2}}(r\bar{b} - b\bar{r}), |3\rangle = \frac{1}{\sqrt{2}}(r\bar{r} - b\bar{b})$$

$$|4\rangle = \frac{1}{\sqrt{2}}(r\bar{g} + g\bar{r}), |5\rangle = -\frac{i}{\sqrt{2}}(r\bar{g} - g\bar{r}), |6\rangle = \frac{1}{\sqrt{2}}(b\bar{g} + g\bar{b})$$

$$|7\rangle = -\frac{i}{\sqrt{2}}(b\bar{g} - g\bar{b}), |8\rangle = \frac{1}{\sqrt{6}}(r\bar{r} + b\bar{b} - 2g\bar{g})$$

Colored Gluons - II

36


A very natural question: Gluons couple to $q\bar{q}$

Since one can decompose the total $q\bar{q}$ color state as:

$$\mathbf{3} \otimes \mathbf{3}^* = \mathbf{1} \oplus \mathbf{8}$$

Then: Where is the singlet gluon?

Does not exist: There are only 8 gluons, not 9



Should the singlet gluon actually exist, it would behave more or less like a “photon”:

Would be ‘white’ (= Singlet)

Would couple to color charges in the same way as photon couples to electric charges

Would give rise to a sort of “QED-like”, long range color interaction, not observed

Colored Gluons - III

37

Non Abelian vertices: Gluon-Gluon scattering *at tree level* (no loops)



@TBA


Vertexes:

$$3\text{-gluons} : A \propto g$$

$$4\text{-gluons} : A \propto g^2$$

Compare:


In QED, photon-photon scattering amplitude occurs at order α^2 through the 1-loop diagram



Comparing QED and QCD - I

38

Compare the different situations:



QED
Photon is *neutral*

Neither sourcing,
nor sinking charge

QCD
Gluon is *colored*

Sourcing color,
sinking anti-color

Comparing QED and QCD - II

39

Comparison of coupling constants:

α vs. α_s Adimensional constants (*Interaction strength*)

Can define elementary charge in terms of α, α_s

Measure particle charge by its ratio to elementary charge:

Number

What are the allowed values for these numbers?

Comparing QED and QCD - III

40

QED: Gauge group is *Abelian*

Electric charge can be *any* number:

No reason for charge quantization → So electric charge quantization is a bit of a mystery

[Tricky business: Sticking to perturbation theory, one must have the SM *anomaly-free* in order to be renormalizable → This in turn *requires* charge quantization.

But: Is the SM just perturbation theory?

At a fundamental level, Grand Unified Theories explain charge quantization based on larger symmetry groups like $SU(5)$.

But: They fail to explain proton stability]

Photon charge is strictly 0

QCD: Gauge group is *non Abelian*

“Color charge” value is *fixed* for every representation

Quarks: $3, 3^*$ → $Q = 4/3$

Gluons: 8 → $Q = 3$

Similar to $I(I+1)$ for any isospin ($SU(2)$) multiplet

Color Factors - I

41

Consider the static interaction between 2 charges:

QED For fixed $|q|$, the 'charge factor' can be defined as:

$$f_{12} = \frac{q_1 q_2}{|q|^2} = \begin{cases} +1 & q_1 q_2 > 0 \\ -1 & q_1 q_2 < 0 \end{cases}$$

Very simple for an Abelian interaction

QCD The 'color factor' depends on the irr.rep. of the color state

Representation dependent

Identical for any transition in a given representation

→*Color Conservation*

Less simple in this non-Abelian interaction


Color Factors - II

42

$$q\bar{q} \rightarrow q\bar{q}$$


$$\mathbf{3} \otimes \mathbf{3^*} = \mathbf{1} \oplus \mathbf{8}$$

Total color conservation: $\begin{cases} \mathbf{1} \rightarrow \mathbf{1} \\ \mathbf{8} \rightarrow \mathbf{8} \end{cases}$



Observe:

Similar to conservation of total I-spin



$$T_{fi} = i \sum_{\alpha, \beta=1}^8 \underbrace{\left[\bar{u}(3)c_3^\dagger \right] \left[-i \frac{g_s}{2} \lambda^\alpha \gamma^\mu \right] \left[u(1)c_1 \right]}_{\text{color current}} \underbrace{\left[-i \frac{g_{\mu\nu} \delta^{\alpha\beta}}{q^2} \right] \left[\bar{v}(2)c_2^\dagger \right] \left[-i \frac{g_s}{2} \lambda^\beta \gamma^\nu \right]}_{\text{propagator}} \underbrace{\left[v(4)c_4 \right]}_{\text{color current}}$$

Sum is over all 8 color matrices

c_i are the color states of initial, final $q\bar{q}$

$$T_{fi} = \frac{-g_s^2}{q^2} \left[\bar{u}(3)\gamma^\mu u(1) \right] \left[\bar{v}(2)\gamma_\mu v(4) \right] \underbrace{\frac{1}{4} \sum_\alpha \left[c_3^\dagger \lambda^\alpha c_1 \right] \left[c_2^\dagger \lambda^\alpha c_4 \right]}_{\text{color factor}}$$

Color Factors - III

43

Octet

$r\bar{b}$

Just as an example: Result is the same for all octet states

$$c_1 = c_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad c_2 = c_4 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \rightarrow f = \frac{1}{4} \sum_{\alpha=1}^8 (1 \ 0 \ 0) \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} (0 \ 1 \ 0) \lambda^\alpha \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\rightarrow f = \frac{1}{4} \sum_{\alpha=1}^8 \lambda_{11}^\alpha \lambda_{22}^\alpha = \frac{1}{4} (\lambda_{11}^3 \lambda_{22}^3 + \lambda_{11}^8 \lambda_{22}^8) = -\frac{1}{6}$$

Color Factors - IV

44

Singlet

$$\frac{1}{\sqrt{3}}(r\bar{r} + b\bar{b} + g\bar{g}) \quad \text{Only this state in the singlet}$$

But: *Any* component can go into *any other..*

$$f_i = \frac{1}{4} \frac{1}{\sqrt{3}} \frac{1}{\sqrt{3}} \sum_{j=1}^3 \sum_{\alpha=1}^8 [c_i^\dagger \lambda^\alpha c_j] [c_j^\dagger \lambda^\alpha c_i], \quad i = 1, 2, 3$$

$$f = \sum_{i=1}^3 f_i = \frac{1}{4} \frac{1}{\sqrt{3}} \frac{1}{\sqrt{3}} \sum_{\alpha=1}^8 \sum_{i,j=1}^3 \lambda_{ij}^\alpha \lambda_{ji}^\alpha = \frac{1}{12} \sum_{\alpha=1}^8 \text{Tr}(\lambda^\alpha \lambda^\alpha)$$

$$\text{Tr}(\lambda^\alpha \lambda^\beta) = 2\delta^{\alpha\beta} \rightarrow \sum_{\alpha=1}^8 \text{Tr}(\lambda^\alpha \lambda^\alpha) = 16$$


$$\rightarrow f = \frac{4}{3}$$

Color Factors - V

45

$qq \rightarrow qq$

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{3}^* \oplus \mathbf{6}$$



$$T_{fi} = \frac{-g_s^2}{q^2} [\bar{u}(3) \gamma^\mu u(1)] [\bar{u}(4) \gamma_\mu u(2)] \underbrace{\frac{1}{4} \sum_{\alpha=1}^8 [c_3^\dagger \lambda^\alpha c_1] [c_4^\dagger \lambda^\alpha c_2]}_{\text{color factor}}$$

$$f_{qq} = \frac{1}{4} \sum_{\alpha=1}^8 (c_3^\dagger \lambda^\alpha c_1) (c_4^\dagger \lambda^\alpha c_2)$$

Color Factors - VI

46

Color states: Triplet and sextet:

$$\mathbf{3}^*: \quad \frac{1}{\sqrt{2}}(rb - br), \frac{1}{\sqrt{2}}(bg - gb), \frac{1}{\sqrt{2}}(gr - rg)$$

Antisymmetric

$$\mathbf{6}: \quad rr, bb, gg, \frac{1}{\sqrt{2}}(rb + br), \frac{1}{\sqrt{2}}(bg + gb), \frac{1}{\sqrt{2}}(gr + rg)$$

Symmetric

Color Factors - VII

47

Sextet

rr

Just as an example: Result is the same for all sextet states

$$c_1 = c_2 = c_3 = c_4 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$
$$f = \frac{1}{4} \sum_{\alpha=1}^8 \left[(1 \ 0 \ 0) \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right] \left[(1 \ 0 \ 0) \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right] = \frac{1}{4} \sum_{\alpha=1}^8 (\lambda_{11}^\alpha \lambda_{11}^\alpha)$$
$$= \frac{1}{4} (\lambda_{11}^3 \lambda_{11}^3 + \lambda_{11}^8 \lambda_{11}^8) = \frac{1}{3}$$

Color Factors - VIII

48

Triplet

$$\frac{1}{\sqrt{2}}(rb - br)$$

Just as an example as before

$$f = \frac{1}{4} \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} \sum_{\alpha=1}^8$$

$$\left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right] \left[\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] - \left[\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right] \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right]$$

$$- \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] \left[\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right] - \left[\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right] \left[\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda^\alpha \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right]$$

$$f = \frac{1}{8} \sum_{\alpha=1}^8 \left\{ \lambda_{11}^\alpha \lambda_{22}^\alpha - \lambda_{21}^\alpha \lambda_{12}^\alpha - \lambda_{12}^\alpha \lambda_{21}^\alpha + \lambda_{22}^\alpha \lambda_{11}^\alpha \right\}$$

$$\rightarrow f = \frac{1}{4} \sum_{\alpha=1}^8 \left\{ \lambda_{11}^\alpha \lambda_{22}^\alpha - \lambda_{12}^\alpha \lambda_{21}^\alpha \right\} = \frac{1}{4} \left\{ \lambda_{11}^3 \lambda_{22}^3 + \lambda_{11}^8 \lambda_{22}^8 - \lambda_{12}^1 \lambda_{21}^1 - \lambda_{12}^2 \lambda_{21}^2 \right\} = -\frac{2}{3}$$

Color Factors - IX

49

Matrix elements just calculated:

Very similar to the corresponding tree-level amplitudes in QED

→Expect similar, Coulomb-like, effective potential in the static limit

Constant depends on the color representation for the quark pair:

$$V_{q\bar{q}} = \begin{cases} -\frac{4}{3} \frac{\alpha_s}{r} & \text{singlet} \\ \frac{1}{6} \frac{\alpha_s}{r} & \text{octet} \end{cases} \quad \text{Attractive}$$

$$V_{qq} = \begin{cases} -\frac{2}{3} \frac{\alpha_s}{r} & \text{triplet} \\ \frac{1}{3} \frac{\alpha_s}{r} & \text{sextet} \end{cases} \quad \text{Attractive}$$

Expect maximal attraction in singlet

Color Factors - X

50

Baryons could be in any one of the **1, 8, 10** representations:

Why only the singlet is observed?

A hint of an explanation:

$$\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = (\mathbf{3} \otimes \mathbf{3}) \otimes \mathbf{3}$$

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{6} \oplus \mathbf{3}^* \rightarrow (\mathbf{3} \otimes \mathbf{3}) \otimes \mathbf{3} = (\mathbf{6} \oplus \mathbf{3}^*) \otimes \mathbf{3}$$

$$\mathbf{6} \otimes \mathbf{3} = \mathbf{10} \oplus \mathbf{8}$$

$$\mathbf{3}^* \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8}$$

1: each qq pair is a triplet \rightarrow attractive

8: qq pair can be triplets, or sextet \rightarrow attractive + repulsive

10: each qq pair is a sextet \rightarrow repulsive


So singlet is the state maximally attractive for 3 quarks

Does this explain the singlet-only mystery of bound states?

Running Coupling: QED - I

51

Consider the *one loop* modification to the photon propagator:



Includes a sum over P , the momentum circulating in the virtual loop. No obvious bounds on P ..

$$M \propto [e\bar{u}(k')\gamma^\mu u(k)] \frac{g_{\mu\nu}}{q^2} \frac{1}{(2\pi)^4} \int d^4P \frac{[e\bar{u}(P)\gamma^\mu u(P-q)]}{P^2 - m^2} \frac{[e\bar{u}(P-q)\gamma^\nu u(P)]}{(P-q)^2 - m^2} \frac{g_{\nu\nu'}}{q^2} [e\bar{u}(p')\gamma^\nu u(p)]$$

Modified propagator:

$$\frac{g_{\mu\nu}}{q^2} \rightarrow \frac{g_{\mu\nu}}{q^2} (1 - I(q^2)), \quad I(q^2) = \frac{\alpha}{3\pi} \int_{m^2}^{\infty} \frac{dp^2}{p^2} - \frac{2\alpha}{\pi} \int_0^1 dx x (1-x) \ln \left[1 - \frac{q^2 x (1-x)}{m^2} \right]$$

Running Coupling: QED - II

52

Take the high q^2 approximation

$$-q^2 \gg m^2 \rightarrow \ln \left[1 - \frac{q^2 x(1-x)}{m^2} \right] \approx \ln \left[-\frac{q^2}{m^2} \right]$$


$$I(q^2) \approx \frac{\alpha}{3\pi} \int_{m^2}^{M^2} \frac{dp^2}{p^2} - \frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \ln \left[\frac{-q^2}{m^2} \right]$$

$$I(q^2) \approx \frac{\alpha}{3\pi} \ln \left(\frac{M^2}{m^2} \right) - \frac{2\alpha}{\pi} \frac{1}{6} \ln \left[\frac{-q^2}{m^2} \right] = \frac{\alpha}{3\pi} \left[\ln \left(\frac{M^2}{m^2} \right) - \ln \left[\frac{-q^2}{m^2} \right] \right] = \frac{\alpha}{3\pi} \ln \left(\frac{M^2}{-q^2} \right)$$

$$M \propto \alpha \left[\bar{u}(k') \gamma^\mu u(k) \right] \frac{g_{\mu\nu}}{q^2} \left[1 - \frac{\alpha}{3\pi} \ln \left(\frac{M^2}{-q^2} \right) \right] \left[\bar{u}(p') \gamma^\nu u(p) \right]$$

Provisional upper bound (cutoff) to make integral converging

Cartoon translation:



Running Coupling: QED - III

53

Extend to diagrams with $2, 3, \dots, n, \dots$ loops: Add up all contributes

Sum of a ‘geometrical series’: Converging ??

$$\text{M}\alpha = \begin{aligned} & \left[1 - \left[\text{loop diagram} \right] + \left[\text{loop diagram} \right]^2 + \dots \right] \\ & = \frac{1}{1 - \left[\text{loop diagram} \right]} \end{aligned}$$

Experts say this is the only contribution to running α to the ‘leading logs’ approximation, which means neglecting the next levels of iteration

Running Coupling: QED - IV

54

$$M \propto [\bar{u}(k')\gamma^\mu u(k)] \frac{g_{\mu\nu}}{q^2} \left[\frac{\alpha}{1 + \alpha/3\pi \ln(M^2/-q^2)} \right] [\bar{u}(p')\gamma^\nu u(p)]$$

What is α ?

Coupling 'constant' we would get should we turn off all loops

Call it α_0 = 'Bare' coupling constant, not physical:

Loops cannot be turned off

Then obtain an effective coupling, not constant but running:

$$\alpha(q^2) = \frac{\alpha_0}{1 + \alpha_0/3\pi \ln(M^2/-q^2)}$$

α is q^2 , or distance, dependent!

Running Coupling: QED - V

55

Running α is still cutoff dependent, which of course is uncomfortable
But: Not a real problem.
Indeed:

$$Q^2 = -q^2 \rightarrow \alpha(Q^2) = \frac{\alpha_0}{1 + (\alpha_0/3\pi) \ln(M^2/Q^2)}$$

Take a particular energy scale : $Q^2 = \mu^2$

$$\rightarrow \alpha(\mu^2) = \frac{\alpha_0}{1 + (\alpha_0/3\pi) \ln(M^2/\mu^2)}$$

Usually choose $\mu^2=0$, i.e. take α at distance $\rightarrow \infty$

Quite natural in QED (but not compulsory)

Running Coupling: QED - VI

56

$$\ln\left(\frac{M^2}{Q^2}\right) = \ln\left(\frac{M^2}{Q^2} \frac{\mu^2}{\mu^2}\right) = \ln\left(\frac{M^2}{\mu^2}\right) + \ln\left(\frac{\mu^2}{Q^2}\right)$$

$$\rightarrow \alpha(Q^2) = \frac{\alpha_0}{1 + (\alpha_0/3\pi)[\ln(M^2/\mu^2) + \ln(\mu^2/Q^2)]}$$

$$\rightarrow \frac{\alpha_0}{\alpha(\mu^2)} = 1 + (\alpha_0/3\pi)\ln(M^2/\mu^2)$$

$$\rightarrow \alpha(Q^2) = \frac{\alpha_0}{\alpha_0/\alpha(\mu^2) + (\alpha_0/3\pi)\ln(\mu^2/Q^2)} = \frac{\alpha(\mu^2)}{1 - [\alpha(\mu^2)/3\pi]\ln(Q^2/\mu^2)}$$

Very interesting result: Running α depends on q^2 , through its own *measured* value at any chosen energy scale μ^2 .

Cutoff has disappeared.


Running Coupling: QED - VII

57

Deep physics involved:

A ∞ number of diagrams can be formally replaced by a single, 1-photon diagram where the coupling 'constant' is running with q^2


Result valid to the 'leading log' approximation




Running Coupling: QED - VIII

58


Virtual (loops) e^+e^- pairs



Effective shielding



@TBA



Picture the QED perturbative vacuum as a sort of (virtual) dielectric medium: Virtual photons and e^+e^- pairs continuously created/annihilated

Bare charge is shielded at large distance by the virtual pairs coming from loops.

The standard e charge is smaller than the bare charge

By probing the electron at smaller and smaller distance, observe an *increasing* effective charge

Running α at LEP - I

59



Experimental method: Bhabha scattering

δ_γ, δ_Z s -channel contributions (small)

ε radiative corrections (known)

Use accurate, differential cross-section measurement to unfold $\alpha(t)$

[Total cross-section measurement would require a luminosity]





Running α at LEP - II

60

Just as an example, take L3 at LEP:
Relying on Bhabha scattering at small angle

$$\sigma = \frac{16\pi\alpha^2}{s} \left(\frac{1}{\theta_{min}^2} - \frac{1}{\theta_{max}^2} \right)$$


Calorimeter (Energy) + Silicon Tracker (Angle + Acceptance)




Running Coupling: QCD - I

61

Repeat all the steps: Loops etc



Except this time one has more loops: Gluons



Running Coupling: QCD - II

62

Turns out gluon loops yield *anti-shielding* effect
With 8 gluons and 6 quark flavors, gluons win

$$\alpha_s(|q^2|) = \frac{\alpha_s(\mu^2)}{1 + (\alpha_s(\mu^2)/12\pi)(33 - 2n_{flavor}) \ln(|q^2|/\mu^2)}$$

Running coupling *decreases* with increasing q^2 (or at smaller distance)
This is known as *asymptotic freedom*:

Large q^2 processes feature small coupling \rightarrow Perturbative!

Most important consequence:

The fundamental hypothesis behind the successful parton model is finally understood and justified

Running Coupling: QCD - III

63

Rather than making reference to a specific value of α_s

$$\alpha_s(|q^2|) = \frac{\alpha_s(\mu^2)}{1 + (\alpha_s(\mu^2)/12\pi)(33 - 2n_{flavor}) \ln(|q^2|/\mu^2)}$$

define a new constant

$$\begin{aligned} \ln \Lambda^2 &= \ln \mu^2 - \frac{12\pi}{(33 - 2n_{flavor})\alpha_s(\mu^2)} \rightarrow \Lambda^2 = \mu^2 e^{-\frac{12\pi}{(33 - 2n_{flavor})\alpha_s(\mu^2)}} \\ \rightarrow \alpha_s(|q^2|) &\simeq \frac{12\pi}{(33 - 2n_{flavor}) \ln(|q^2|/\Lambda^2)} = \frac{12\pi}{21 \ln(|q^2|/\Lambda^2)}, \quad |q^2| \gg \Lambda^2 \end{aligned}$$

Λ = Renormalization scale \rightarrow Fixes α_s at all q^2


$\Lambda \approx 200 \text{ MeV}$ yields the correct α_s at $\mu^2 = M_{Z^0}^2$

Funny behavior, known as 'Dimensional Transmutation':

From an adimensional constant to a dimensional one $\alpha_s \rightarrow \Lambda$

Running Coupling: QCD - IV

64



Sources:

Jets

DIS

Quarkonium

Annihilation: Muons vs Quarks

65

Electron-Positron annihilation:
Electroweak process

Low energy ($E_{CM} \ll M_{Z^0}$): Mostly electromagnetic

High energy ($E_{CM} \sim M_{Z^0}$): Mostly neutral current

Final state: *Fermion / Antifermion* pair

Muon vs quark pairs

Best observed at e^+e^- colliders

Annihilation Cross-Section - I


66

Apply crossing symmetry to electron-muon scattering, take pure e.m. amplitude at tree level

$$e^- + \mu^- \rightarrow e^- + \mu^-$$

$$e^- + [e^-] \xrightarrow{\text{crossed}} [\mu^-] + \mu^- \equiv e^- + e^+ \rightarrow \mu^+ + \mu^- \quad \text{B: Annihilation}$$

A: Scattering



Amplitude for scattering:


$$T_{fi}(s, s', r, r') = (-e) \bar{u}_{(\mu)}(p_2', s') \gamma^\mu u_{(\mu)}(p_2, s) \frac{-ig_{\mu\nu}}{q^2} (-e) \bar{u}_{(e)}(p_1', r') \gamma^\nu u_{(e)}(p_1, r)$$

$$\begin{aligned} q &= p_1 - p_1' \rightarrow q^2 = (p_1 - p_1')^2 = p_1^2 + p_1'^2 - 2p_1 \cdot p_1' \\ q^2 &= 2m_e^2 - 2(E_1 E_1' - \mathbf{p}_1 \cdot \mathbf{p}_1') \underset{E \gg m}{\simeq} -2(E_1 E_1' - \mathbf{p}_1 \cdot \mathbf{p}_1') < 0 \end{aligned} \quad q = 4\text{-momentum transfer}$$

Amplitude for annihilation:


$$T_{fi}(s, s', r, r') = (-e) \bar{u}_{(\mu)}(p_2', s') \gamma^\mu v_{(\mu)}(p_1', r') \frac{-ig_{\mu\nu}}{q^2} (-e) \bar{v}_{(e)}(p_1, s) \gamma^\nu u_{(e)}(p_2, r)$$

$$\begin{aligned} q &= p_1 + p_2 \rightarrow q^2 = (p_1 + p_2)^2 = p_1^2 + p_2^2 + 2p_1 \cdot p_2 \\ q^2 &= 2m_e^2 + 2(E_1 E_2 - \mathbf{p}_1 \cdot \mathbf{p}_2) \underset{E \gg m}{\simeq} 2(E_1 E_2 - \mathbf{p}_1 \cdot \mathbf{p}_2) > 0 \end{aligned} \quad q = \text{total 4-momentum}$$



Annihilation Cross-Section - II

67



$$T_{fi} = \bar{v}(p_1, s') \gamma^\mu u(p_2, s) \frac{e^2}{q^2} \bar{u}(p_2', r) \gamma_\mu v(p_1', r')$$

$$|T_{fi}|^2 = \frac{e^4}{q^4} [\bar{v}(p_1, s') \gamma^\mu u(p_2, s) \bar{u}(p_2', s) \gamma^\nu v(p_1', s')] [\bar{v}(p_1, r) \gamma_\mu u(p_2, r) \bar{u}(p_2', r) \gamma_\nu v(p_1', r)]$$

Annihilation Cross-Section - III

68

$$|T_{fi}|^2 = \frac{e^4}{q^4} [\bar{v}(p_1, s') \gamma^\mu u(p_2, s) \bar{u}(p_2', s) \gamma^\nu v(p_1', s')] [\bar{v}(p_1, r) \gamma_\mu u(p_2, r') \bar{u}(p_2', r') \gamma_\nu v(p_1', r)]$$

$$\langle |T_{fi}|^2 \rangle = \frac{1}{4} \sum_{s,s',r,r'} |T_{fi}|^2 = \frac{e^4}{4q^4} Tr[(\not{p}_1 - m) \gamma^\mu (\not{p}_2 + m) \gamma^\nu] Tr[(\not{p}_2' + M) \gamma_\mu (\not{p}_1' - M) \gamma_\nu]$$

$$Tr[(\not{p}_1 - m) \gamma^\mu (\not{p}_2 + m) \gamma^\nu] = 4 [p_1^\mu p_2^\nu + p_1^\nu p_2^\mu - g^{\mu\nu} (p_2 \cdot p_1 + m^2)]$$

$$Tr[(\not{p}_2' + M) \gamma_\mu (\not{p}_1' - M) \gamma_\nu] = 4 [p_1'^\mu p_2'^\nu + p_1'^\nu p_2'^\mu - g^{\mu\nu} (p_2' \cdot p_1' + M^2)]$$

$$\rightarrow \langle |T_{fi}|^2 \rangle = \frac{8e^4}{q^4} [(p_1 \cdot p_1') (p_2 \cdot p_2') + (p_1 \cdot p_2') (p_2 \cdot p_1') + M^2 (p_1 \cdot p_2)] \quad m \approx 0$$

$$CM : \begin{cases} p_1 = (E \ 0 \ 0 \ E), p_2 = (E \ 0 \ 0 \ -E) \\ |\mathbf{p}'| = |\mathbf{p}_1'| = |\mathbf{p}_2'| = \sqrt{E^2 - M^2} \end{cases}$$

$$p_1' = (E \ |\mathbf{p}'| \sin \theta \ 0 \ |\mathbf{p}'| \cos \theta), p_2' = (E \ -|\mathbf{p}'| \sin \theta \ 0 \ -|\mathbf{p}'| \cos \theta)$$

$$\rightarrow \langle |T_{fi}|^2 \rangle = e^4 \left[1 + \frac{M^2}{E^2} + \left(1 - \frac{M^2}{E^2} \right) \cos^2 \theta \right]$$

Annihilation Cross-Section - IV

69

$$\frac{d\sigma}{d\Omega} \Big|_{CM} = \frac{\alpha^2}{4s} \sqrt{1 - \frac{M^2}{E^2}} \left[1 + \frac{M^2}{E^2} + \left(1 - \frac{M^2}{E^2} \right) \cos^2 \theta \right]$$


$$\rightarrow \frac{d\sigma}{d\Omega} \Big|_{CM} \approx \frac{\alpha^2}{4s} (1 + \cos^2 \theta), \quad E \gg M$$

$$\sigma = \int_{\Omega} \frac{d\sigma}{d\Omega} d\Omega = \frac{4\pi\alpha^2}{3s} \sqrt{1 - \frac{M^2}{E^2}} \left(1 + \frac{1}{2} \frac{M^2}{E^2} \right)$$

$$\gamma = \frac{E}{M} = \frac{1}{\sqrt{1 - \beta^2}}$$

$$\rightarrow \sigma = \frac{4\pi\alpha^2}{3s} \underbrace{\sqrt{\frac{\gamma^2 - 1}{\gamma^2}}}_{\beta} \left(\frac{1 + 2\gamma^2}{2\gamma^2} \right) = \frac{4\pi\alpha^2}{3s} \beta \left(\frac{1}{2\gamma^2} + 1 \right) = \frac{4\pi\alpha^2}{3s} \beta \left(\frac{3 - \beta^2}{2} \right)$$

$$\rightarrow \sigma \approx \frac{4\pi\alpha^2}{3s} = \frac{87}{s [GeV^2]} nb, \quad E \gg M$$



Annihilation Cross-Section - V

70





Figure 12b)




τ lepton discovery, mass & spin determination:

$$R_{ex}^{2p} \simeq \frac{\sigma(\tau^+ \tau^-)}{\sigma(\mu^+ \mu^-)} \simeq \sqrt{1 - \frac{M_\tau^2}{E^2}} \left(1 + \frac{1}{2} \frac{M_\tau^2}{E^2} \right), M_\mu \approx 0$$

Annihilation Cross-Section - VI

71



$\mu^+ \mu^-$ event: L3 detector at LEP

Annihilation Cross-Section - VII

72

Total cross-section vs s :
Low energy






Fig. 3

ADONE ('70s)

Annihilation Cross-Section - VIII

73

Total cross-section vs s :
Higher energy



Annihilation Cross-Section - IX

74

Angular distribution: Low energy
1-photon, forward/backward symmetric





Fig. 15. MARK II $e^+e^- \rightarrow \mu^+\mu^-$ at $\langle E_{\text{miss}} \rangle^2 = 5.847$ compared to $1 + \cos^2 \theta$.

SPEAR ('70s)

Annihilation Cross-Section - X

75

Angular distribution: Higher energy:
Some contribution from Z^0 , forward/backward asymmetric



Annihilation Cross-Section - XI

76

Forward/Backward asymmetry: Important subject

Effective tool for precision tests of SM

May probe physics BSM

Interesting point:

Some tiny asymmetry expected from pure QED

Coming from diagrams with >1 photon (Radiative correction)

Dominated by interference terms


between amplitudes (a) and (b)-(g):

Opposite charge parity

Recent surge of interest from large asymmetry

found at Tevatron in $t\bar{t}$ production by 1 and 2 gluons:

Similar physics, not fully understood






PQCD: Jets in e^+e^- Collisions - I

77

e^+e^- annihilation into hadrons:

At the parton level = Crossed Deep Inelastic Scattering




Understood as annihilation into a $q\bar{q}$ pair,
followed by quark fragmentation into hadrons

PQCD: Jets in e^+e^- Collisions - II

78

Picture of quark fragmentation



PQCD: Jets in e^+e^- Collisions - III

79

By ignoring *quark fragmentation* details

$$e^+ + e^- \rightarrow q + \bar{q} \rightarrow \text{hadrons}$$

$$\frac{d\sigma}{d\Omega} = \frac{3\alpha^2}{4s} (1 + \cos^2 \theta) \sum_{flavor} e_{flavor}^2$$

$$\rightarrow \sigma(s) = \frac{4\pi\alpha^2}{s} \sum_{flavor} e_{flavor}^2$$

Due to color confinement, quark/antiquark unobservable as free particles

Expect hadron debris from quark fragmentation

→ *Jets*


Hadrons: Trend towards grouping into narrow pencils of particles

Naive prediction: 2 jets

At high energy, expect hadronic annihilation events to be *non-spherical*

PQCD: Jets in e^+e^- Collisions - IV

80



Define *sphericity* of events:

$$S = \min \frac{3}{2} \frac{\sum_i p_{\perp i}^2}{\sum_i p_i^2} \rightarrow \begin{cases} 0 & \text{2-jets} \\ 1 & \text{spherical} \end{cases}$$

min : Choose axes which minimize S (\leftarrow Iterative)


PQCD: Jets in e^+e^- Collisions - V

81


Interesting observable: *Thrust*

$$T = \frac{1}{Q} \max_{\vec{n}} \sum_i |\vec{n} \cdot \vec{p}_i|$$
$$\tau = 1 - T$$


$\tau = 0$ $\tau = \frac{1}{3}$



$$\tau = 1 - \frac{1}{\sqrt{3}} = 0.42$$




$$\tau = 0.48$$



$$\tau = \frac{M_1^2 + M_2^2}{Q^2}$$

PQCD: Jets in e^+e^- Collisions - VI


82



ALEPH

PQCD: Jets in $e^+ e^-$ Collisions - VII

83




PQCD: Jets in $e^+ e^-$ Collisions - VIII

84


For 2 jets events

$$\left. \begin{aligned} \frac{d\sigma}{d\Omega} &\propto 1 + \cos^2 \theta & \text{quark spin} &= 1/2 \\ \frac{d\sigma}{d\Omega} &\propto 1 - \cos^2 \theta & \text{quark spin} &= 0 \end{aligned} \right\} \equiv 1 + \alpha \cos^2 \theta$$



@TBA

Mark I (SPEAR)
 $E = \text{few GeV}$



ALEPH (LEP)
 $E = 90 \text{ GeV}$


PQCD: Jets in e^+e^- Collisions - IX

85



PQCD: Jets in e^+e^- Collisions - X

86



PQCD: Jets in $e^+ e^-$ Collisions - XI

87

Total hadronic cross section $\leftrightarrow R$ Ratio

Reminder:

Time scale of hard interaction

$$T \sim \frac{1}{Q} = \frac{1}{\sqrt{s}} \sim \frac{1}{\text{Many GeV}} \rightarrow \text{Very small}$$

Time scale of soft hadronization

$$\tau \sim \frac{1}{\Lambda} \sim \frac{1}{1 \text{ GeV}} \rightarrow \text{Large}$$

\rightarrow Perturbative cross section OK

Expect at 30-40 GeV:

$$R = \frac{\sigma_{q\bar{q}}}{\sigma_{\mu\mu}} = 3 \sum_{i=u,d,s,c,b} Q_i^2 = \frac{11}{3} \approx 3.67 (+ 0.05 \text{ coming from } Z^0)$$

Measure :


$$R \approx 3.9$$

$\rightarrow QCD$ Correction required


PQCD: Jets in $e^+ e^-$ Collisions - XII

88

QCD corrections Next to Leading Order (NLO):



Virtual gluons



Real gluons

Real gluons: 3 particles in the final state

Some kinematics:

$$x_1 = \frac{2E_1}{\sqrt{s}}, x_2 = \frac{2E_2}{\sqrt{s}}$$

$$\rightarrow 0 \leq x_1, x_2 \leq 1, x_1 + x_2 \geq 1$$

$$x_3 = \frac{2E_g}{\sqrt{s}} = 2 - x_1 - x_2$$

PQCD: Jets in $e^+ e^-$ Collisions - XIII

89

Observe:

Plane (2D) event

Within the event plane: 2 degrees of freedom

Differential cross section:

$$\frac{1}{\sigma} \frac{d^2\sigma}{dx_1 dx_2} = \frac{2\alpha_s}{3\pi} \frac{x_1^2 + x_2^2}{(1-x_1)(1-x_2)}$$

Basic remark:

$$x_1, x_2 \rightarrow 1 \quad \rightarrow \quad \sigma \rightarrow \infty$$

Also true to higher perturbative orders

\rightarrow 2 jets dominant over everything else

PQCD: Jets in $e^+ e^-$ Collisions - XIV

90

Total hadronic cross section:

$$\sigma^{q\bar{q}g} = \sigma_0 3 \sum_q Q_q^2 \int \frac{d^2\sigma}{dx_1 dx_2} dx_1 dx_2 = \sigma_0 3 \sum_q Q_q^2 \int \frac{2\alpha_s}{3\pi} \frac{x_1^2 + x_2^2}{(1-x_1)(1-x_2)} dx_1 dx_2$$

In order to regularize diverging integrals: Funny and smart idea

Shift to $4-2\varepsilon$ space-time dimensions, make them nicely converging..

Diagrams with real gluons:

$$\sigma^{q\bar{q}g} = \sigma_0 3 \sum_q Q_q^2 \frac{2\alpha_s}{3\pi} H(\varepsilon) \left[\frac{2}{\varepsilon^2} - \frac{3}{\varepsilon} + \frac{19}{2} + O(\varepsilon) \right]$$

Diagrams with virtual gluons:

$$\sigma^{q\bar{q}g} = \sigma_0 3 \sum_q Q_q^2 \frac{2\alpha_s}{3\pi} H(\varepsilon) \left[-\frac{2}{\varepsilon^2} + \frac{3}{\varepsilon} - 8 + O(\varepsilon) \right]$$

Adding everything up, and reverting to 4D:

$$R \xrightarrow[\varepsilon \rightarrow 0]{} 3 \sum_q Q_q^2 \left\{ 1 + \frac{\alpha_s}{\pi} + O(\alpha_s^2) \right\} \simeq 3 \cdot \frac{11}{9} \left(1 + \frac{0.14}{3.14} \right) = 3.83$$

$$R + R_{Z^0} \simeq 3.83 + 0.05 = 3.88 !!!$$

PQCD: Jets in $e^+ e^-$ Collisions - XV

91

Absolute number of jets ill defined, both in theory and experiment

Can define number of jets as a function of some *resolution parameter*, according to a *clustering algorithm*

Example : *Durham algorithm*

Take $q\bar{q}g$ final state

By fixing a y parameter as

$$m_{thresh}^2 = ys$$

compare the (invariant mass)² of each parton pair to m_{thresh}^2

$$(p_i + p_j)^2 > ys \quad i, j = q, \bar{q}, g \quad 3 \text{ comb/event}$$

Then:

$$N_{hit} = N_{jet}$$

Of course,

$$\begin{aligned} R_{2jet} &= R_{2jet}(y) \\ R_{3jet} &= R_{3jet}(y) \end{aligned}$$

Extend to n partons \rightarrow QCD predicts $R_{k-jet}(y)!$


PQCD: Jets in $e^+ e^-$ Collisions - XVI

92

Jet algorithm for data: (modified) *Durham*

Define y_{cut}

Loop over all particle pairs




$$y_{ij} = \frac{2E_i E_j (1 - \cos \theta_{ij})}{s} \simeq \frac{m_{ij}^2}{s}$$

$$p_i = (E_i, \vec{p}_i)$$
$$\theta_{ij}$$
$$p_j = (E_j, \vec{p}_j)$$

Exit with 4-momenta of n jets

PQCD: Jets in $e^+ e^-$ Collisions - XVII


93




PQCD: Jets in $e^+ e^-$ Collisions - XVIII

94

Exceptional 3-jet event from OPAL



@TBA



PQCD: Jets in e^+e^- Collisions - XIX



95

Get a measurement of α_s :

$$\alpha_s(35\text{GeV}) = 0.146 \pm 0.03$$

$$\alpha_s(M_{Z^0}) = 0.124 \pm 0.0043$$

$$R_3 \equiv \frac{\sigma_{3\text{-jet}}}{\sigma_{\text{tot}}} \propto \alpha_s(E_{\text{cm}}) \propto \frac{1}{\ln E_{\text{cm}}}$$




PQCD: Jets in e^+e^- Collisions- XX

96


Is QCD really $SU(3)$?

Test for non-Abelian couplings at LEP: 4 jets events


Special angular correlation from 3-gluon vertex amplitude



(a)



(b)





(c)

PQCD: Jets in $e^+ e^-$ Collisions - XXI

97

Look at distribution of a special angle, sensitive to non-Abelian couplings:



Quark Parton Model - I

98

Write down F_2 in terms of PDFs

$$\frac{F_2(x)}{x} = \sum_i z_i^2 \underbrace{n_i \delta\left(x - \frac{m_i}{M}\right)}_{\text{PDFs}} \rightarrow \frac{F_2(x)}{x} = \sum_i z_i^2 \underbrace{q_i(x)}_{\text{PDFs}}$$

$p = uud$

$$F_2^p(x) = x \left[\left(\frac{2}{3}\right)^2 u_p(x) + \left(-\frac{1}{3}\right)^2 d_p(x) \right]$$

$$\rightarrow F_2^p(x) = x \left[\frac{4}{9} u_p(x) + \frac{1}{9} d_p(x) \right]$$

$n = ddu$

$$F_2^n(x) = x \left[\left(-\frac{1}{3}\right)^2 d_n(x) + \left(\frac{2}{3}\right)^2 u_n(x) \right]$$

From isospin symmetry:

$$F_2^n(x) = x \left[\left(-\frac{1}{3}\right)^2 u_p(x) + \left(\frac{2}{3}\right)^2 d_p(x) \right]$$

$$\rightarrow F_2^n(x) = x \left[\frac{1}{9} u_p(x) + \frac{4}{9} d_p(x) \right]$$

Quark Parton Model - II

99

Consider the deuteron structure function:

$$\begin{aligned} F_2^d(x) &= \frac{1}{2}(F_2^p + F_2^n) = \frac{5}{9}\frac{x}{2}[u_p(x) + d_p(x)] \\ \rightarrow F_2^n(x) &= F_2^d(x) - F_2^p(x) \\ &= \frac{5}{18}x[u_p(x) + d_p(x)] - \frac{1}{9}x[u_p(x) - 4d_p(x)] \\ &= \frac{3}{18}x[u_p(x) - d_p(x)] \end{aligned}$$

Finally extract PDFs from measured F_2

$$xu_p(x) = xd_n(x) = 3F_2^p(x) - \frac{6}{5}F_2^d(x)$$

$$xu_n(x) = xd_p(x) = 3F_2^p(x) + \frac{24}{5}F_2^d(x)$$

Quark Parton Model - III

100

Take a Hydrogen atom:

Common wisdom: “A bound state of proton + electron”

But: Consider the effect of radiative corrections (e.g. loops)

Then we should be more precise:

Hydrogen = (Proton+Electron)_{Valence} + (Positrons+Electrons+Photons)_{Sea}

Can we say valence and sea particles are fundamentally different? Well,...

In a bound state, both are off mass shell

Both are active in yielding measurable effects (Coulomb levels vs. Lamb shift,...)

Sea particles yield small corrections to levels determined by valence e+p


Take a hadron:

Hadron = (Quarks/Antiquarks)_{Valence} + (Quarks/Antiquarks+Gluons)_{Sea}

Since $a_s \gg a$, **sea effects are much larger in QCD**

Quark Parton Model - IV

101



Among parton model predictions: *Sum Rules* (= Integral relations) for PDFs
 Examples: Proton quark content is *uud*

$$\int [u_p(x) - \bar{u}_p(x)] dx = 2$$


$$\int [d_p(x) - \bar{d}_p(x)] dx = 1$$

$$\int [s_p(x) - \bar{s}_p(x)] dx = 0$$


PQCD: DIS Scaling Violations - I

102

Our picture of structure functions



Observe small deviations from scaling: $F_2(x) \rightarrow F_2(x, Q^2)$



PQCD: DIS Scaling Violations - II

103

QCD on $F_2(x, Q^2)$:

x -dependence \rightarrow Non perturbative \rightarrow Not predicted

Q^2 -dependence \rightarrow Perturbative \rightarrow Predicted !




Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) Equations:

Successful prediction of Q^2 evolution of structure functions


PQCD: DIS Scaling Violations - III

104

First order (NLO) QCD corrections to naive Quark Parton Model:



QPM



QCD

PQCD: DIS Scaling Violations - IV

105

The bottom line:

Measured structure functions at any given Bjorken x depend on quark & gluon densities taken at higher fractional momentum $y>x$

This originates a slow Q^2 dependence

Core physics: Probabilities of QCD radiative/scattering processes $P_{qq}(x/y)$, $P_{gq}(x/y)$
Usually called *splitting functions*

$$\frac{F_2(x)}{x} = 2F_1(x) = \sum_i e_i^2 q_i(x) \quad \text{Quark-Parton Model}$$

$$\rightarrow \frac{F_2(x)}{x} = \sum_i e_i^2 \int_0^1 q_i(y) \delta(x-y) dy \quad \text{Reminder:}$$


$$\begin{aligned} \rightarrow \frac{F_2(x)}{x} &= \sum_i e_i^2 \int_0^1 q_i(y) \delta\left(1 - \frac{x}{y}\right) \frac{dy}{y} \\ z &= \frac{x}{y} \\ &\rightarrow \delta(x-y) = \delta\left[y\left(1 - \frac{x}{y}\right)\right] = \frac{1}{y} \delta\left(1 - \frac{x}{y}\right) \end{aligned}$$

$$\rightarrow \frac{F_2(x)}{x} = \sum_i e_i^2 \int_0^1 q_i(y) \left[\delta\left(1 - \frac{x}{y}\right) + \sigma_{qq}(z) \right] \frac{dy}{y} \quad \text{QCD corrections}$$

PQCD: DIS Scaling Violations - V


Just as an example: Gluon radiation splitting function at leading order (LO)

Almost carbon-copy of Compton effect



$$k \leftrightarrow q \\ k' \leftrightarrow p' \\ u = (k-p')^2 \\ t = (q-p')^2$$

$$\gamma^*(k) q(p) \rightarrow \gamma^*(k') q(p')$$




$$\gamma^*(q) q(p) \rightarrow q(p') g(k)$$

$$\left(\frac{d\sigma}{d\Omega} \right)_{\gamma q \rightarrow \gamma q} = \frac{\alpha^2 e_q^2}{2s} \left(\frac{-u}{s} - \frac{s}{u} - \frac{2tq^2}{su} \right)$$

$$\left(\frac{d\sigma}{d\Omega} \right)_{qg \rightarrow gg} = \frac{C_F \alpha \alpha_s e_q^2}{2s} \left(\frac{-t}{s} - \frac{s}{t} - \frac{2uq^2}{st} \right)$$

PQCD: DIS Scaling Violations - VI



$$dp_T^2 = d(p^2 \sin^2 \theta) = 2p^2 \sin \theta \underbrace{\cos \theta}_{\approx 1} d\theta = 2p^2 d(\cos \theta)$$

$$\rightarrow dp_T^2 = \frac{s}{2} d(\cos \theta) = \frac{s}{2} \frac{d\Omega}{4\pi} \rightarrow \frac{d\sigma}{d\Omega} = \frac{s}{8\pi} \frac{d\sigma}{dp_T^2} \rightarrow \frac{d\sigma}{dp_T^2} = \frac{8\pi}{s} \frac{d\sigma}{d\Omega}$$

Reminder (Bjorken): $x = -\frac{q^2}{2P \cdot q}$

Define: $z = -\frac{q^2}{2p \cdot q} = -\frac{q^2}{s - q^2}$

$$\rightarrow \left(\frac{d\sigma}{dp_T^2} \right)_{\gamma q \rightarrow gq} \propto \frac{C_F \alpha \alpha_s e_q^2}{p_T^2} P_{qq}(z), \quad P_{qq}(z) \equiv \frac{1+z^2}{1-z^2}$$

PQCD: DIS Scaling Violations - VII

108

Integrate 'Compton-like' differential cross-section between:

$$\begin{cases} \lambda \text{ lower cutoff } (\leftarrow \text{no divergences}) \\ \frac{\hat{s}}{4} \text{ upper cutoff } (\leftarrow \text{kinematical}), \hat{s} \text{ partonic CM energy squared} \end{cases}$$

$$\sigma_{qq}(z) = \int_{\lambda}^{\hat{s}/4} \frac{d\sigma}{dp_T^2} dp_T^2 \propto \frac{C_F \alpha \alpha_S e_q^2}{s} P_{qq}(z) \ln\left(-\frac{q^2}{\lambda}\right)$$

$$\text{Redefine : } P_{qq}(z) \equiv \frac{\alpha e_q^2 C_F}{2\pi s} \frac{1+z^2}{1-z^2}$$

$$\rightarrow \frac{F_2(x)}{x} = \sum_i e_i^2 \int_x^1 q_i(y) \left[\delta(1-\frac{x}{y}) + \frac{\alpha_s}{2\pi} P_{qq}(z) \ln\left(\frac{Q^2}{\lambda}\right) \right] \frac{dy}{y}$$

$$\rightarrow \frac{F_2(x)}{x} = \sum_i e_i^2 \underbrace{\left[q_i(x) + \int_x^1 q_i(y) \frac{\alpha_s}{2\pi} \ln\left(\frac{Q^2}{\lambda}\right) P_{qq}\left(\frac{x}{y}\right) \frac{dy}{y} \right]}_{q_i(x, Q^2)}$$

$$q_i(x, Q^2) = q_i(x) + \int_x^1 q_i(y) \frac{\alpha_s}{2\pi} \ln\left(\frac{Q^2}{\lambda}\right) P_{qq}\left(\frac{x}{y}\right) \frac{dy}{y}$$

PQCD: DIS Scaling Violations -VIII

109

Evolution equation for each quark flavor:

$$\frac{dq_i(x, Q^2)}{d \ln(Q^2)} = \int_x^1 q_i(y, Q^2) \frac{\alpha_s}{2\pi} P_{qq} \left(\frac{x}{y} \right) \frac{dy}{y}$$

Observe: Since $q_i(x) \rightarrow q_i(x, Q^2)$, the evolution equation should involve $q_i(x, Q^2)$, rather than $q_i(x)$, under the integral symbol

This is actually incomplete:

We forgot to add the contribution from the 2nd diagram..

$$\rightarrow \frac{dq(x, Q^2)}{d \ln(Q^2)} = \frac{\alpha_s(Q^2)}{2\pi} \int_x^1 \left[q(y, Q^2) P_{qq} \left(\frac{x}{y} \right) + g(y, Q^2) P_{gq} \left(\frac{x}{y} \right) \right] \frac{dy}{y}$$



And there is another equation for the evolution of the *gluon* density:

$$\frac{dg(x, Q^2)}{d \ln(Q^2)} = \frac{\alpha_s(Q^2)}{2\pi} \int_x^1 \left[\sum_i q_i(y, Q^2) P_{qg} \left(\frac{x}{y} \right) + g(y, Q^2) P_{gg} \left(\frac{x}{y} \right) \right] \frac{dy}{y}$$

Altarelli - Parisi, or *DGLAP*, equations for the parton densities

PQCD: DIS Scaling Violations -IX


110



PQCD: DIS Scaling Violations -X


111

PDF Evolution with Q^2



PQCD: Drell-Yan - I

112




Angular distribution in the pair rest frame
Expect $\propto 1 + \cos^2 \theta^*$ as usual for Fermion-Antifermion

$$\frac{d\sigma(q\bar{q} \rightarrow l^+l^-)}{dq^2} = \frac{4\pi\alpha^2}{3q^2} e_q^2 \delta(q^2 - s_{q\bar{q}})$$

x_1, x_2 Bjorken x for q, \bar{q}

$$s_{q\bar{q}} = (p_q + p_{\bar{q}})^2 = (x_1 p_1 + x_2 p_2)^2 \simeq x_1 x_2 s$$

$$\sigma(q\bar{q} \rightarrow l^+l^-) = \frac{4\pi\alpha^2}{3q^2} e_q^2, \quad e_q = \text{quark charge in } e \text{ units}$$



@TBA


PQCD: Drell – Yan - II

113

Reverse $e^+e^- \rightarrow q\bar{q}$ process: $q\bar{q} \rightarrow e^+e^-$

Obtained by any reaction

hadron + hadron $\rightarrow l^+ l^- + X$



Ignore parton transverse momentum

Ignore non-annihilating partons (\rightarrow "spectators")

Ignore parton fragmentation

PQCD: Drell – Yan - III

114

$e^+e^- \rightarrow \mu^+\mu^-$:

$$\frac{d\sigma}{d\Omega^*} = \frac{\alpha^2}{4s} (1 + \cos^2 \theta^*)$$

$$\sigma = \frac{4\pi\alpha^2}{3s}$$

$q\bar{q} \rightarrow \mu^+\mu^-$:

$$\frac{d\sigma_q}{d\Omega^*} = \frac{Q_q^2 \alpha^2}{4M^2} (1 + \cos^2 \theta^*) \cdot \frac{1}{3}$$

$Q_q e$: Quark charge

$\frac{1}{3}$: Color factor

M^2 : $\mu^+\mu^-$ invariant mass = Total energy in partonic CM

PQCD: Drell – Yan - IV

115

$$p_1 = x_1 P_1 \simeq x_1 (E, 0, 0, P)$$

$$p_2 = x_2 P_2 \simeq x_2 (E, 0, 0, -P)$$

$$q = x_1 P_1 + x_2 P_2 \simeq [(x_1 + x_2) E, 0, 0, (x_1 - x_2) P]$$

$$M^2 = q^2 \simeq (x_1 + x_2)^2 E^2 - (x_1 - x_2)^2 P^2 \simeq 4x_1 x_2 E^2 = x_1 x_2 s \equiv \tau s$$

Switch to more useful kinematical variables:

Either

$$\begin{cases} x_F \equiv x_1 - x_2 = \pm \frac{2|\mathbf{q}|}{\sqrt{s}} & \text{Feynman } x \text{ of parton pair} \\ M^2 = sx_1 x_2 \end{cases}$$

Or:

$$\begin{cases} y = \frac{1}{2} \ln \frac{E + p_3}{E - p_3} \simeq \frac{1}{2} \ln \frac{x_1}{x_2} & \text{Rapidity of parton pair (neglecting rest mass)} \\ M = \sqrt{sx_1 x_2} \end{cases}$$

PQCD: Drell – Yan - V

116

Inclusive cross-section:

Contribution by parton pair with (x_1, x_2) fractional momenta

$$d^2\sigma(pp \rightarrow \mu^+ \mu^- + X) = \frac{4\pi\alpha^2}{9M^2} \sum_q Q_q^2 [f_q(x_1) f_{\bar{q}}(x_2) + f_q(x_2) f_{\bar{q}}(x_1)] dx_1 dx_2$$

$$\begin{cases} y = \frac{1}{2} \ln \frac{E + p_3}{E - p_3} \simeq \frac{1}{2} \ln \frac{x_1}{x_2} \rightarrow x_1 = x_2 e^{2y} \\ M = \sqrt{s x_1 x_2} \end{cases} \rightarrow M = \sqrt{s} x_2 e^y \rightarrow x_2 = \frac{M}{\sqrt{s}} e^{-y}, x_1 = \frac{M}{\sqrt{s}} e^{+y}$$

$$dx_1 dx_2 = J dM dy$$

$$J = \frac{\partial(x_1, x_2)}{\partial(M, y)} = \begin{vmatrix} \frac{\partial x_1}{\partial M} & \frac{\partial x_1}{\partial y} \\ \frac{\partial x_2}{\partial M} & \frac{\partial x_2}{\partial y} \end{vmatrix} = \frac{\partial x_1}{\partial M} \frac{\partial x_2}{\partial y} - \frac{\partial x_1}{\partial y} \frac{\partial x_2}{\partial M}$$

$$\rightarrow J = \frac{1}{\sqrt{s}} e^{+y} \left(-\frac{M}{\sqrt{s}} e^{-y} \right) - \frac{M}{\sqrt{s}} e^{+y} \frac{1}{\sqrt{s}} e^{-y} = -\frac{2M}{s} = -2\sqrt{\frac{x_1 x_2}{s}} \rightarrow dx_1 dx_2 = \left(-2\sqrt{\frac{x_1 x_2}{s}} \right) dM dy$$

PQCD: Drell – Yan - VI

117

$$\rightarrow d^2\sigma = \frac{4\pi\alpha^2}{9M^2} \left| -2\sqrt{\frac{x_1 x_2}{s}} \right| \sum_q Q_q^2 [f_q(x_1) f_{\bar{q}}(x_2) + f_q(x_2) f_{\bar{q}}(x_1)] dM dy$$

$$s\tau = M^2 \rightarrow M = \sqrt{s\tau}$$

$$\rightarrow \frac{d^2\sigma}{dM dy} = \frac{8\pi\alpha^2}{9M^2} \frac{\tau}{M} \sum_q Q_q^2 [f_q(x_1) f_{\bar{q}}(x_2) + f_q(x_2) f_{\bar{q}}(x_1)]$$

$$\rightarrow \frac{d^2\sigma}{dM dy} = \frac{8\pi\alpha^2}{9Ms} \sum_q Q_q^2 [f_q(x_1) f_{\bar{q}}(x_2) + f_q(x_2) f_{\bar{q}}(x_1)]$$

Central events:

$$y = 0, x_1 = x_2 = \sqrt{\tau}$$

$$\rightarrow \frac{d^2\sigma}{dM dy} \Big|_{y=0} = \frac{8\pi\alpha^2}{9Ms} \sum_q Q_q^2 [f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau}) + f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau})]$$

$$\rightarrow s^{3/2} \frac{d^2\sigma}{dM dy} \Big|_{y=0} = \frac{8\pi\alpha^2}{9} \frac{1}{\sqrt{\tau}} \sum_q Q_q^2 [f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau}) + f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau})]$$

PQCD: Drell – Yan - VII

118

$$M = \sqrt{s\tau} \rightarrow dM = \sqrt{s}d(\sqrt{\tau})$$



$$\rightarrow s^{3/2} \frac{d^2\sigma}{dMdy} \Big|_{y=0} = s^{3/2} \frac{d^2\sigma}{\sqrt{s}d(\sqrt{\tau})dy} \Big|_{y=0} = s \frac{d^2\sigma}{d(\sqrt{\tau})dy} \Big|_{y=0}$$

$$\rightarrow s \frac{d^2\sigma}{d(\sqrt{\tau})dy} \Big|_{y=0} = \frac{8\pi\alpha^2}{9} \frac{1}{\sqrt{\tau}} \sum_q Q_q^2 [f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau}) + f_q(\sqrt{\tau}) f_{\bar{q}}(\sqrt{\tau})]$$

Scaling behavior: Compare to DIS

PQCD: Drell – Yan - VIII

119



PQCD: Drell – Yan - IX

120


NLO QCD corrections:



(a)



(b)



(c)

Quite similar to QCD corrections to:

$$e^+ e^- \rightarrow q\bar{q}$$

PQCD: Drell – Yan - X

121

Total rate:

Same effect as for

$$e^+ e^- \rightarrow q\bar{q}$$

Real gluons compensate virtual gluons

$$\sigma(\text{real}) + \sigma(\text{virtual}) = \frac{2\alpha_s}{3\pi} \sigma_0 \left(\frac{4\pi^2}{3} - \frac{7}{2} \right)$$


→ Overall effect lumped into a K -factor

$$K_{DY}^{(1)} = 1 + \frac{\alpha_s}{\pi} \left(\frac{8\pi^2}{9} - \frac{7}{3} \right) \approx 1 + 2.05\alpha_s \sim 2$$

PQCD: Drell – Yan - XI

122

DY Scaling violation



$$\text{Feynman } x: x_F = \frac{2}{\sqrt{s}}(p_+ + p_-) \approx x_1 - x_2$$

PQCD: Hadron Collisions - I

123

Historically best observed and studied at hadron colliders

ISR = Intersecting Storage Ring (CERN '70s)

pp 31 GeV / beam

S $p\bar{p}S$ = Super p \bar{p} Synchrotron (CERN '80s)

p \bar{p} 270 - 310 GeV / beam

Tevatron (Fermilab early '90s - 2011)

p \bar{p} 1 TeV / beam

RHIC = Relativistic Heavy Ion Collider (BNL 3rd Millennium)

ions 200 GeV / nucleon * beam

LHC = Large Hadron Collider (CERN 3rd Millennium)

pp 7 TeV / beam (presently 4 TeV)

ions 2.7 TeV / nucleon * beam

PQCD: Hadron Collisions - II

124

CM frame: usually identical to LAB

Important exception: ISR (collision angle 15^0)

Not relevant for LHC (collision angle 0.01^0)

But: Partonic collision CM \neq Event CM

→ E_{top} , p of parton collision unknown

→ Initial state only partially known

→ Separate collision kinematics into:

Transverse

Longitudinal

Introduce useful kinematical variables: *Rapidity*, *Transverse momentum*

PQCD: Hadron Collisions - III

125

Lorentz transformation $S \rightarrow S'(\beta)$:

$$\begin{pmatrix} z' \\ t' \end{pmatrix} = \begin{pmatrix} \gamma & -\gamma\beta \\ -\gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} z \\ t \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{1-\beta^2}} & -\frac{\beta}{\sqrt{1-\beta^2}} \\ -\frac{\beta}{\sqrt{1-\beta^2}} & \frac{1}{\sqrt{1-\beta^2}} \end{pmatrix} \begin{pmatrix} z \\ t \end{pmatrix}$$

Compare:

$$\gamma^2 - \gamma^2\beta^2 = \frac{1}{1-\beta^2} - \frac{\beta^2}{1-\beta^2} = 1 \Leftrightarrow \cosh^2 y - \sinh^2 y = 1$$

$$\rightarrow \begin{cases} \gamma = \cosh y \\ \beta\gamma = \sinh y \end{cases} \rightarrow \beta = \tanh y, \quad y \text{ rapidity}$$

$$\rightarrow \begin{pmatrix} z' \\ t' \end{pmatrix} = \begin{pmatrix} \cosh y & -\sinh y \\ -\sinh y & \cosh y \end{pmatrix} \begin{pmatrix} z \\ t \end{pmatrix}$$

PQCD: Hadron Collisions - IV

126

Most important properties:

Rapidity is *additive* under Lorentz boosts

Transverse momentum is *invariant* under Lorentz boosts

$$y = \frac{1}{2} \ln \frac{E + p_{\parallel}}{E - p_{\parallel}}$$

Boost along z by β, γ

$$\rightarrow y' = \frac{1}{2} \ln \frac{\gamma(E + \beta p_{\parallel}) + \gamma(p_{\parallel} + \beta E)}{\gamma(E + \beta p_{\parallel}) - \gamma(p_{\parallel} + \beta E)}$$

$$\rightarrow y' = \frac{1}{2} \ln \frac{(E + p_{\parallel})(1 + \beta)}{(E - p_{\parallel})(1 - \beta)} = \underbrace{\frac{1}{2} \ln \frac{(E + p_{\parallel})}{(E - p_{\parallel})}}_y + \underbrace{\frac{1}{2} \ln \frac{(1 + \beta)}{(1 - \beta)}}_{y_{boost}}$$

Indeed:

$$y \rightarrow y' = y + y_b$$

$$\rightarrow dy' = dy, \quad \Delta y' = \Delta y$$

PQCD: Hadron Collisions - V

127

Separate longitudinal/transverse momentum components:

$$E^2 = m^2 + |\mathbf{p}|^2$$

$$|\mathbf{p}|^2 = p_{\parallel}^2 + p_{\perp}^2$$

$$\rightarrow E^2 = m^2 + p_{\parallel}^2 + p_{\perp}^2 = m_{\perp}^2 + p_{\parallel}^2$$

$$\rightarrow \left(\frac{E}{m_{\perp}} \right)^2 - \left(\frac{p_{\parallel}}{m_{\perp}} \right)^2 = 1$$

$$\rightarrow \begin{cases} E = m_{\perp} \cosh y \\ p_{\parallel} = m_{\perp} \sinh y \end{cases}$$

$$\rightarrow p_{\parallel} = E \tanh y$$

$$\rightarrow \begin{pmatrix} p_{\parallel}' \\ E' \end{pmatrix} = \begin{pmatrix} \cosh y & -\sinh y \\ -\sinh y & \cosh y \end{pmatrix} \begin{pmatrix} p_{\parallel} \\ E \end{pmatrix}$$

PQCD: Hadron Collisions - VI

128

Rapidity:

$$p_{\parallel} = E \tanh y \leftrightarrow y = \frac{1}{2} \ln \frac{E + p_{\parallel}}{E - p_{\parallel}}$$

Consistency check:

For momentum along z

$$\begin{aligned} y &= \frac{1}{2} \ln \frac{E + p}{E - p} = \frac{1}{2} \ln \frac{\gamma m + \beta \gamma m}{\gamma m - \beta \gamma m} = \frac{1}{2} \ln \frac{1 + \beta}{1 - \beta} \\ &\rightarrow \frac{1 + \beta}{1 - \beta} = e^{2y} \rightarrow 1 + \beta = (1 - \beta)e^{2y} \rightarrow \beta(1 + e^{2y}) = e^{2y} - 1 \end{aligned}$$

$$\rightarrow \beta = \frac{e^{2y} - 1}{e^{2y} + 1} = \tanh y$$

$$\rightarrow \gamma = \cosh y$$

→ OK

PQCD: Hadron Collisions - VII


129

Pseudo-rapidity η :

$$y = \frac{1}{2} \ln \frac{E + p_{\parallel}}{E - p_{\parallel}} \approx \frac{1}{2} \ln \frac{1 + \cos \theta}{1 - \cos \theta} = -\frac{1}{2} \ln \frac{1 - \cos \theta}{1 + \cos \theta} = -\frac{1}{2} \ln \frac{\sin^2 \theta/2}{\cos^2 \theta/2}$$

$$\rightarrow y \approx -\frac{1}{2} \ln (\tan^2 \theta/2) = -\ln (\tan \theta/2) = \eta$$

1-to-1 correspondance to polar angle



PQCD: Hadron Collisions - VIII

130

Interesting processes: *Inclusive*

Ex: Inclusive production of $c \rightarrow a + b \rightarrow c + X$

→ Inclusive, invariant cross-sections

Reminder: $\frac{d^3\mathbf{P}}{E}$ Lorentz invariant quantity

Elementary volume (impulse space):

$$d^3\mathbf{P} = P^2 dP d\Omega = dP_{||} P_T dP_T d\varphi \rightarrow \frac{d^3\mathbf{P}}{E} = \frac{dP_{||} P_T dP_T d\varphi}{E}$$

$$dy = \frac{dP_{||}}{E} \rightarrow \frac{d^3\mathbf{P}}{E} = dy P_T dP_T d\varphi$$

$$\int (dy P_T dP_T) d\varphi = 2\pi dy P_T dP_T = 2\pi dy \frac{1}{2} d(P_T^2) = \pi dy d(P_T^2)$$

Inclusive, invariant differential cross-section:

$$\underbrace{\frac{d\sigma}{d^3\mathbf{P}}}_{\substack{\text{Lorentz} \\ \text{inv.}}} = E \frac{d\sigma}{d^3\mathbf{P}} = \frac{1}{\pi} \frac{d\sigma}{dy d(P_T^2)} = \frac{1}{2\pi P_T} \frac{d\sigma}{dy dP_T}$$

PQCD: Hadron Collisions - IX

131

$$\begin{pmatrix} E' \\ p_{\parallel}' \end{pmatrix} = \begin{pmatrix} \gamma & -\gamma\beta_{CM} \\ -\gamma\beta_{CM} & \gamma \end{pmatrix} \begin{pmatrix} E \\ p_{\parallel} \end{pmatrix} = \begin{pmatrix} \cosh y_{CM} & -\sinh y_{CM} \\ -\sinh y_{CM} & \cosh y_{CM} \end{pmatrix} \begin{pmatrix} E \\ p_{\parallel} \end{pmatrix}$$

$$p_T^2 = p_x^2 + p_y^2$$

$$p_T = p \sin \theta$$

$$E_T^2 = p_x^2 + p_y^2 + m^2 = p_T^2 + m^2 = E^2 - p_{\parallel}^2$$

$$p_{\parallel} = E \tanh y$$

$$E_T^2 = E^2 - p_{\parallel}^2 = E^2 - E^2 \tanh^2 y$$

$$\rightarrow E = E_T \cosh y$$

$$\rightarrow p_{\parallel} = E_T \sinh y$$

$$y \approx \eta = -\ln(\tan \theta/2) \rightarrow E_T = E(1 - \tanh^2 \eta)^{1/2} \approx E \left(1 - \frac{\frac{\cos \theta/2}{\sin \theta/2} - \frac{\sin \theta/2}{\cos \theta/2}}{\frac{\cos \theta/2}{\sin \theta/2} + \frac{\sin \theta/2}{\cos \theta/2}} \right)^{1/2}$$

$$E_T = E \left[1 - (\cos^2 \theta/2 - \sin^2 \theta/2) \right]^{1/2} \rightarrow E_T \approx E \sin \theta$$

PQCD: Hadron Collisions - X

132

Express 4-momentum in terms of longitudinal, transverse quantities

$$p = \left(E, \underbrace{P_x, P_y}_{P_T^2 = P_x^2 + P_y^2}, \underbrace{P_z}_{P_{\parallel} = P_z} \right)$$

$$P_T = \sqrt{P^2 - P_{\parallel}^2}$$

$$\rightarrow \begin{cases} P = P_T \cosh \eta \\ P_{\parallel} \approx P_T \sinh \eta \end{cases}$$

$$E \approx P, E_T \approx P_T$$

$$\rightarrow p \approx \left(E_T \cosh \eta, \underbrace{E_T \sin \phi, E_T \cos \phi}_{E_T}, E_T \sinh \eta \right)$$

Useful in clustering algorithms

PQCD: Hadron Collisions - XI

133

Partonic kinematics: Relevant for 'hard' collisions

Event CM frame:

$$\begin{cases} P_A = (E_A, 0, 0, p_A) \\ P_B = (E_A, 0, 0, -p_A) \end{cases} \quad \text{4-momenta incident particles}$$

$$\begin{cases} p_1 = x_1 P_A \\ p_2 = x_2 P_B \end{cases} \quad \text{4-momenta incident partons}$$

$$\beta_{CM} = \frac{x_1 - x_2}{x_1 + x_2} \quad \text{Parton CM speed as seen by event CM (= LAB for most colliders)}$$

$$y = \frac{1}{2} \ln \frac{1 + \beta_{CM}}{1 - \beta_{CM}} = \frac{1}{2} \ln \frac{x_1}{x_2} \quad \text{Parton CM rapidity as seen by event CM}$$

x_1, x_2 varying on event-by-event basis

→ y not fixed, statistically distributed

Distribution depending on event type

PQCD: Hadron Collisions - XII

134

→ *Hadronic, inclusive* cross sections for hard processes at any given partonic CM energy \hat{s}
result from a convolution of *partonic, exclusive* cross sections with pdf over full range of (x_1, x_2) :

$$d\sigma_{A+B \rightarrow f+X} = \int d\sigma_{ij \rightarrow f} f_1(x_1) f_2(x_2) dx_1 dx_2$$

$d\sigma_{ij \rightarrow f}$ depending on kinematical variables of the final state f

Partonic CM energy, Feynman x :

$$\sqrt{\hat{s}} = \sqrt{sx_1 x_2}$$

$$x_F = x_1 - x_2$$

Introduce τ, y :

$$\begin{cases} \tau = x_1 x_2 \\ y = \frac{1}{2} \ln \frac{x_1}{x_2} \end{cases} \rightarrow \begin{cases} x_1 = \sqrt{\tau} e^y \\ x_2 = \sqrt{\tau} e^{-y} \end{cases} \rightarrow dx_1 dx_2 = d\tau dy$$

$$\rightarrow \sqrt{\hat{s}} \equiv \sqrt{\tau s}$$

For any given τ : Bounds & Distribution of y

$$|y| < -\frac{1}{2} \ln \tau$$

$$\frac{1}{\sigma} \frac{d\sigma}{dy} = f_1(\sqrt{\tau} e^y) f_2(\sqrt{\tau} e^{-y})$$

PQCD: Hadron Collisions - XIII


135

Examples of rapidity distribution:

Produzione di top al Tevatron

$$q\bar{q} \rightarrow t\bar{t}$$

$$\tau = 4m_t^2 / s \approx 0.2^2$$




distribuzione
come $(1-x)^k$

Produzione di b a LHC

$$q\bar{q}, gg \rightarrow b\bar{b}$$

$$\tau = 4m_b^2 / s \approx (7 \times 10^{-4})^2$$



interazione più probabile:
un quark di valenza con uno del mare.

PQCD: Hadron Collisions - XIV

136

Introduce differential luminosity for parton collision occurring within $(\tau, \tau + d\tau)$:


$$\frac{dL}{d\tau} = \int dx_1 dx_2 f_1(x_1) f_2(x_2) \delta(x_1 x_2 - \tau) = \int_0^1 dx_1 \int_0^1 dx_2 f_1(x_1) f_2(x_2) \frac{1}{x_1} \delta\left(x_2 - \frac{\tau}{x_1}\right)$$

$$\rightarrow \frac{dL}{d\tau} = \int_{\tau}^1 dx_1 f_1(x_1) f_2\left(\frac{\tau}{x_1}\right) \frac{1}{x_1}$$

$$\rightarrow d\sigma_{A+B \rightarrow f+X} = d\sigma_{ij \rightarrow f} \frac{dL}{d\tau}$$

Define parton luminosity:

$$\frac{\tau}{\hat{s}} \frac{dL}{d\tau} = \frac{1}{s} \frac{dL}{d\tau} = \frac{dL}{d\hat{s}}$$




PQCD: Hadron Collisions - XV

137

Example: Drell-Yan from pp collisions


$$\begin{aligned} M &= \sqrt{\tau s} & \frac{d\tau}{dM} &= 2 \frac{M}{s} = 2 \frac{\tau}{M} \\ \frac{d\sigma}{dM} &= \frac{2}{M} \frac{4\pi\alpha^2}{9M^2} e_q^2 \tau \int_{\frac{1}{2}\ln\tau}^{\frac{1}{2}\ln\tau} dy f_q(\sqrt{\tau}e^y) f_{\bar{q}}(\sqrt{\tau}e^{-y}) \\ &= \frac{2}{M} \frac{4\pi\alpha^2}{9M^2} e_q^2 \tau \int_{\tau}^1 \frac{dx}{x} f_q(x) f_{\bar{q}}(\tau/x) \\ &= \frac{2}{M} \frac{4\pi\alpha^2}{9} e_q^2 \left[\frac{\tau}{M^2} \frac{d\mathcal{L}_{q\bar{q}}}{d\tau} \right] \end{aligned}$$



PQCD: Hadron Collisions - XVI

138

Effect of scaling violations:



PQCD: Hadron Collisions - XVII

139

Consider all the 2-body processes in QCD:

$$q q \rightarrow q q, q \bar{q} \rightarrow q \bar{q}$$

$$q g \rightarrow q g, \bar{q} g \rightarrow \bar{q} g, g g \rightarrow g g, q \bar{q} \rightarrow g g, g g \rightarrow q \bar{q}$$

Quarks only

Quarks and/or Gluons

All yielding 2 jets to LO



Figure 1: Feynman diagram for $q_i q_j \rightarrow q_i q_j$, $i \neq j$




Figure 2: Feynman diagrams for $q_i q_i \rightarrow q_i q_i$




Figure 5: Feynman diagrams for $q_i \bar{q}_i \rightarrow g g$

PQCD: Hadron Collisions - XVIII


140

When quark only processes can be identified, expect

$$\frac{d\sigma}{d(\cos \theta^*)} = \frac{\pi \alpha_s^2}{2 s_{ij}} |M|^2$$

$$s_{ij} = (x_i p_i + x_j p_j)^2 \approx x_i x_j s$$


$$\rightarrow |M|^2 \propto 1 + \cos^2 \theta^* \text{ as usual}$$



@TBA

PQCD: Hadron Collisions - XIX

141



@TBA

PQCD: Hadron Collisions - XX


142

Complete jet evolution cannot be computed in a full QCD framework:

As shown in the picture, *increasing* time scales correspond to *decreasing* Q^2 scale, down to a region where perturbative expansion cannot be granted

Conversion of quarks and gluons into hadrons (*Fragmentation* or *Hadronization*) must be evaluated by ad-hoc prescriptions (models).

QCD-inspired, successful models often based on string-like behavior of $q\bar{q}$ pairs



@TBA


PQCD: Hadron Collisions - XXI

143

Typical model implemented in fragmentation Montecarlo programs

$q\bar{q}$ Interaction

QED-like at small distance



Gluon self-interaction yields *string* (flux tube) pattern at large distance: $F = \text{const}$



Picture baryons as ‘mesons’:

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{3}^* \oplus \mathbf{6}$$

$$qqq = \underbrace{qq}_{\sim \bar{q}} + q$$

Confinement - I

144

$$\alpha_s(|q^2|) \simeq \frac{12\pi}{21 \ln(|q^2|/\Lambda^2)}, \quad |q^2| \gg \Lambda^2$$

When $|q^2| \sim \Lambda^2$, the previous expression does not apply

$\alpha_s(\Lambda^2)$ is large

Strong interaction is strong

Cannot rely on perturbative expansion

In a general sense, expect Λ to mark the low energy range, corresponding to soft (low q^2) processes

Bound states: Non-perturbative, ‘white’, energy scale $\approx \Lambda$

Does $a_s(\Lambda^2)$ correspond to the color confinement range?

Very likely. But remember:

It is not yet convincingly shown that QCD is a confining theory

Confinement - II

145

QCD: At large color charges separation, field lines compressed to tube-like regions

Reason: Gluon-gluon interaction

→ ~ String



→ $F(r) \approx \text{const} \rightarrow V(r) = kr$ Linearly confining potential

$k \sim 1 \text{ GeV / fm}$


Confinement - III

146

Regge trajectories: Old concept, adapted by potential scattering theory

Very general property, not related to any constituent model:


Linear relationship between angular momentum and $(\text{mass})^2$ of resonances



Confinement - IV

147

String model of mesons: Simple ‘explaination’ of Regge trajectories



$$F^\mu = \frac{dP^\mu}{d\tau}, F^\mu = (\gamma \mathbf{F} \cdot \mathbf{u}, \gamma \mathbf{F}) \text{ relativistic 2nd law}$$

$$\rightarrow m = E = W = 2 \int_0^R \gamma \mathbf{F} \cdot d\mathbf{r}$$

$$\mathbf{F} = k \hat{\mathbf{r}} \leftrightarrow \text{linear potential}$$

$$\rightarrow m = E = 2 \int_0^R \gamma k \hat{\mathbf{r}} \cdot d\mathbf{r} = 2 \int_0^R \frac{k}{\sqrt{1-\beta^2}} dr$$

$$\beta = \frac{r}{R} \rightarrow m = E = 2k \int_0^R \frac{dr}{\sqrt{1-\left(\frac{r}{R}\right)^2}} = \pi k R$$

$$J = 2k \int_0^R \frac{\frac{r}{R}}{\sqrt{1-\left(\frac{r}{R}\right)^2}} dr = \frac{1}{2} \pi k R^2 = \frac{m^2}{2\pi k}$$

Gluonia - I

148

QCD : Leading to predict new, ‘exotic’ ($=$ non $q\bar{q}$) mesonic states

Quarkless mesons: no valence quarks

\rightarrow *Gluonium*, aka *Glueball*

Expect, among others, exotic quantum numbers

Flavor: **1** Singlet (\leftarrow no quark)

Color: Bound state \rightarrow Must be **1** Singlet (\leftarrow ‘white’)

\rightarrow 2 g at least

$$\mathbf{8} \otimes \mathbf{8} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10} \oplus \overline{\mathbf{10}} \oplus \mathbf{27}$$

Pick singlet:

1 \leftrightarrow Symmetric

Bose statistics \rightarrow Spin \times Orbital: Symmetric

Observe:

1 of $SU(3)_C$ exchange-symmetric
when originated by

$$\mathbf{8} \otimes \mathbf{8} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10} \oplus \mathbf{10}^* \oplus \mathbf{27}$$

Gluonia - II

149

$$\mathbf{J} = \mathbf{L} + \mathbf{S}$$

By taking S – wave (space symmetric):

$$L = 0 \rightarrow J = S = 1 \oplus 1 = 0, 1, 2$$

$S = 0, 2$ Symmetric → OK

$$\left. \begin{aligned} P &= (-1)^L = +1 \\ C &= (-1)^2 = +1 \end{aligned} \right\} \rightarrow 0^{++}, 2^{++}$$


By taking P – wave (space antisymmetric):

$$L = 1 \rightarrow J = 1 \oplus 1 \oplus 1 = 0, 1, 2, 3$$

$S = 1$ Antisymmetric → OK

$$\left. \begin{aligned} P &= (-1)^L = -1 \\ C &= (-1)^2 = +1 \end{aligned} \right\} \rightarrow 1^{-+}, \text{ Exotic!}$$

Compare to $q\bar{q}$, standard mesons:



1×1	$\begin{matrix} 2 \\ +2 \end{matrix}$	$\begin{matrix} 2 & 1 \\ +1 & +1 \end{matrix}$	$\begin{matrix} 2 \\ 0 \\ 0 \end{matrix}$	$\begin{matrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{matrix}$	$\begin{matrix} 2 \\ 0 \\ 0 \end{matrix}$	$\begin{matrix} 1 & 0 \\ 0 & 0 \end{matrix}$	$\begin{matrix} 1/6 & 1/2 & 1/3 \\ 2/3 & 0 & 1/3 \end{matrix}$	$\begin{matrix} 2 & 1 \\ -1 & -1 \end{matrix}$	$\begin{matrix} 2 \\ -2 \end{matrix}$	$\begin{matrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{matrix}$	$\begin{matrix} 2 \\ -2 \end{matrix}$

Gluonia - III

150

Indeed, build $2g$ state out of single gluon states with defined helicity:

$$U_P |\mathbf{k}, R; -\mathbf{k}, R\rangle = |-\mathbf{k}, L; \mathbf{k}, L\rangle \quad \text{not a } U_P \text{ eigenstate}$$

$$U_P |\mathbf{k}, L; -\mathbf{k}, L\rangle = |-\mathbf{k}, R; \mathbf{k}, R\rangle \quad \text{not a } U_P \text{ eigenstate}$$

$$U_P |\mathbf{k}, R; -\mathbf{k}, L\rangle = |-\mathbf{k}, L; \mathbf{k}, R\rangle \equiv |\mathbf{k}, R; -\mathbf{k}, L\rangle$$

$\rightarrow U_P$ eigenstate, $\eta_P = +1$, $J_3 = +2 \rightarrow J = 2$

$$U_P |\mathbf{k}, L; -\mathbf{k}, R\rangle = |-\mathbf{k}, R; \mathbf{k}, L\rangle \equiv |\mathbf{k}, L; -\mathbf{k}, R\rangle$$

$\rightarrow U_P$ eigenstate, $\eta_P = +1$ $J_3 = -2 \rightarrow J = 2$

$$|\mathbf{k}, R; -\mathbf{k}, R\rangle \pm |\mathbf{k}, L; -\mathbf{k}, L\rangle$$

$\rightarrow U_P$ eigenstates, $\eta_P = \pm 1$, $J_3 = 0 \rightarrow J = 0, 2$

\rightarrow Pick $|\mathbf{k}, R; -\mathbf{k}, R\rangle + |\mathbf{k}, L; -\mathbf{k}, L\rangle$ (symmetric) $\rightarrow \eta_P = +1$

Quarkonium - I

151

Small distance: Perturbative!

Indeed: Large quark mass \rightarrow Large Q^2

\rightarrow One gluon exchange OK

Non relativistic effective potential \sim Coulomb-like

$$\rightarrow V \left(r \ll \frac{1}{m_q} \right) \approx -\frac{4}{3} \frac{\alpha_s}{r}$$

Add phenomenological confining term: String inspired

\rightarrow Full potential:

$$V(r) \approx -\frac{4}{3} \frac{\alpha_s}{r} + kr$$


Most interesting $q\bar{q}$ physics case: Heavy, neutral, flavorless mesons

In order to better understand it, revert for a while to simple QED bound state: *Positronium*

Quarkonium - II

152

Bound state of electron - positron: Similar to Hydrogen atom




Quarkonium - III

153


Family portrait of *-onia*:

@TBA




Quarkonium - IV

154



@TBA



Striking similarity, \approx same energy scale *above ground state*

Quarkonium - V

155


Original interest for non-relativistic, Schrodinger equation approach:

$$\Delta E(\text{charm}) \approx \Delta E(\text{bottom})$$


Amazingly close

E288	$M(\Upsilon') - M(\Upsilon)$	$M(\Upsilon'') - M(\Upsilon)$
Two-level fit	$650 \pm 30 \text{ MeV}$	
Three-level fit	$610 \pm 40 \text{ MeV}$	$1000 \pm 120 \text{ MeV}$
$M(\psi') - M(\psi)$	$\approx 590 \text{ MeV}$	

Yesterday 1977



Today 2012



Quarkonium - VI

156

Reminder: Bohr radius

$$R = \frac{n^2}{\alpha m} \rightarrow R_0 = \frac{1}{\alpha m}$$

Consequence of Coulomb potential, static limit of 1 photon diagram

If 1 gluon exchange approximation can be granted, expect for quarkonium

$$R_{q\bar{q}} \sim \frac{1}{\alpha_s m}$$

Observe: m large $\rightarrow R$ small $\rightarrow \alpha_s$ small \rightarrow 1 gluon appr. OK: Self-consistent

Use phenomenological, $q\bar{q}$ confining term like this:

$$V(r) = -\frac{4}{3} \frac{\alpha_s}{r} + kr$$

Solve Schrodinger equation with these terms

(Add more terms to take into account relativistic & color-hyperfine effects)

Question: Which form of effective potential would yield m -independent ΔE ??

Quarkonium - VII

157

Scaling Schroedinger:

$$\psi(r) = R(r)Y_{lm}(\theta, \varphi), \quad u(r) = rR(r)$$

$$\mu = \frac{m}{2} \quad \text{Reduced mass}$$

$$V(r) = \lambda r^\nu \quad \text{Power law potential}$$

$$\frac{\hbar^2}{2\mu} \frac{d^2u}{dr^2} + \left[E - \lambda r^\nu - \frac{l(l+1)}{2\mu r^2} \right] u(r) = 0 \quad \text{Radial Schrodinger equation}$$

$$r = \rho \left(\frac{\hbar}{2\mu|a|} \right)^{\frac{1}{2+\nu}} \quad \text{Scale radial distance}$$

$$E = \varepsilon \left(\frac{\hbar}{2\mu|\lambda|} \right)^{-\frac{2}{2+\nu}} \frac{\hbar^2}{2\mu} \quad \text{Scale energy}$$

$$w(\rho) = u(r)$$

$$\rightarrow \frac{d^2w}{d\rho^2} + \left[\varepsilon - \rho^\nu - \frac{l(l+1)}{\rho^2} \right] w(\rho) = 0 \quad \text{Adimensional radial equation}$$

3 parameters:

$$\mu \text{ reduced mass } \equiv \frac{m_q}{2} \text{ for } q\bar{q}$$

λ strength

ν exponent

Quarkonium - VIII

158

Scaling laws

Length: $L \propto (\mu |\lambda|)^{-\frac{1}{\nu+2}}$

Energy: $E \propto (\mu)^{-\frac{\nu}{\nu+2}} |\lambda|^{\frac{2}{\nu+2}}$

Coulomb $(\mu |\lambda|)^{-1}$ $\mu |\lambda|^2$

Logarithmic $(\mu |\lambda|)^{-\frac{1}{2}}$ $\mu^0 |\lambda|$

Linear $(\mu |\lambda|)^{-\frac{1}{3}}$ $\mu^{-\frac{1}{3}} |\lambda|^{\frac{2}{3}}$


Harmonic $(\mu |\lambda|)^{-\frac{1}{4}}$ $\mu^{-\frac{1}{2}} |\lambda|^{\frac{1}{2}}$

Well $(\mu |\lambda|)^0$ μ^{-1}

Quarkonium - IX

159

Cornell potential



Quarkonium - X

160

Several interesting applications:

1) Logarithmic potential yielding ΔE mass independent

Also obtained by properly fitted Cornell potential

$$\rightarrow \text{Fit} \quad \left\{ \begin{array}{l} \alpha_s \left(q^2 = (2m_q)^2 \right) \sim 0.25 - 0.35 \\ k \sim 1 \text{ GeV / fm} \end{array} \right.$$


2) Extra bonus:

$$\text{Probability density} \propto L^{-3} \rightarrow |\psi(0)|^2 \sim (\mu|\lambda|)^{\frac{3}{\nu+2}}$$

→ Fix partial width to e^+e^- of vector mesons


Crystal Ball - I

161



@TBA

94% of solid angle covered



Sodium Iodide

$NaI(Tl)$: Inorganic scintillating crystal; Tl is an activator

Merits:

Can grow large crystals

Lots of light

Crystal Ball - II

162

672 optically isolated NaI(Tl) crystals, 15.7 radiation lengths thick


Inner radius 25.3 cm; Outer radius 66.0 cm

CB geometry: Based on *icosahedron*.

Each of the 20 triangular faces (major triangles) is divided into four minor triangles, each consisting of nine separated crystals.

Each crystal: Truncated triangular pyramid, 40.6 cm high, pointing towards the center of the Ball.

Side on the inner end: 5.1 cm ; Side on the outer end: 12.7 cm




Crystal & Photomultiplier

Crystal Ball - III

163

Icosahedron magic: Platonic solid (!) , 20 equilateral triangle faces




Triangle count:

Large triangle 20

Small triangle 80


Crystal < 720 (edges)



@TBA


Crystal Ball - IV

164




Inclusive photon spectrum

@TBA




Most important results, among many:
Tune beam energy as to form $\psi'(3686)$
Observe decays into photon + X



Crystal Ball - V

165

After moving the CB detector to DORIS (DESY, Hamburg): Bottomonium!
Observation of the P-wave triplets



@TBA

Figure 11.2: The photon spectrum from Υ' decays obtained by the Crystal Ball Collaboration at DORIS II. A triplet of lines corresponding to $\Upsilon' \rightarrow \gamma \chi_b({}^3P_{2,1,0})$ is seen between 100 and 200 MeV. The decays $\chi_b \rightarrow \gamma \Upsilon$ produce the unresolved signal between 400 and 500 MeV [R. Nernst *et al.*, *Phys. Rev. Lett.* **54**, 2195 (1985)].

Non-Perturbative QCD - I

166

Needed to deal with *bound states* and *soft interaction* regime

Very difficult problem

Different approaches available:

Lattice QCD

Chiral Perturbation Theory

Non-Relativistic QCD

Heavy Quark Effective Theory


...

Deep waters, not even surfed in this course

Non Perturbative QCD - II

167

Perform QCD calculations over a discretized space-time (lattice)



$q\bar{q}$ potential from lattice


$$V(r) = -\frac{4}{3} \frac{\alpha_s}{r} + ar : \text{Not a bad idea after all...}$$

Non Perturbative QCD - III

168

Examples:

Charmonium levels from lattice



Predicted glueball spectrum from lattice

