

Elementary Particles II

3 – The Higgs Particle at the LHC

Predictions, Strategies, Machines, Detectors, Data,
Measurements, Tests of SM

SSB - I

Field theory: Real scalar field

$$L = \frac{1}{2}(\partial_\mu \phi)^2 - V(\phi)$$

$$V(\phi) = \frac{1}{2}\mu^2\phi^2 + \frac{1}{4}\lambda^2\phi^4, \lambda > 0$$

Reflection symmetric: $V(\phi) = V(-\phi)$

V Minima:

$$\mu^2 > 0 : \phi = 0$$


$$\mu^2 < 0 : \phi = \nu = \pm \sqrt{-\frac{\mu^2}{\lambda}}$$

V Minima: Defining *vacuum* state (\leftarrow Cannot have less energy)

$\mu^2 > 0$: Vacuum (non degenerate) \equiv Zero field

$\mu^2 < 0$: Vacuum (degenerate!) $= \nu \neq$ Zero field !!

ν = Vacuum Expectation Value (VEV) of ϕ



SSB - II

Fall 2016

Choose vacuum state:

$$\langle \phi(x) \rangle_0 = v \quad \text{Spontaneous Symmetry Breaking}$$

Define: $\phi(x) = v + \eta(x)$

$$\rightarrow L = \frac{1}{2}(\partial_\mu \eta)^2 - \lambda \left(v^2 \eta^2 - v \eta^3 - \frac{1}{4} \eta^4 \right) = \left[\frac{1}{2}(\partial_\mu \eta)^2 - \lambda v^2 \eta^2 \right] - \lambda v \eta^3 - \frac{1}{4} \lambda \eta^4$$


$$L = \frac{1}{2}(\partial_\mu \eta)^2 - \lambda v^2 \eta^2 \cancel{+ \text{higher powers of } \eta}$$

→ Free Klein-Gordon equation

→ Scalar quantum field

$$m^2 = 2\lambda v^2 \rightarrow m = \sqrt{-2\mu^2} > 0$$

[Observe: $\mu^2 < 0 \rightarrow$ Imaginary mass in original $L!$]



KO: $L(\eta) \neq L(-\eta)$

Reflection symmetry *spontaneously broken*

SSB - III

Fall 2016

Field theory: Complex scalar field

$$\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2), \phi^* = \frac{1}{\sqrt{2}}(\phi_1 - i\phi_2)$$

$$L = (\partial_\mu \phi)(\partial^\mu \phi)^* - V(\phi)$$

$$V(\phi) = \mu^2 \phi^* \phi + \lambda (\phi^* \phi)^2, \lambda > 0, \mu^2 < 0$$


$U(1)$ symmetric: $\phi \rightarrow \phi' = e^{i\alpha} \phi$

Observe: $U(1)$ continuous symmetry

V Minima:

$$\mu^2 < 0: \phi_1^2 + \phi_2^2 = -\frac{\mu^2}{\lambda}$$

→ Vacuum *infinitely* degenerate



Choose vacuum = $(v, 0)$

→ $U(1)$ symmetry spontaneously broken

SSB - IV

Fall 2016

Define:


$$\phi(x) = \frac{1}{\sqrt{2}} [v + \xi(x) + i\eta(x)]$$

$$L = \frac{1}{2} \left[(\partial_\mu \xi)^2 + (\partial_\mu \eta)^2 \right] + \mu^2 \eta^2 \cancel{+ \text{higher powers of } \eta}$$

Free Klein-Gordon equations for (ξ, η)

But: "Kinetic energy" terms for both ξ, η ; Mass term only for η

- η field excitations: *massive* scalar particles
- ξ field excitations: *massless* scalar particles, aka *Goldstone Bosons*



SSB - V

Fall 2016

SSB of a continuous, global (\leftarrow non local) symmetry

$\rightarrow \infty$ degenerate vacuum states

Symmetry generators transform any vacuum state into another one

Indeed, for a non-degenerate vacuum:

$\langle \phi \rangle_0$ Invariant under G :

$$\rightarrow e^{i\alpha G} \langle \phi \rangle_0 \simeq (1 + i\alpha G) \langle \phi \rangle_0 = \langle \phi \rangle_0 \leftrightarrow G \langle \phi \rangle_0 = 0$$

$\rightarrow G$ does annihilate a non-degenerate vacuum

For a degenerate vacuum:

$\langle \phi \rangle_0$ Non-invariant under G :

\rightarrow Goldstone Theorem:

n generators not annihilating the vacuum \rightarrow Appearance of n massless scalars

Also called *Goldstone bosons*

SSB - VI

Fall 2014

Example from condensed matter physics: Ferromagnet

Interaction rotationally invariant

But, below Curie temperature

→ Spontaneous magnetization in a random orientation


→ Rotational symmetry lost

→ Degenerate vacuum

→ Goldstones

Non-relativistic version of *massless*

Spin waves \equiv Zero energy-gap *quasi-particles* (i.e. lattice excitation)



Higgs Mechanism - I

Goldstone theorem would seem to definitely destroy our hint of a Standard Model:

- 3 + 1 massless gauge bosons, only one observed
- 4 massless scalar bosons, none observed

Local gauge invariance + SSB:
Higgs mechanism
evading Goldstone's theorem

Simple, yet subtle way of giving mass to gauge bosons without spoiling gauge invariance (and renormalizability)

Higgs Mechanism - II

Fall 2014

Example by Higgs :

$U(1)$ gauge group, require *local* symmetry:

Gauge vector boson A_μ to be introduced, coupling to some current as usual

Now: Add "sombrero" potential for a complex, scalar field $\phi = \phi_1 + i\phi_2$

$$L = \underbrace{[\partial_\mu - ieA_\mu]\phi^*}_{EM \text{ interaction of (charged) } \phi} [\partial^\mu + ieA^\mu]\phi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \mu^2\phi^*\phi - \lambda(\phi^*\phi)^2 + \underbrace{\dots}_{\text{Current-field interaction etc}}$$

As found before:

Degenerate vacuum state \rightarrow SSB picks as vacuum state $(v, 0)$

Higgs Mechanism - III

$$\rightarrow \phi = v + \eta_1 + i\eta_2$$

L written in terms of η_1, η_2 :

Upon quantization, 2 scalar particles $\rightarrow m_1 = \sqrt{2\lambda v^2}, m_2 = 0$

Plugging $\phi = v + \eta_1 + i\eta_2$ into L :

$$L = \frac{1}{2}(\partial_\mu \eta_1)(\partial^\mu \eta_1) - \frac{1}{2} \overbrace{2\lambda v^2}^{m_1^2} \eta_1^2 + \frac{1}{2}(\partial_\mu \eta_2)(\partial^\mu \eta_2) \\ + \underbrace{\frac{1}{2} \overbrace{(ev)^2}^{m_V^2} A^\mu A_\mu}_{\text{Massive vector!}} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \underbrace{eva^\mu \partial_\mu \eta_2}_{??} + \dots$$

Attempting to understand L :

Massive vector field A^μ + Massive scalar field η_1 + Massless scalar field η_2

Troubling term coupling A^μ and η_2

Higgs Mechanism - IV

Use gauge invariance:

$$\begin{cases} \phi \rightarrow \phi' = e^{-ie\theta(x)}\phi \\ A^\mu \rightarrow A'^\mu = A^\mu + \partial_\mu \theta \end{cases}$$

Choose θ to make ϕ real: Then $\eta_2 \equiv 0$ (\leftarrow Unitary gauge)

$$\rightarrow L = \frac{1}{2}(\partial_\mu \eta_1)(\partial^\mu \eta_1) - \frac{1}{2}2\lambda v^2 \eta_1^2 + \underbrace{\frac{1}{2}(ev)^2 A^\mu A_\mu}_{\text{Massive vector!}} - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \dots$$

Massive vector field A^μ + Massive scalar field η_1

Counting degrees of freedom:

$$\underbrace{2}_{A_\mu} + \underbrace{1+1}_{\phi} = \underbrace{3}_{A_\mu} + \underbrace{1}_{\eta_1} \quad \text{OK}$$

Standard picture:

By effect of a smart gauge transformation,
 the massless vector field A_μ has eaten the Goldstone boson η_2
 to become massive

Higgs Mechanism - V

Fall 2016

Attempting to dissipate some misunderstandings likely to sneak in:
Mostly related to our naive perception of what is really a 'particle'

1) Where is the mass?

To identify mass terms in L : Not necessarily a trivial task

Key point: Particle content *only meaningful in perturbative expansion*

Example: $L = (\partial_\mu \phi)(\partial^\mu \phi)^* - (-\mu^2 \phi^* \phi) - \lambda (\phi^* \phi)^2, \lambda > 0, \mu^2 > 0$

$-\mu^2 \rightarrow$ Imaginary mass \rightarrow Nonsense $\rightarrow ???$

But: To use this form of L to extract Feynman rules, should expand around $|\phi| = 0$

Unstable extremum \rightarrow Can't make it

Rewrite by expanding around $\eta = 0$: $L = \frac{1}{2}(\partial_\mu \eta)^2 - \lambda \nu^2 \eta^2 + \dots$

Stable extremum \rightarrow OK

\rightarrow Particle content should be identified in this form

Higgs Mechanism - VI

Fall 2016

2) What's so special in unitary gauge?

Nothing: L invariant under local gauge transformations,
including the one to unitary gauge:

→ L describe the same physics before and after the gauge transformation

But: Particle content much easier to extract in the unitary gauge

3) Disappearing Goldstones !?

Indeed: And re-appearing as extra degrees of freedom for massive gauge bosons

See comment above on the tricky business of defining what is a particle...

4) What decides which vacuum is selected among the many?

Not really relevant: Any choice yields identical results

5) Could we make it with the SM without SSB and all that complicated swapping of degrees of freedom?

Actually no: SSB is an *intrinsic* feature of certain quantum systems
(with an infinite number of degrees of freedom)

SM Reminder - I

Fall 2014

Extend Abelian Higgs model to non-Abelian gauge symmetry:

$$\text{Gauge group} = SU(2)_L \otimes U(1)_Y$$

To add SSB to the Standard Model :

Add a doublet of complex, scalar fields:

$$\phi(x) = \begin{pmatrix} \phi^+(x) \\ \phi^0(x) \end{pmatrix}$$

$$\text{Assuming } y=1 \rightarrow \begin{cases} Q[\phi^+(x)] = +1 \\ Q[\phi^0(x)] = 0 \end{cases}$$

SM Reminder - II

Fall 2014

$SU(2)_L \otimes U(1)_Y$ Gauge transformation of doublet:

$$\phi \rightarrow \phi' = \exp \left\{ -i \left[\frac{g}{2} \mathbf{a}(x) \cdot \boldsymbol{\tau} + \frac{g'}{2} y \theta(x) I \right] \right\} \phi$$

$SU(2)_L \otimes U(1)_Y$ Covariant derivative:

$$D^\mu = \partial^\mu + i \left[\frac{g}{2} \boldsymbol{\tau} \cdot \mathbf{W}^\mu + \frac{g'}{2} y B^\mu \right]$$

→ Additional term to EW lagrangian:

$$L_H = D_\mu \phi^\dagger D^\mu \phi - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2$$

Take $\mu^2 < 0$, $\lambda > 0$:

$$|\phi|_{\min}^2 = -\frac{\mu^2}{2\lambda} = \frac{v^2}{2} \rightarrow v = \sqrt{-\frac{\mu^2}{\lambda}}$$

Pick ground state (= vacuum) as

$$\langle \phi \rangle_0 = \begin{pmatrix} 0 \\ v \\ \sqrt{2} \end{pmatrix} \rightarrow \text{SSB of Electroweak gauge symmetry}$$

$$v = 246 \text{ GeV}$$

SM Reminder - III

Introduce field deviation from vacuum:

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} \sigma_1 + i\sigma_2 \\ v + \eta_1 + i\eta_2 \end{pmatrix} \rightarrow V = -\frac{\mu^4}{4\lambda} + \lambda v^2 \eta_1^2 + \lambda v \eta_1 (\sigma_1^2 + \sigma_2^2 + \eta_1^2 + \eta_2^2) + \frac{\lambda}{4} (\sigma_1^2 + \sigma_2^2 + \eta_1^2 + \eta_2^2)^2$$

After properly 'rotating' to Unitary Gauge:

1 massive scalar: $\eta_1, m_{\eta_1} \equiv m_H = \sqrt{2\lambda v^2} \leftarrow \text{The Higgs}$

2 massive, charged vectors: $W^\pm, m_W = \frac{g}{2} \sqrt{-\frac{\mu^2}{\lambda}}$

1 massive, neutral vector: $Z^0, m_Z = \frac{\sqrt{(g^2 + g'^2)}}{2} \sqrt{-\frac{\mu^2}{\lambda}}$

→ Relating model parameters to independently measured constants $e, G_F, \sin \theta_W$:

$$M_W = \sqrt{\frac{\sqrt{2}g^2}{8G_F}} = \sqrt{\frac{\sqrt{2}e^2}{8G_F \sin^2 \theta_W}} \simeq 77.5 \text{ GeV}, \quad M_Z = \frac{M_W}{\cos \theta_W} \simeq 88.4 \text{ GeV}$$

$$M_H = \frac{\sqrt{2}\lambda}{G_F} = ???$$

SM Reminder - IV

Fall 2014

Gauge terms of L in the unitary gauge, in terms of the physical fields:

$$\begin{aligned} L_B + L_H &= -\frac{1}{4} F_{\mu\nu}(x) F^{\mu\nu}(x) && \text{Photon} \\ &- \frac{1}{2} F_W{}_{\mu\nu}(x) F^{W\dagger\mu\nu}(x) + \frac{1}{2} m_W^2 W_\mu^\dagger W^\mu && W^\pm \text{ boson} \\ &- \frac{1}{4} Z_{\mu\nu}(x) Z^{\mu\nu}(x) + \frac{1}{2} m_Z^2 Z_\mu Z^\mu && Z^0 \text{ boson} \\ &+ (\partial_\mu \sigma)(\partial^\mu \sigma) - \frac{1}{2} m_H^2 \sigma^2 && H \text{ Higgs boson} \\ &+ L_{BB}^I + L_{HH}^I + L_{HB}^I && \text{Gauge-Higgs, Higgs self-, Gauge self-interactions} \end{aligned}$$

SM Reminder - V

Fall 2014

Fermion masses: Yukawa (scalar) coupling

Describing interaction between Dirac and scalar fields:

Example: Single lepton family

$$L_{HL} = -g_l \left[\bar{\Psi}_l^L \Phi \psi_l^R + \bar{\psi}_l^R \Phi^\dagger \Psi_l^L \right] - \underbrace{g_{\nu_l}}_{=0 \text{ for massless neutrino}} \left[\bar{\Psi}_l^L \tilde{\Phi} \psi_{\nu_l}^R + \bar{\psi}_{\nu_l}^R \tilde{\Phi}^\dagger \Psi_l^L \right], \quad \tilde{\Phi} = \begin{pmatrix} \phi_b^* \\ -\phi_a^* \end{pmatrix}$$

In the unitary gauge:

$$L_{HL} = -\frac{1}{v} m_l \bar{\psi}_l \psi_l \sigma - \frac{1}{v} m_{\nu_l} \bar{\psi}_{\nu_l} \psi_{\nu_l} \sigma$$

Lepton masses in terms of model parameters:


$$m_l = \frac{v g_l}{\sqrt{2}}, \quad m_{\nu_l} = \frac{v g_{\nu_l}}{\sqrt{2}} (= 0 \text{ in the Minimal Standard Model})$$

g_l individual constant

SM Reminder - VI

Fall 2014

Higgs vertexes:



Observe: Amplitude \sim mass(fermions), mass²(bosons)

About the Higgs Field - I

Fall 2014

Universal, constant field

Lorentz scalar → Same value in any frame, rotation invariant

Non-standard feature:

Vacuum expectation value $v \neq 0$

Usual analogy: Spontaneously magnetized ferromagnet:

$\mathbf{M} \neq 0$ below Curie temperature

→ Pick up a direction

Ground state rotationally not symmetric, in spite of H being symmetric

About the Higgs Field - II


Fall 2014

Better analogy: Superconductor

Energy difference between normal and s.c. state at two different temperatures

$$\Delta E = a(T)|\psi|^2 + \frac{1}{2}b(T)|\psi|^4 + \dots \quad \text{Landau theory of phase transitions}$$

ψ is the Cooper pair ‘wave function’ $\rightarrow |\psi|^2 \sim \text{density of Cooper pairs}$



Below T_c , the minimum energy state (‘vacuum’) occurs for $\psi = \psi_0 \neq 0$, phase undefined
 $\rightarrow U(1) QED$ gauge invariance spontaneously broken

\rightarrow Photon becomes ‘massive’ $\rightarrow \mathbf{B} = 0$ inside

About the Higgs Field - III

Fall 2014

ψ ‘Higgs field’ of superconductivity: $\langle \psi \rangle \neq 0 \leftrightarrow$ Permanent supercurrents

Superconductive state:

‘Higgs field’ = ‘Wave function’ of Cooper pairs

→ Not a fundamental field

→ ‘Composite’ field of fundamentals fermions (electrons)

Why there is the composite?

e - e effective interaction: Attractive (!) due to e – lattice interaction

Is the ‘real’ Higgs field a genuine, fundamental field or a composite?

Good question..No answer (yet):

Take it as a fundamental field

About the Higgs Field - IV

Fall 2014

A couple of questions:

1) What about the nonzero VEV of the Higgs field?

Higgs: Unique field whose VEV $\neq 0$

Similar to magnetization $\mathbf{M} \neq 0$ in a ferromagnet

But:

In a vacuum \rightarrow Not related to many body effects

Lorentz scalar \rightarrow No preferred direction, reference frame

2) Does it involve a new force? 'Giving mass to \approx all the fundamental constituents' ??

- Part of the standard EW interaction, often as a negligible contribution:

Higgs *particle* exchange diagrams between Fermion lines normally strongly suppressed by m_f/m_w factors as compared to γ, Z^0, W^\pm exchange (Not true for t quark!)

3- & 4-boson diagrams with and without H similar

- Crucial role as 'Background' interaction:

For most particles Higgs *field* coupling translates into *inertial mass* !

About the Higgs Field - V

Fall 2014

Apparently contributing to vacuum energy density:

Beware: Take *potential energy* $V(\phi)$

$$\begin{aligned}V_{\min} = V(v) &= \frac{1}{2}\mu^2 v^2 + \frac{1}{4}\lambda v^4 \quad \text{use } \mu^2 = -\lambda v^2 \\&= -\frac{1}{4}\lambda v^4 \quad \text{use } m_h^2 = 2\lambda v^2 \\&= -\frac{1}{8}m_h^2 v^2\end{aligned}$$

Constant term: Usually not considered

Does not enter field equations, where only energy *differences* count

But: Taken into account by gravity

→ Cosmological term ?

Cosmological constant : Possibly additional term in Einstein's field equations

May yield long range attraction/repulsion, according to sign

Invented by Einstein in order to guarantee static universes

Rejected by Einstein at the time of discovery of expansion of the Universe

Recently resurrected following the discovery of accelerated expansion

About the Higgs Field - VI

Fall 2014

$$\text{Zero point energy} = -\frac{1}{8}m_h^2v^2 \sim \rho_{Higgs}$$

$$\text{Indeed: } \left[-\frac{1}{8}m_h^2v^2 \right] = \underbrace{E^4}_{GeV^4} = E(E^{-1})^{-3} \rightarrow \frac{E}{L^3} \quad \text{Energy density}$$

$$\rho_{Higgs} \sim 1.210^8 \text{ GeV}^4$$

By assuming ρ to be a cosmological term, compare:

$$\rho_{observed} \sim 10^{-47} \text{ GeV}^4 !$$

$\rightarrow \rho_{Higgs}$ 55 orders of magnitude too big

(and with the wrong sign...)

Quick fix: $V(v)$ can be set = 0 by adding a constant to $V(\phi)$

Constant apparently unrelated to $m_h, v\dots$

...to be chosen to an accuracy of 1 part out of 10^{55} !

Fine tuning problem, still essentially unsolved

Something missing?

About the Higgs Field - VII

Fall 2014

Higgs boson: Quantum excitation of the field, mass m_H *not* given by the field

Further issue:

$V(\phi)$ appearing in L : *Classical* potential

→ Must be quantized

→ Will be used perturbatively

→ Radiative corrections will modify the classical $V(\phi)$

Similar to vacuum polarization corrections to Coulomb potential in QED

(Uheling potential & Lamb shift)

Standard effect:

Running constants, including λ

$$L_H = D_\mu \phi^\dagger D^\mu \phi - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2$$

$$\lambda = \lambda(q^2)$$

→ m_H modified by radiative corrections

Upon taking $\mu^2 < 0$, $\lambda(0) > 0$

→ λ evolution depending on β -functions

About the Higgs Field - VIII

Fall 2014

Running couplings and β -functions:

$$\frac{dg^2}{d \ln Q^2} \equiv 4\pi\beta(g^2) = \underbrace{bg^4}_{1 \text{ loop}} + \underbrace{O(g^6)}_{2 \text{ loop}} + \dots$$

$$\rightarrow \frac{dg_i^2}{d \ln Q^2} = 4\pi\beta_i(g_i^2) \simeq b_i g_i^4$$

For the EW interaction:


$$b_g = -\frac{19}{6 \cdot 16\pi^2}, \quad b_{g'} = +\frac{41}{6 \cdot 16\pi^2}$$

Higgs couplings:

$$\frac{d\lambda}{d \ln Q^2} = \frac{1}{32\pi^2} \left[24(\lambda^2 + h_t^2 - h_t^4) - 3\lambda(3g^2 + g'^2) + \frac{3}{8}(2g^4 + (g^2 + g'^2)^2) \right] \text{ Self}$$

$$\frac{dh_t}{d \ln Q^2} = \frac{1}{32\pi^2} \left[9h_t^3 - h_t \left(8g_s^2 + \frac{9}{4}g^2 + \frac{17}{12}g'^2 \right) \right] \text{ Top (Yukawa)}$$

$$\rightarrow m_H < \left(\frac{2\sqrt{2}\pi^2}{3G_F \ln \frac{\Lambda}{\nu}} \right)^{1/2} \sim \begin{cases} O(140 \text{ GeV}), \Lambda \sim m_{Planck} \approx 1.2 \cdot 10^{19} \text{ GeV} \\ O(650 \text{ GeV}), \Lambda \sim 1 \text{ TeV} \end{cases}$$



About the Higgs Field - IX

Fall 2014

$$\frac{d\lambda}{d \ln Q^2} \sim \frac{3\lambda^2}{4\pi^2} \quad \text{Neglect smaller contributions at large } \lambda$$

$$\frac{d\lambda}{\lambda^2} \sim \frac{3}{4\pi^2} d \ln Q^2 \rightarrow -\frac{1}{\lambda(Q^2)} + \frac{1}{\lambda(\nu^2)} \sim \frac{3}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

$$\rightarrow \frac{1}{\lambda(Q^2)} \sim \frac{1}{\lambda(\nu^2)} - \frac{3}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

$$\lambda(\nu^2) = \frac{G_F m_H^2}{\sqrt{2}}$$

$$\rightarrow \lambda(Q^2) \sim \frac{\lambda(\nu^2)}{1 - \frac{3}{4\pi^2} \lambda(\nu^2) \ln \frac{Q^2}{\nu^2}}$$

$$\lambda \rightarrow \infty \text{ as } \frac{3}{4\pi^2} \lambda(\nu^2) \ln \frac{Q^2}{\nu^2} \rightarrow 1 \quad \text{Diverging at 'Landau pole'} \quad Q_{LP} = v \exp \left(\frac{2\pi^2}{3\lambda(\nu^2)} \right) = v \exp \left(\frac{2\sqrt{2}\pi^2}{3G_F m_H^2} \right)$$

$$\rightarrow \text{New physics required at scale } \Lambda < Q_{LP} \rightarrow \ln \frac{\Lambda}{v} < \left(\frac{2\sqrt{2}\pi^2}{3G_F m_H^2} \right) \rightarrow m_H < \left(\frac{2\sqrt{2}\pi^2}{3G_F \ln \frac{\Lambda}{v}} \right)^{1/2}$$

E.Menichetti - Universita' di Torino

About the Higgs Field - X

Fall 2014

$$\frac{d\lambda}{d \ln Q^2} \sim -\frac{3h_t^4}{4\pi^2} \text{ Neglect smaller contributions at small } \lambda$$

$$\rightarrow d\lambda \sim -\frac{3h_t^4}{4\pi^2} d \ln Q^2$$

$$\rightarrow \lambda(Q^2) \sim \lambda(\nu^2) - \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

λ must stay +ve in order to keep vacuum stable (!):

Don't like a too quick End of the World

$$\rightarrow \lambda(\nu^2) > \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2} \rightarrow \frac{G_F m_H^2}{\sqrt{2}} > \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2} \text{ for some } Q \sim \Lambda$$



$$\rightarrow m_H > \left(\frac{3h_t^4}{\sqrt{2}\pi^2 G_F} \ln \frac{\Lambda}{\nu} \right)^{1/2}$$

About the Higgs Field - XI


Fall 2014

Running couplings:

Sombrero: $\lambda > 0$
Relax



Dog Bowl: $\lambda < 0$
End of the World
(sometime)



E.Menichetti - Universita' di Torino


About the Higgs Field - XI

Fall 2014

Radiative corrections leading to major changes in the effective Higgs potential at large ϕ values:

Details tied to m_H, m_t


Might induce vacuum instability/metastability through fast/slow tunneling



About the Higgs Field - XII

Fall 2014

Theoretical upper & lower bounds on m_H :



E.Menichetti - Universita' di Torino

Hunting the Higgs

Fall 2014

Try to sketch some guidelines:

- (i) Production modes → Machines
- (ii) Decay modes → Detectors

Both related to couplings


Compare rates, backgrounds

- Sensitivity
- Further observables (Spin/Parity, Width, Branching Ratios,..)


Higgs Production - I

Fall 2014

Start from H coupling to Fermions:



Compare to coupling to Z^0 :




→ H coupling down by a factor $\sim \frac{m_f}{m_W}$ as compared to Z^0

Higgs Production - II

Fall 2014

First mode:

s-channel formation:



Ideal for lineshape scan, provided cross-section is big enough

Lepton colliders:


Tough requirements on luminosity, energy resolution

Higgs Production - III

Fall 2014

Second mode:

H radiation from quarks, sizeable contribution from Top:



$t\bar{t}$ signature might be useful to tag

Higgs Production - IV

Fall 2014

Shift to gauge bosons:

Exclude massless photon, gluon *at tree level*

[Photon, gluon *loop* contribution to be taken into account: See later]

More promising: W, Z mass very large

$$V^\mu \text{---} H = 2i \frac{M_V^2}{v} g^{\mu\nu}$$




$$V^\mu \text{---} H = 2i \frac{M_V^2}{v^2} g^{\mu\nu}$$

Higgs Production - V

Fall 2014

Best modes:

'Higgsstrahlung', 'Gauge boson fusion'




Higgs Production - VI

Fall 2014

Beyond tree level: Very Important Loops


Lepton machines:

Interesting diagrams, also quite relevant to detection




Parton machines:

Dominant diagram at LHC




Higgs Decays - I

Fall 2014




$$\Gamma(H \rightarrow f\bar{f}) = N_C \frac{1}{8\pi} \frac{m_f^2}{v^2} M_H \left(1 - \frac{4m_f^2}{M_H^2}\right)^{3/2}$$




$$\begin{aligned} \Gamma(H \rightarrow WW) &= \frac{1}{4\pi} \frac{M_W^4}{M_H v^2} \left(1 - \frac{4M_W^2}{M_H^2}\right)^{1/2} \left(3 + \frac{1}{4} \frac{M_H^4}{M_W^4} - \frac{M_H^2}{M_W^2}\right) \\ \Gamma(H \rightarrow ZZ) &= \frac{1}{8\pi} \frac{M_Z^4}{M_H v^2} \left(1 - \frac{4M_Z^2}{M_H^2}\right)^{1/2} \left(3 + \frac{1}{4} \frac{M_H^4}{M_Z^4} - \frac{M_H^2}{M_Z^2}\right) \end{aligned}$$

Higgs Decays - II

Fall 2014




$$\Gamma(H \rightarrow gg) = \frac{M_H^3}{8\pi v^2} \left(\frac{\alpha_s}{\pi} \right)^2 n^2 |D(n)|^2$$




Higgs Decays - III

Fall 2014



$$\Gamma(H \rightarrow W tb) = N_C \frac{g^2}{2v^2} \frac{M_W^4}{M_H} |V_{tb}|^2 \int dQ_3 \frac{G^{\beta\nu} T_{\beta\nu}}{[s_{34} - M_W^2]^2}$$

3-body phase space factor




$$\Gamma(H \rightarrow W f_u f_d) = \frac{g^2}{v^2} \frac{3M_W^2}{256\pi^3} M_H S(x)$$

3-body phase space factor


Higgs Decays - IV

Fall 2014



$$\Gamma(H \rightarrow Z f\bar{f}) = \frac{g^2}{v^2} \frac{3M_Z^2}{256\pi^3} M_H S(x) \frac{R(\theta_w)}{\cos^2 \theta_w} \left(\frac{7}{12} - \frac{10}{9} \sin^2 \theta_w + \frac{40}{27} \sin^4 \theta_w \right)$$

3-body phase space factor




E.Menichetti - Universita' di Torino

Higgs Decays - V

Fall 2014


H decays entirely determined by Higgs mass:



Higgs Decays - VI

Fall 2014

H branching ratios:




Higgs Decays - VII

Fall 2014

E.Menichetti - Universita' di Torino

46



Lepton Collider - I

Fall 2014

Leptons: Easier to handle

Taking lifetime into account: Restrict to electron, muon (?)

Linear vs. Circular

s – channel formation:

Not feasible at e^-e^+ colliders:

Factor $\frac{m_e^2}{M_W^2} \sim 4 \cdot 10^{-11}$ → Tiny cross section

Better chance for a $\mu^-\mu^+$ collider


$\frac{m_\mu^2}{M_W^2} \sim 1.6 \cdot 10^{-6}$

Other channels more promising

Lepton Collider - II

Fall 2014


Electron collider: Tree level contributing diagrams



Lepton Collider - III

Fall 2014



Electron collider: 'Low energy' option



Lepton Collider - IV

Fall 2014



Electron collider: 'High energy' option



Lepton Collider - V

Fall 2014

m_H dependence:



Lepton Collider - VI

Fall 2014

\sqrt{s} (GeV)	$\langle L \rangle$ (ab $^{-1}$ /year)*	Rate (Hz) $e^+e^- \rightarrow$ hadrons	Years	Statistics
90	5.6	$2 \cdot 10^4$	1	$2 \cdot 10^{11}$ Z decays
160	1.6	25	1-2	$2 \cdot 10^7$ W pairs
240	0.5	3	5	$5 \cdot 10^5$ HZ events
350	0.13	1	5	$2 \cdot 10^5$ ttbar

* each interaction point

- ❑ Precise measurement (0.1% to 1%) of the Higgs Couplings
- ❑ Improve precision (statistics $\times 10^5$) on the measurements of the Z parameters [M_Z , Γ_Z , R_ℓ , R_b , R_c , Asymmetries & weak mixing angle]. Z rare decays.
- ❑ Scan W threshold (aiming at 0.5 MeV precision). W rare decays
- ❑ Scan ttbar threshold (aiming at 10 MeV)

Lepton Collider - VII

Fall 2014

Circular collider: Two main issues, among many

Bending field: Must keep the beam on orbit

$$\text{Orbit radius: } R = \frac{3.3p}{B} \quad m, GeV, T$$

First look:

Either low B , large R or high B , small R

RF power: Must provide energy to beam up to max. energy

(also compensating for synchrotron radiation loss)

Ex: LEP I

$$B = \frac{3.3 \cdot 45}{4300} \approx 0.034 \text{ T}$$

128 2 m cavities

Typical cavity max field: 1.5 MV / m

Typical beam current: 6+6 mA

→ Max. energy gain $\sim 128 \cdot 3 \sim 375 \text{ MeV / turn}$; RF max power $\sim 125 \text{ kW / cavity}$


Lepton Collider - VIII

Fall 2014

Another crucial point: Minimize *synchrotron radiation* loss

Process related to EM interaction of ultrarelativistic charged particles moving in a **B** field:

Similar to Bremsstrahlung



Critical energy:

$$\epsilon_c = 3hc\gamma^3 / (2R)$$

$$\epsilon_c(\text{keV}) = 2.218 E^3(\text{GeV}) / R(\text{m})$$

Ex: LEP II

$$E = 104 \text{ GeV} \rightarrow \epsilon_c \sim 580 \text{ keV}$$

Energy loss per particle, per turn:

$$\Delta E(\text{KeV}) = \frac{e^2 \gamma^4}{3\epsilon_0 R} = \begin{cases} 88.5 \frac{E(\text{GeV})^4}{R(\text{m})} & \text{Electrons} \\ 6.03 \frac{E(\text{TeV})^4}{R(\text{m})} & \text{Protons} \end{cases}$$

Power loss by a beam current I_b , to be restored by RF:

$$P(\text{kW}) = \frac{e\gamma^4}{3\epsilon_0 R} I_b = \begin{cases} 88.5 \frac{E(\text{GeV})^4}{R(\text{m})} I_b (\text{A}) & \text{Electrons} \\ 6.03 \frac{E(\text{TeV})^4}{R(\text{m})} I_b (\text{A}) & \text{Protons} \end{cases}$$


→ Pointing to large $R \rightarrow$ low B

Lepton Collider - IX

Fall 2014

Electron collider: Linear

International Linear Collider (Japan?)




- Superconducting RF cavities
- Gradient 32 MV/m
- $\sqrt{s} \leq 500$ GeV (1 TeV upgrade option)
- $L \sim 2 \cdot 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Focus on ≤ 500 GeV, physics studies also for 1 TeV
- Length ~ 31 km (500 GeV)

Lepton Collider - X

Fall 2014

Electron collider: Linear


Compact LInear Collider (CERN?)



- Copper RF cavities
- Gradient 150 MV/m (!)
- $\sqrt{s} = 3$ TeV
- $L \sim 6 \cdot 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Length ~ 41 km (500 GeV)

Lepton Collider - XI

Fall 2014




E.Menichetti - Universita' di Torino

Lepton Collider - XII

Fall 2014

Muon collider: Overall layout



Critical points:

- Proton LINAC+Booster (4 MW)

$$16 \text{ GeV} \times 1.5 \times 10^{15} \text{ pps} = 16 \text{ GeV} \times 250 \mu\text{A} = 4 \text{ MW} !$$

- Cooling (fast, large)

Ionization: dE / dx reducing both $p_{||}$ and p_{\perp}

RF restoring $p_{||} \rightarrow \frac{p_{\perp}}{p_{||}}$ reduced

Lepton Collider - XIII

Fall 2014

Muon collider

Pros:

Large H cross section in the s -channel $(\sigma_{\mu\mu} \approx 4 \cdot 10^4 \sigma_{ee})E$

Best energy resolution ('*Beamstrahlung*' strongly suppressed)

→ Unique feature: Can perform full scan of lineshape

Main requirements:

Energy resolution:
$$\begin{cases} E = \frac{m_H}{2} \simeq 63 \text{ GeV} \\ \sigma_E \leq \Gamma_H = 4.2 \text{ MeV} \end{cases} \rightarrow R = \frac{\sigma_E}{E} \leq 5 \cdot 10^{-5}$$

Feasible, but $L \sim 10^{31} \text{ cm}^{-2} \text{s}^{-1}$

→ 1 y data taking $\sim 10^{38} \text{ cm}^2 = 100 \text{ pb}^{-1}$

→ $N_H \sim 2000 \text{ y}^{-1}$

→ $\sigma_M \sim 100 \text{ keV}, \sigma_\Gamma \sim 200 \text{ keV}, \sigma_{\mu \text{ coupling}}$ to 3% in 1 year


Lepton Collider - XIV

Fall 2014

Breit-Wigner/Gaussian convolution:

$$\sigma(\mu^+ \mu^- \rightarrow h \rightarrow X) = \frac{4\pi \Gamma_h^2 \text{Br}(h \rightarrow \mu^+ \mu^-) \text{Br}(h \rightarrow X)}{(\hat{s} - m_h^2)^2 + \Gamma_h^2 m_h^2}.$$


$$\begin{aligned} \sigma_{\text{eff}}(s) &= \int d\sqrt{\hat{s}} \frac{dL(\sqrt{s})}{d\sqrt{\hat{s}}} \sigma(\mu^+ \mu^- \rightarrow h \rightarrow X) \\ &\propto \begin{cases} \Gamma_h^2 B/[(s - m_h^2)^2 + \Gamma_h^2 m_h^2] & (\Delta \ll \Gamma_h), \\ B \exp[-\frac{(m_h - \sqrt{s})^2}{2\Delta^2}] (\frac{\Gamma_h}{\Delta})/m_h^2 & (\Delta \gg \Gamma_h). \end{cases} \end{aligned}$$




Photon Collider - I

Fall 2014

General idea: Compton back-scattering of intense laser beam by the e beam





$$x = \left(\frac{4E_b\omega_0}{m_e^2} \right) \cos^2(\alpha/2) \approx 15.3 \left(\frac{E_b}{\text{TeV}} \right) \left(\frac{\omega_0}{\text{eV}} \right)$$




Photon Collider - II


Fall 2014



Most important loop diagrams for the 'blob': Dominated by largest mass fermions, gauge bosons



Making for a particularly clean study of H self-coupling



E.Menichetti - Universita' di Torino

Photon Collider - III

Fall 2014



Higgs self-couplings: Very important for $\begin{cases} \text{radiative corrections to } m_h \\ \text{signs of new physics} \end{cases}$

Origin of 3- and 4-linear self-couplings:

Re-writing L upon introducing shifted field, 1D example

$$\begin{aligned} \text{Kinetic term: } \mathcal{L}_{\text{kin}}(\eta) &= \frac{1}{2}(\partial_\mu(\eta + v)\partial^\mu(\eta + v)) \\ &= \frac{1}{2}(\partial_\mu\eta)(\partial^\mu\eta) \end{aligned}$$

$$\begin{aligned} \text{Potential term: } V(\eta) &= +\frac{1}{2}\mu^2(\eta + v)^2 + \frac{1}{4}\lambda(\eta + v)^4 \\ &= \lambda v^2\eta^2 + \lambda v\eta^3 + \frac{1}{4}\lambda\eta^4 + \frac{1}{4}\lambda v^4 \end{aligned}$$




Parton Collider - I


Fall 2014

Dominant diagrams for H production:



Gluon-Gluon fusion



Vector Boson fusion



Higgsstrahlung



Parton Collider - II

Fall 2014

Basic ingredient: PDFs

Quarks: Look for heavy ones

Tree diagrams: Best bet is with b

Factor $\frac{m_b^2}{M_W^2} \sim 3 \cdot 10^{-3}$ encouraging

But: No b -quark beams, must rely on $b\bar{b}$ sea inside the nucleon

b -quark partonic density small...

Taking H production at small rapidity $y \sim 0$, with a 7 TeV beam $x \sim 10^{-2}$

→ Incident flux of sea b -quarks very small

Gluons: Main contribution


Loop diagrams, dominated by t quark

PDF somewhat dependent on Q^2

Parton Collider- III

Fall 2014

E.Menichetti - Universita' di Torino




Parton Collider- IV

Fall 2014

E.Menichetti - Universita' di Torino



67



Parton Collider - V


Fall 2014

Parton densities:



Parton Collider - VI


Fall 2014



Parton Collider - VII

Fall 2014


Expected cross sections for parton colliders:



Parton Collider - VIII

Fall 2014


Results for LHC cross sections



Parton Collider - IX

Fall 2014



Cross-sections for a 126 GeV Higgs



E.Menichetti - Universita' di Torino

H Searches - I

Direct searches at LEP I:
Higgsstrahlung
Z off-shell detected by lepton decay





Final LEP I result:
 $m_H < 65 \text{ GeV}$ excluded at 95% CL

H Searches - II

Fall 2014


Direct searches at LEP II:
Higgsstrahlung
Z on-shell detected by lepton decay



E.Menichetti - Universita' di Torino

H Searches - III

Fall 2014



Best channel:



$$H \rightarrow b\bar{b}$$

$$Z \rightarrow q\bar{q}$$

	Higgs 115 GeV	Z boson
qq		70
bb	74	15
cc	4	12
gg	6	0
$\ell^+\ell^-$		10
$\tau^+\tau^-$	7	3
$\nu\bar{\nu}$		20
WW*	8	
ZZ*	1	

H Searches - IV

Fall 2014



H Searches - V


Fall 2014

Likelihood ratio test

$$Q = \frac{L_{sign+bckg}}{L_{bckg}} \rightarrow -2 \ln Q \approx \Delta\chi^2$$

Final LEP II result:

$m_H > 114.4 \text{ GeV}$ at 95% CL



A false alarm:

'Signal' = Best fit

But: Excess at 115 GeV expected in 9% of cases from pure background

E.Menichetti - Universita' di Torino

H Searches - VI

Fall 2014

Tevatron direct searches:

Two large experiments, CDF & D0

Among tens of channels investigated, main results from:

$$q\bar{q} \rightarrow WH \rightarrow l\nu b\bar{b}$$

$$gg \rightarrow H \rightarrow WW \rightarrow l\nu l\nu$$

Rather complex topologies, heavy use of *neural networks*

Sophisticated, parallel logic networks capable of handling many parameters in order to select candidate events

Can be 'trained' by tuning selection criteria across Montecarlo samples of signal and background

H Searches - VII


Fall 2014

CDF :

$$q\bar{q} \rightarrow WH \rightarrow l\nu b\bar{b}$$

Neural Network tagging

LEP 'indication' at 115 GeV: Expect 4.8 events, observe 5.6



H Searches - VIII


Fall 2014

CDF :

$H \rightarrow WW \rightarrow l\nu l\nu$

Neural Network tagging

Consistent with $t\bar{t}$ background




H Searches - IX

Fall 2014

Main Tevatron result : "Brazilian Flag" plot

Continuous line: Upper limit of (Higgs+SM Background) observed
Dashed line: SM Background ($\pm 1\sigma$, $\pm 2\sigma$) expected
Unit: Expected SM Higgs signal as a function of m_H




H Searches - X

Fall 2014

Indirect searches:

Radiative corrections to many EW observables receiving contributions $\propto \ln m_H, m_t^2$

Ex: m_W





H Searches - XI

Fall 2014

Global fit to ~ 20 EW observables:


Best indirect estimate of m_H



H Searches - XII

Fall 2014


χ^2 minimization:



$$m_H = 90^{+36}_{-27} \text{ GeV}, \text{ } 68\% \text{ CL}$$

LHC: Machine - I

Fall 2014



E.Menichetti - Universita' di Torino

LHC: Machine - II

Fall 2014

LHC dipole field 8.3 T

(HERA/Tevatron ~4 T)


LHC pp $\sim 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

(Tevatron pp $3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$)

(SppbarS pp $6 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$)

LHC Layout

- 8 arcs.
- 8 straight sections (LSS),
~ 700 m long.
- The beams exchange their
positions (inside/outside) in 4
points to ensure that both
rings have the same
circumference !



E.Menichetti - Universita' di Torino

LHC: Machine - III

Fall 2014

$R = L\sigma$ Rate, Luminosity,Cross-Section

$$L = \frac{kN^2 f}{4\pi\sigma_x^* \sigma_y^*}$$

k = number of bunches = 2808

N = no. protons per bunch = 1.15×10^{11}

f = revolution frequency = 11.25 kHz

σ_x^*, σ_y^* = beam sizes at collision point (hor./vert.) = 16 mm

High L:

Many bunches (k)

Many protons per bunch (N)

A small beam size $\sigma_u^* = (\beta^* \epsilon)^{1/2}$

β^* : Beam envelope (optics)

ϵ : Phase space volume occupied
by the beam (constant along the ring)

High beam “brilliance” N/ϵ
(particles per phase space volume)
→ Injector chain performance

Small envelope
→ Strong focusing

Optics property

Beam property

LHC: Machine - IV

Fall 2014




$$B\rho = \frac{mv}{e} = \frac{p}{e}$$

LHC: $\rho = 2.8$ km given by LEP tunnel

To reach $p = 7$ TeV/c given a bending radius of $\rho = 2805$ m:


Bending field : $B = 8.33$ T

→ Superconducting magnets



LHC: Machine - V

Fall 2014




E.Menichetti - Universita' di Torino

LHC: Machine - VI

Fall 2014

Superconducting coils:





E.Menichetti - Universita' di Torino

LHC: Machine - VII

Fall 2014

LHC main dipole:

Two magnets in a single module



LHC: Machine - VIII

Fall 2014


RF system:

4 + 4 Superconducting RF cavities

400 MHz

~ 0.5 MeV/turn

20 minutes for 450 GeV \rightarrow 7 TeV



LHC @ 3.5 TeV

LHC @ 7 TeV

LEP @ 104 GeV

Synchrotron
radiation loss

0.42 keV/turn


6.7 keV /turn

~ 3 GeV /turn

LHC: Machine - IX

Fall 2014

Superconducting cavity



H - I

Selecting best decay channels for detection:

Strongly dependent on (unknown) M_H

By taking $M_H < 2M_W$

$b\bar{b}$: Large $BR > 50 \%$, good signature (secondary vertexes), *lots* of QCD background

$\tau^+\tau^-$: Large $BR \sim 7 \%$, somewhat harder than $b\bar{b}$ (neutrinos)

$\gamma\gamma$: Tiny $BR \sim 2 \cdot 10^{-3}$, small background, experimentally challenging

gg : Large $BR \sim 5 \%$, 2 jets, *lots* of QCD background

ZZ^* : Small $BR \sim 3 \%$, small background in the 4 leptons mode



WW^* : Large $BR \sim 20 \%$, sizeable QCD background in the 4 jets mode, harder than ZZ^* in leptonic modes (neutrinos)

H - II

Fall 2014

Total integrated luminosity:

$\sim 30 \text{ fb}^{-1}$ / experiment





Phenomenal performance:

- Record luminosity ($> 5 \times 10^{33}$) obtained soon after startup in 2012
- Sustained data collection rate of $> 1.0 \text{ fb}^{-1} / \text{wk}$
- Delivered/recoded @ 8 TeV = [23.3 / 21.3 (ATLAS) , 21.8 (CMS)] fb⁻¹


H - III

4 leptons:
 $\sim 7\sigma$ observation



H - IV

2 γ 's:
 ~6 σ observation



H - V

Signal strength:

	ATLAS (expected)	ATLAS (observed)	CMS (expected)	CMS (observed)
$h \rightarrow \gamma\gamma$	4.1	7.4	5.2	5.7
$h \rightarrow ZZ$	4.4	6.6	6.7	6.8
$h \rightarrow WW$	3.7	3.8	5.8	4.3
$h \rightarrow \tau\tau$	3.2	4.1	3.6	3.4
$h \rightarrow bb$	1.6	~0	2.1	2.1

H - VI

Fall 2014

Combined mass: All modes

$$m_H \text{ (ATLAS)} = 125.36 \pm 0.37 \text{ (stat)} \pm 0.18 \text{ (syst)}$$

$$m_H \text{ (CMS)} = 125.03^{+0.26}_{-0.27} \text{ (stat)}^{+0.13}_{-0.15} \text{ (syst)}$$


Many more results on spin/parity, couplings, width...

....Next time!

Future Collider Comparison - I

Fall 2014

Comparison of different colliders:



Future Collider Comparison - II

Fall 2014

Accelerator → Physical Quantity ↓	LHC 300 fb ⁻¹ /expt	HL-LHC 3000 fb ⁻¹ /expt	ILC 250 GeV 250 fb ⁻¹ 5 yrs	Full ILC 250+350+ 1000 GeV 5 yrs each	CLIC 350 GeV (500 fb ⁻¹) 1.4 TeV (1.5 ab ⁻¹) 5 yrs each	LEP3, 4 IP 240 GeV 2 ab ⁻¹ (*) 5 yrs	TLEP, 4 IP 240 GeV 10 ab ⁻¹ 5 yrs (*) 350 GeV 1.4 ab ⁻¹ 5 yrs (*)
N _H	1.7×10^7	1.7×10^8	6×10^4 ZH	10^3 ZH 1.4×10^5 Hvv	7.5×10^4 ZH 4.7×10^5 Hvv	4×10^5 ZH	2×10^6 ZH 3.5×10^4 Hvv
m _H (MeV)	100	50	35	35	100	26	7
$\Delta\Gamma_H / \Gamma_H$	--	--	10%	3%	ongoing	4%	1.3%
$\Delta\Gamma_{\text{inv}} / \Gamma_H$	Indirect (30%?)	Indirect (10%?)	1.5%	1.0%	ongoing	0.35%	0.15%
$\Delta g_{H\gamma\gamma} / g_{H\gamma\gamma}$	6.5 – 5.1%	5.4 – 1.5%	--	5%	ongoing	3.4%	1.4%
$\Delta g_{H\pi\pi} / g_{H\pi\pi}$	11 – 5.7%	7.5 – 2.7%	4.5%	2.5%	< 3%	2.2%	0.7%
$\Delta g_{H\eta\eta} / g_{H\eta\eta}$	5.7 – 2.7%	4.5 – 1.0%	4.3%	1%	~1%	1.5%	0.25%
$\Delta g_{HZZ} / g_{HZZ}$	5.7 – 2.7%	4.5 – 1.0%	1.3%	1.5%	~1%	0.65%	0.2%
$\Delta g_{HHH} / g_{HHH}$	--	< 30% (2 expts)	--	~30%	~22% (~11% at 3 TeV)	--	--
$\Delta g_{H\mu\mu} / g_{H\mu\mu}$	< 30%	< 10%	--	--	10%	14%	7%
$\Delta g_{H\tau\tau} / g_{H\tau\tau}$	8.5 – 5.1%	5.4 – 2.0%	3.5%	2.5%	$\leq 3\%$	1.5%	0.4%
$\Delta g_{Hcc} / g_{Hcc}$	--	--	3.7%	2%	2%	2.0%	0.65%
$\Delta g_{Hbb} / g_{Hbb}$	15 – 6.9%	11 – 2.7%	1.4%	1%	1%	0.7%	0.22%
$\Delta g_{Htt} / g_{Htt}$	14 – 8.7%	8.0 – 3.9%	--	5%	3%	--	30%