

Elementary Particles II

Higgs Short Notes

Higgs Field, Vacuum Nightmares, End of the World

Production, Decays

Strategies, Detectors

Observation

About the Higgs Field - I

Universal, constant field

Lorentz scalar → Same value in any frame, rotation invariant

Non-standard feature:

Vacuum expectation value $v \neq 0$

Usual analogy: Spontaneously magnetized ferromagnet:

$\mathbf{M} \neq 0$ below Curie temperature

→ Pick up a direction

Ground state rotationally not symmetric, in spite of H being symmetric


About the Higgs Field - II

Better analogy: Superconductor

Energy difference between normal and s.c. state at two different temperatures

$$\Delta E = a(T)|\psi|^2 + \frac{1}{2}b(T)|\psi|^4 + \dots \quad \text{Landau theory of phase transitions}$$

ψ is the Cooper pair ‘wave function’ $\rightarrow |\psi|^2 \sim \text{density of Cooper pairs}$



Below T_c , the minimum energy state (‘vacuum’) occurs for $\psi = \psi_0 \neq 0$, phase undefined

$\rightarrow U(1)$ QED gauge invariance spontaneously broken

\rightarrow Photon becomes ‘massive’ $\rightarrow B = 0$ inside

About the Higgs Field - III

ψ ‘Higgs field’ of superconductivity: $\langle \psi \rangle \neq 0 \leftrightarrow$ Permanent supercurrents

Superconductive state:

‘Higgs field’ = ‘Wave function’ of Cooper pairs

→ Not a fundamental field

→ ‘Composite’ field of fundamentals fermions (electrons)

Why there is the composite?

e - e effective interaction: Attractive (!) due to e – lattice interaction

Is the ‘real’ Higgs field a genuine, fundamental field or a composite?

Good question..No answer (yet):

Take it as a fundamental field

About the Higgs Field - IV

Fall 2020

A couple of questions:

1) What about the nonzero VEV of the Higgs field?

Higgs: Unique field whose VEV $\neq 0$

Similar to magnetization $\mathbf{M} \neq 0$ in a ferromagnet

But:

In a vacuum \rightarrow Not related to many body effects

Lorentz scalar \rightarrow No preferred direction, reference frame

2) Does it involve a new force? 'Giving mass to \approx all the fundamental constituents' ??

- Part of the standard EW interaction, often as a negligible contribution:

Higgs *particle* exchange diagrams between Fermion lines normally strongly suppressed

by m_f/m_W factors as compared to γ, Z^0, W^\pm exchange (Not true for t quark!)

3- & 4-boson diagrams with and without H similar

- Crucial role as 'Background' interaction:

For most particles Higgs *field* coupling translates into *inertial mass* !

About the Higgs Field - V

Fall 2020

Apparently contributing to vacuum energy density:

Beware: Take *potential energy* $V(\phi)$

$$\begin{aligned}V_{\min} = V(v) &= \frac{1}{2}\mu^2 v^2 + \frac{1}{4}\lambda v^4 \quad \text{use } \mu^2 = -\lambda v^2 \\&= -\frac{1}{4}\lambda v^4 \quad \text{use } m_h^2 = 2\lambda v^2 \\&= -\frac{1}{8}m_h^2 v^2\end{aligned}$$

Constant term: Usually not considered

Does not enter field equations, where only energy *differences* count

But: Taken into account by gravity → Cosmological term?

Cosmological constant: Possibly additional term in Einstein's field equations

May yield long range attraction/repulsion, according to sign

Invented by Einstein in order to guarantee static universes

Rejected by Einstein at the time of discovery of expansion of the Universe

Recently resurrected following the discovery of accelerated expansion

E.Menichetti - Universita' di Torino

About the Higgs Field - VI

Fall 2020

$$\text{Zero point energy} = -\frac{1}{8}m_h^2v^2 \sim \rho_{Higgs}$$

$$\text{Indeed: } \left[-\frac{1}{8}m_h^2v^2 \right] = \underbrace{E^4}_{GeV^4} = E(E^{-1})^{-3} \rightarrow \frac{E}{L^3} \quad \text{Energy density}$$

$$\rho_{Higgs} \sim 1.2 \cdot 10^8 \text{ GeV}^4$$

By assuming ρ to be a cosmological term, compare:

$$\rho_{observed} \sim 10^{-47} \text{ GeV}^4 !$$

$\rightarrow \rho_{Higgs}$ 55 orders of magnitude too big

(and with the wrong sign...)

Quick fix: $V(v)$ can be set = 0 by adding a constant to $V(\phi)$

Constant apparently unrelated to $m_h, v...$

...to be chosen to an accuracy of 1 part out of 10^{55} !

Fine tuning problem, still essentially unsolved

Something missing?

About the Higgs Field - VII

Fall 2020

Higgs boson: Quantum excitation of the field, mass m_H *not* given by the field

Further issue: $V(\phi)$ appearing in L : *Classical* potential

→ Must be quantized

→ Will be used perturbatively

→ Radiative corrections will modify the classical $V(\phi)$

Similar to vacuum polarization corrections to Coulomb potential in QED
(Uehling potential & Lamb shift)

Standard effect: Running constants, including λ

$$L_H = D_\mu \phi^\dagger D^\mu \phi - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2$$

$$\lambda = \lambda(q^2)$$

→ m_H modified by radiative corrections

Upon taking $\mu^2 < 0$, $\lambda(0) > 0$

→ λ evolution depending on β -functions

About the Higgs Field - VIII

Running couplings and β -functions:

$$\frac{dg^2}{d\ln Q^2} \equiv 4\pi\beta(g^2) = \underbrace{bg^4}_{1 \text{ loop}} + \underbrace{O(g^6)}_{2 \text{ loop}} + \dots$$

$$\rightarrow \frac{dg_i^2}{d\ln Q^2} = 4\pi\beta_i(g_i^2) \simeq b_i g_i^4$$


For the EW interaction:

$$b_g = -\frac{19}{6 \cdot 16\pi^2}, \quad b_{g'} = +\frac{41}{6 \cdot 16\pi^2}$$

Higgs couplings:

$$\frac{d\lambda}{d\ln Q^2} = \frac{1}{32\pi^2} \left[24(\lambda^2 + h_t^2 - h_t^4) - 3\lambda(3g^2 + g'^2) + \frac{3}{8}(2g^4 + (g^2 + g'^2)^2) \right] \text{ Self}$$

$$\frac{dh_t}{d\ln Q^2} = \frac{1}{32\pi^2} \left[9h_t^3 - h_t \left(8g_s^2 + \frac{9}{4}g^2 + \frac{17}{12}g'^2 \right) \right] \text{ Top (Yukawa)}$$



About the Higgs Field - IX

$$\frac{d\lambda}{d \ln Q^2} \sim \frac{3\lambda^2}{4\pi^2} \text{ Neglect smaller contributions at large } \lambda$$

$$\frac{d\lambda}{\lambda^2} \sim \frac{3}{4\pi^2} d \ln Q^2 \rightarrow \frac{1}{\lambda(Q^2)} + \frac{1}{\lambda(\nu^2)} \sim \frac{3}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

$$\rightarrow \frac{1}{\lambda(Q^2)} \sim \frac{1}{\lambda(\nu^2)} - \frac{3}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

$$\lambda(\nu^2) = \frac{G_F m_H^2}{\sqrt{2}} \rightarrow \lambda(Q^2) \sim \frac{\lambda(\nu^2)}{1 - \frac{3}{4\pi^2} \lambda(\nu^2) \ln \frac{Q^2}{\nu^2}}$$

$$\lambda \rightarrow \infty \text{ as } \frac{3}{4\pi^2} \lambda(\nu^2) \ln \frac{Q^2}{\nu^2} \rightarrow 1 \quad \text{Diverging at 'Landau pole'} \quad Q_{LP} = v \exp \left(\frac{2\pi^2}{3\lambda(\nu^2)} \right) = v \exp \left(\frac{2\sqrt{2}\pi^2}{3G_F m_H^2} \right)$$

$$\rightarrow \text{New physics required at scale } \Lambda < Q_{LP} \rightarrow \ln \frac{\Lambda}{v} < \left(\frac{2\sqrt{2}\pi^2}{3G_F m_H^2} \right)^{1/2} \rightarrow m_H < \left(\frac{2\sqrt{2}\pi^2}{3G_F \ln \frac{\Lambda}{v}} \right)^{1/2} \sim \begin{cases} O(140 \text{ GeV}), \Lambda \sim m_{Planck} \approx 1.2 \cdot 10^{19} \text{ GeV} \\ O(650 \text{ GeV}), \Lambda \sim 1 \text{ TeV} \end{cases}$$

About the Higgs Field - X

$$\frac{d\lambda}{d \ln Q^2} \sim -\frac{3h_t^4}{4\pi^2} \text{ Neglect smaller contributions at small } \lambda$$

$$\rightarrow d\lambda \sim -\frac{3h_t^4}{4\pi^2} d \ln Q^2$$

$$\rightarrow \lambda(Q^2) \sim \lambda(\nu^2) - \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2}$$

λ must stay +ve in order to keep vacuum stable (!):

Don't like a too quick End of the World



$$\rightarrow \lambda(\nu^2) > \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2} \rightarrow \frac{G_F m_H^2}{\sqrt{2}} > \frac{3h_t^4}{4\pi^2} \ln \frac{Q^2}{\nu^2} \text{ for some } Q \sim \Lambda$$

$$\rightarrow m_H > \left(\frac{3h_t^4}{\sqrt{2}\pi^2 G_F} \ln \frac{\Lambda}{\nu} \right)^{1/2}$$


About the Higgs Field - XI

Running couplings:

Sombrero: $\lambda > 0$
Relax



Dog Bowl: $\lambda < 0$
End of the World
(sometime)




About the Higgs Field - XII

Radiative corrections leading to major changes in the effective Higgs potential at large ϕ values:


Details tied to m_H, m_t

Might induce vacuum instability/metastability through fast/slow tunneling



About the Higgs Field - XIII


Upper & lower bounds on m_H :




Higgs Production - I

Fall 2020

Start from H coupling to Fermions:



Compare to coupling to Z^0 :




$\rightarrow H$ coupling down by a factor $\sim \frac{m_f}{m_W}$ as compared to Z^0

Higgs Production - II

First mode:

s-channel formation:



Ideal for lineshape scan, provided cross-section is big enough

Lepton colliders:


Tough requirements on luminosity, energy resolution

Higgs Production - III

Fall 2020

Second mode:

H radiation from quarks, sizeable contribution from Top:



$t\bar{t}$ signature might be useful to tag

Higgs Production - IV

Fall 2020

Shift to gauge bosons:

Exclude massless photon, gluon *at tree level*

[Photon, gluon *loop* contribution to be taken into account: See later]

More promising: W, Z mass very large

A Feynman diagram showing the annihilation of two virtual photons (V^μ and V^ν) into a Higgs boson (H). The interaction is represented by a dashed line. The equation associated with this diagram is:

$$H = 2i \frac{M_V^2}{v} g^{\mu\nu}$$


A Feynman diagram showing the annihilation of a virtual photon (V^μ) and a virtual Z boson (V^ν) into a Higgs boson (H). The interaction is represented by a dashed line. The equation associated with this diagram is:

$$= 2i \frac{M_V^2}{v^2} g^{\mu\nu}$$

Higgs Production - V

Best modes:

'Higgsstrahlung', 'Gauge boson fusion'




Higgs Production - VI

Fall 2020

Beyond tree level: Very Important Loops


Lepton machines:

Interesting diagrams, also quite relevant to detection




Parton machines:


Dominant diagram at LHC



Higgs Decays - I




$$\Gamma(H \rightarrow f\bar{f}) = N_C \frac{1}{8\pi} \frac{m_f^2}{v^2} M_H \left(1 - \frac{4m_f^2}{M_H^2}\right)^{3/2}$$




$$\begin{aligned} \Gamma(H \rightarrow WW) &= \frac{1}{4\pi} \frac{M_W^4}{M_H v^2} \left(1 - \frac{4M_W^2}{M_H^2}\right)^{1/2} \left(3 + \frac{1}{4} \frac{M_H^4}{M_W^4} - \frac{M_H^2}{M_W^2}\right) \\ \Gamma(H \rightarrow ZZ) &= \frac{1}{8\pi} \frac{M_Z^4}{M_H v^2} \left(1 - \frac{4M_Z^2}{M_H^2}\right)^{1/2} \left(3 + \frac{1}{4} \frac{M_H^4}{M_Z^4} - \frac{M_H^2}{M_Z^2}\right) \end{aligned}$$

Higgs Decays - II


Fall 2020



$$\Gamma(H \rightarrow gg) = \frac{M_H^3}{8\pi v^2} \left(\frac{\alpha_s}{\pi} \right)^2 n^2 |D(n)|^2$$




Higgs Decays - III



$$\Gamma(H \rightarrow W tb) = N_C \frac{g^2}{2v^2} \frac{M_W^4}{M_H} |V_{tb}|^2 \int dQ_3 \frac{G^{\beta\nu} T_{\beta\nu}}{[s_{34} - M_W^2]^2}$$

3-body phase space factor




$$\Gamma(H \rightarrow W f_u f_d) = \frac{g^2}{v^2} \frac{3M_W^2}{256\pi^3} M_H S(x)$$

3-body phase space factor

Higgs Decays - IV


Fall 2020



$$\Gamma(H \rightarrow Z f\bar{f}) = \frac{g^2}{v^2} \frac{3M_Z^2}{256\pi^3} M_H S(x) \frac{R(\theta_w)}{\cos^2 \theta_w} \left(\frac{7}{12} - \frac{10}{9} \sin^2 \theta_w + \frac{40}{27} \sin^4 \theta_w \right)$$


3-body phase space factor

$$\Gamma(H \rightarrow Z/\gamma, \gamma)$$




Higgs Decays - V

H decays entirely determined by Higgs mass:




Higgs Decays - VI

H branching ratios:



Higgs Decays - VII


Fall 2020



Parton Collider - I

Fall 2020

Dominant diagrams for H production:



Parton Collider - II

Basic ingredient: PDFs

Quarks: Look for heavy ones

Tree diagrams: Best bet is with b

Factor $\frac{m_b^2}{M_W^2} \sim 3 \cdot 10^{-3}$ encouraging

But: No b -quark beams, must rely on $b\bar{b}$ sea inside the nucleon

b -quark partonic density small...

Taking H production at small rapidity $y \sim 0$, with a 7 TeV beam $x \sim 10^{-2}$


→ Incident flux of sea b -quarks very small

Gluons: Main contribution


Loop diagrams, dominated by t quark

PDF somewhat dependent on Q^2

Parton Collider- III




Parton Collider- IV




Parton Collider - V

Parton densities:



Parton Collider - VI


Fall 2020



Parton Collider - VII

Fall 2020


Expected cross sections for parton colliders:



Parton Collider - VIII


Fall 2020

Results for LHC cross sections




Parton Collider - IX

Cross-sections for a 126 GeV Higgs



LHC: Machine - I



LHC: Machine - II


LHC dipole field 8.3 T
(HERA/Tevatron ~4 T)

LHC pp $\sim 10^{34}$
 $\text{cm}^{-2} \text{s}^{-1}$

(Tevatron pp 3×10^{32}
 $\text{cm}^{-2} \text{s}^{-1}$) (SppbarS pp
 $6 \times 10^{30} \text{ cm}^{-2} \text{s}^{-1}$)

LHC Layout

- ❑ 8 arcs.
- ❑ 8 straight sections (LSS),
~ 700 m long.
- ❑ The beams exchange their
positions (inside/outside) in 4
points to ensure that both
rings have the same
circumference !



LHC: Machine - III

$R=L\sigma$ Rate, Luminosity,Cross-Section

$$L = \frac{kN^2 f}{4\pi\sigma_x^* \sigma_y^*}$$

k = number of bunches = 2808

N = no. protons per bunch = 1.15×10^{11}

f = revolution frequency = 11.25 kHz

σ_x^*, σ_y^* = beam sizes at collision point (hor./vert.) = 16 mm

High L:

Many bunches (k)

Many protons per bunch (N)

A small beam size $\sigma_u^* = (\beta^* \varepsilon)^{1/2}$

β^* : Beam envelope (optics)

ε : Phase space volume occupied
by the beam (constant along the
ring)

-
- High beam “brilliance” N/ε
(particles per phase space volume)
→ Injector chain performance
 - Small envelope
→ Strong focusing
Optics
 - property
 - Beam
 - property

LHC: Machine - IV

Fall 2020



$$B\rho = \frac{mv}{e} = \frac{p}{e}$$

LHC: $\rho = 2.8$ km given by LEP tunnel

To reach $p = 7$ TeV/c given a bending radius of $\rho = 2805$ m:


Bending field : $B = 8.33$ T

→ Superconducting magnets



LHC: Machine - V

Fall 2020




E.Menichetti - Universita' di Torino

41

LHC: Machine - VI

Superconducting coils:





LHC: Machine - VII

Fall 2020

LHC main dipole:

Two magnets in a single module



LHC: Machine - VIII


RF system:

4+4 Superconducting RF cavities

400 MHz

~ 0.5 MeV/turn



20' for 450 GeV \rightarrow 7 TeV



	Synchrotron radiation loss
LHC @ 3.5 TeV	0.42 keV/turn
LHC @ 7 TeV	6.7 keV /turn
LEP @ 104 GeV	~ 3 GeV /turn

LHC: Machine - IX

Superconducting cavity



H - |

Selecting best decay channels for detection:



Strongly dependent on (unknown) M_H

By taking $M_H < 2M_W$

- $b\bar{b}$: Large $BR > 50\%$, good signature (secondary vertexes), *lots* of QCD background
- $\tau^+\tau^-$: Large $BR \sim 7\%$, somewhat harder than $b\bar{b}$ (neutrinos)
- $\gamma\gamma$: Tiny $BR \sim 2 \cdot 10^{-3}$, small background, experimentally challenging
- gg : Large $BR \sim 5\%$, 2 jets, *lots* of QCD background
- ZZ^* : Small $BR \sim 3\%$, small background in the 4 leptons mode
- WW^* : Large $BR \sim 20\%$, sizeable QCD background in the 4 jets mode, harder than ZZ^* in leptonic modes (neutrinos)

H - II

Total integrated luminosity: $\sim 30 \text{ fb}^{-1}$ / experiment



Phenomenal performance:

Record luminosity ($> 5 \times 10^{33}$) obtained soon after startup in 2012




Sustained data collection rate of $> 1.0 \text{ fb}^{-1} / \text{wk}$

Delivered/recoded @ 8 TeV = [23.3 / 21.3 (ATLAS) , 21.8 (CMS)] fb⁻¹

H - III

Fall 2020



4 leptons: $\sim 7\sigma$ observation



H - IV

Fall 2020

2γ 's: $\sim 6\sigma$ observation



$H - V$

Signal strength:

	ATLAS (expected)	ATLAS (observed)	CMS (expected)	CMS (observed)
$h \rightarrow \gamma\gamma$	4.1	7.4	5.2	5.7
$h \rightarrow ZZ$	4.4	6.6	6.7	6.8
$h \rightarrow$ WW	3.7	3.8	5.8	4.3
$h \rightarrow \tau\tau$	3.2	4.1	3.6	3.4
$h \rightarrow bb$	1.6	~ 0	2.1	2.1

H - VI

Combined mass: All modes

$$m_H(\text{ATLAS}) = 125.36 \pm 0.37 \text{ (stat)} \pm 0.18 \text{ (syst)}$$

$$m_H(\text{CMS}) = 125.03^{+0.26}_{-0.27} \text{ (stat)}^{+0.13}_{-0.15} \text{ (syst)}$$

Many more results on spin/parity, couplings, width...

....Next time!