2F, 4F AND ALL THAT

Giampiero PASSARINO[ab],

CERN March 12, 1999

[a] Dipartimento di Fisica Teorica, Università di Torino, Torino, Italy

[b] INFN, Sezione di Torino, Torino, Italy

Summary

Why are we here if our MC doesn't start with K?

- Introduction to the problem.
- What is available and what is not.

PROTOTYPE of comparisons for 2F calculations can be found in hep-ph/9902452, D. Bardin, M. Grünewald and G. P., based on `TOPAZ0` versus `ZFITTER`

This is LEP 1 with a particular emphasis on demanding the FOLLOWING CRITERIA:

- **Comparisons, after any upgrading, should be consistently better than what they where in earlier studies.**

- **AT THE PEAK all relative deviations among TOTAL CROSS-SECTIONS and absolute deviations among ASYMMETRIES should be below $\boxed{0.1\ \textbf{per-mill}}$.**

- **AT THE WINGS, typically $\sqrt{s} = M_z^2 \pm 1.8\,$GeV, they should be below $\boxed{0.3\ \textbf{per-mill}}$.**

A new 2F comparison for LEP 2 energies, upgrading the one performed in F. Boudjema et al., ***Report of the Workshop in Physics at LEP2*****, must be started**

TH problems and EXP questions:

- **I/F INTERFERENCE is currently treated dubiously by the experiments: The MC used to extrapolate for efficiency and possibly acceptance do not contain I/F interference (eg, KO-RALZ in multiphoton mode). Thus extrapolated and quoted results somehow miss I/F interference.** `TOPAZ0` **and** `ZFITTER` **will extend their comparison to LEP 2 (+ activity of Zeuthen group)**

- **There is a claim of some serious discrepancy between** `KORAL` / `TOPAZ0` / `ZFITTER` **in** $\sigma_{\rm int}, A^{\rm int}_{\rm FB}$ **at** $M^2_{_Z} - 3 \le \sqrt{s} \le M^2_{_Z} + 3\,$**GeV for muons. The same applies at LEP 2. It has to do with** *exponentiation* **of** `ISR` $\otimes$ `FSR`

- **ISPP is the less satisfactory ingredient among all radiative corrections. What we have now is KKKS or JMS.**

$\boxed{\text{The total hadronic cross section}}$ is the sum of the $\boxed{\text{two fermion cross-section}}$ plus $\boxed{\sigma(\overline{q}qX)}$

Its is therefore an inclusive measurement of hadron production in e^+e^--annihilation in which production thresholds can be seen, e.g. $\boxed{W\text{-pair production}}$ with at least one of the W bosons decaying hadronically, or $\boxed{ZZ}$ or $\boxed{\text{other background}}$

COMMENT: The following result is obtained with codes which are based on the *CC03* or *NC02*, $\boxed{\text{double-resonant}}$, approximation. $\boxed{\text{Our goal}}$ will be the *perfect* definition of $\sigma(\overline{q}qX)$.

What has been done so far in the $\boxed{\text{Experimental Collaborations?}}$

- **Hadronic events are selected based on $\boxed{\text{FS particle multiplicity}}$ in the detector, so both**
- **genuine high energy $\overline{q}q$ events,**
- **radiative returns and**
- **4F hadronic events**

are selected for the $\boxed{\text{hadronic lineshape}}$

This gives the $\boxed{\text{total sample}}$: the $\boxed{\text{background}}$ is subtracted on a MC basis, using *CC03* $WW \to$ all, and *NC02* $ZZ \to$ all and hadronic two-photon collisions. After making a cut for non-radiative events, the hadronic cross-section at $\sqrt{s} = 161.3\,$Gev is $35.3\,$pb $(s'/s > 0.8)$. The principal background is an estimated $2.6\,$pb arising from 4F events.

Clearly the above strategy is $\boxed{\text{good enough}}$ so far, but wrong $\boxed{\text{in principle}}$

The interplay between $\boxed{\text{2F}}$ and $\boxed{\text{4F}}$ FS in e^+e^- annihilation at LEP 2.

There are $\boxed{\text{several components}}$ in the radiative corrections to $\boxed{\text{fermion pair production}}$: the less satisfactory, from a theoretical point of view, is IS (or FS) $\boxed{\text{fermion-pair production}}$.

$\boxed{\textbf{For definiteness consider } e^+e^- \to \bar{b}b \textbf{ with radiation of an } e^+e^- \textbf{ pair}}$

The $\boxed{\text{background}}$ is therefore represented by the full 4F process the so called *NC48* process, which is built from $\boxed{\text{48 Feynman diagrams}}$.

For studies around the Z resonance we only included pairs from initial state and a $\boxed{\text{cut was selected}}$ so that $M(\overline{b}b) > 0.25\,s$.

Therefore, the $z_{\rm min}$ cut is on the Q^2 of the $\boxed{\text{primary pair}}$. We don't know what to do if somebody wants to cut in q^2 of the $\boxed{\text{secondary pair}}(e^+e^-)$.

At $\boxed{\text{LEP2 energies or higher}}$ we need a $\boxed{\text{more precise}}$ separation between

- **radiative corrections to 2F production and**

- **real 4F events**



We will denote the $\boxed{\text{evaluation}}$ of any one-loop corrected cross section, e.g. $e^+e^- \to b\bar{b}$ as a $\boxed{\text{2F calculation}}$. A typical example, IS pair production, is shown in figure

By $\boxed{\text{4F calculation}}$ we mean the $\boxed{\text{tree level}}$ evaluation, e.g. $e^+e^- \to b\bar{b}e^+e^-$.

- **We note that the soft pairs $\gamma^* \to e^+e^-$ are $\boxed{\text{divergent}}$ in the limit of zero e^+e^- invariant mass and therefore any simulation of very soft pairs with MASSLESS 4F calculations is bound to produce wrong results.**



Therefore we may ask the question: $\boxed{\text{is a massive 4F calculation enough?}}$ No, because if pairs are soft enough (e^+e^- pairs down to $M(e^+e^-) = 2\,m_e$)

- **then we must include VIRTUAL PAIRS as well, as shown for instance in figure.**

$\boxed{\text{Are soft+virtual I/F pairs in a 2F calculation enough?}}$

No upper cut is imposed on the invariant mass $M(e^+e^-)$, so that all pairs compatible with the request $M(\bar{b}b) >$ some thresholds are accepted.

Thus there is more than $\boxed{\text{S+V I/F PAIRS}}$, there are many topologies for $\boxed{\text{H pairs}}$ and some of them requires a finite m_e also for H pairs. Indeed in *NC48* there are MULTI-PERIPHERAL diagrams which diverge for $m_e \to 0$.

$\boxed{\sigma(\bar{q}qX) \text{ requires}}$

A **Include VIRTUAL+SOFT (up to some invariant mass Δ) I/F state pairs with a complete 2F process calculation.**

B **The contributions to $e^+e^- \to \bar{b}be^+e^-$ not in [A] are then included with a constraint on $M(\bar{b}b)$, but with no further restriction on $M(e^+e^-)$, with a complete (i.e. fully massive) 4F calculation.**

C **The contributions to $e^+e^- \to \bar{b}be^+e^-$ that are already contained in [A], are included with $M(e^+e^-) > \Delta$.**

STEP B REQUIRES EVALUATION OF THE FOLLOWING 4F PROCESSES

– **FULLY HADRONIC, with at least one invariant mass passing the cut, i.e.**

1. **All CC11 processes, like** $d\overline{u}c\overline{s}$ **and flavor permutations**
2. **All NC32 processes, like** $s\overline{s}u\overline{u}$**, etc.**
3. **All Mix43 processes, like** $d\overline{u}u\overline{d}$**, etc.**
4. **All NC64 processes, like** $u\overline{u}u\overline{u}$**, etc.**

– **SEMI-LEPTONIC, with** $M(q_j\overline{q}_i)$ **passing the cut, i.e.**

1. **All CC11 processes, like** $\mu^-\overline{\nu}_\mu u\overline{d}$**, etc.**
2. **The CC20 process,** $e^-(e^+)\overline{\nu}_e(\nu_e)u(\overline{u})\overline{d}(d)$**.**
3. **All NC24 processes, like** $u\overline{u}\mu^+\mu^-$**, etc.**
4. **All NC19 processes, like** $u\overline{u}\nu_e\overline{\nu}_e$**, etc.**
5. **All NC48 processes, like** $u\overline{u}e^+e^-$**, etc.**

- **TO SUMMARIZE: around the $\boxed{Z \text{ resonance}}$ the rate for $\boxed{\text{Real}}$ and $\boxed{\text{Virtual}}$ radiation is known and included in the existing calculations.**

- **Both are enhanced by $\boxed{\text{large logarithms}}$ but they $\boxed{\text{cancel}}$ to a large extent, leading to a $\boxed{\text{small contribution}}$ to the inclusive decay rates.**

The $\boxed{\text{complete evaluation}}$ of $\sigma(\overline{q}qX)$ would be relatively easy if we could separate sub-classes of diagrams, e.g. $\boxed{\text{primary}}$ from $\boxed{\text{secondary}}$ production.

For that it is necessary that the $\boxed{\text{INTERFERENCE}}$ between them be $\boxed{\text{zero}}$, or $\boxed{\text{very small}}$ or, at least $\boxed{\text{non-singular}}$

- **In the limit of $\boxed{\text{massless}}$ fermions, $\boxed{\text{singularities}}$ will arise from**

  1. **s-channel $\gamma(g)$**
  2. **t-channel γ with outgoing $e^{\pm}$ lost in the beam pipe.**

This consideration suggests the appropriate strategy:

- **EVERYTHING should be included**

- **classes of diagrams showing a MASS SINGULARITY must be included through some $\boxed{\textbf{analytical}}$ calculation which also accounts for $\mathcal{O}(\alpha^2)$ $\boxed{\textbf{virtual}}$ radiation.**

- **All INTERFERENCES exhibiting mass singularities belong to this category**

- **The rest, including most of the interferences, are accounted for by some (numerical) massless 4F calculation**

In order to identify mass singular contributions we are guided by $\boxed{\textbf{IR power-counting}}$, which is further simplified by recalling that a cross section is a sum of cut diagrams.

$\boxed{\text{EXAMPLES OF IR POWER-COUNTING}}$

1. A TRIVIAL ONE



- **Propagators are:**

$$
\begin{aligned}
& q^2 \\
(q_+ p_-)^2 + m^2 &= q^2 + 2\,p_- \cdot q \sim 2\,p_- \cdot q \\
(q_- p_+)^2 + m^2 &= q^2 - 2\,p_- \cdot q - \sim 2\,p_+ \cdot q \\
(q_+ p_- - q_+)^2 + M_V^2 &= q^2 - t + M_V^2 + 2\,(p_- - q_-) \cdot q
\end{aligned}
$$

- **behaving as**

$$
\begin{aligned}
\int \frac{d^4 q}{q^4}\ , &\qquad \textbf{IR divergent} \\
\int \frac{d^4 q}{q^5}\ , &\qquad \textbf{for} \quad t = M_V = 0.
\end{aligned}
$$

Connection with $\boxed{\text{Bhabha}}$ through K-diagrams



2. SINGLE PHOTON EMISSION



- **For IR power-counting the behavior is**

$$
\begin{aligned}
&= \int d^4Q d^4q \frac{1}{(q^2+m^2)^2} \delta(Q^2)\delta((Q+q)^2+m^2)\delta((q+p)^2+m^2), \\
&\sim \int d^4q\, \delta(q^2+m^2)\, \delta((q+p)^2+m^2) \int \frac{d^3Q}{Q} \frac{1}{(Q\cdot q)^2}, \\
&\sim \int \frac{dQ}{Q}, \qquad \textbf{IR singular}
\end{aligned}
$$

Only the sum over $\boxed{\text{all cuts}}$ is IR save

3. (Virtual) PAIR RADIATION



- **This $\mathcal{O}(\alpha^2)$ correction to the fermion form factor from the vacuum polarization is obtained from the $\mathcal{O}(\alpha)$ result by replacing**

$$
\begin{aligned}
\frac{\delta_{\mu\nu}}{q^2} &\to \frac{\delta_{\mu\alpha}}{q^2}\,(q^2\delta_{\alpha\beta} - q_\alpha q_\beta)\frac{\alpha}{\pi}\Pi(q^2)\frac{\delta_{\beta\nu}}{q^2}, \\
\Pi(q^2) &= q^2 \int_{4m_2^2}^{\infty} \frac{d\tau}{\tau}\,\frac{R(\tau)}{q^2+\tau-i\,\varepsilon}
\end{aligned}
$$

- **exchanging order of integration we obtain**

$$
\frac{\alpha}{\pi} \quad \int_{4m_2^2}^{\infty} \frac{d\tau}{\tau}\,\frac{\alpha}{\pi}\,F(q^2,\tau)
$$

- **where F is the vertex correction due to the exchange of a boson of mass $\sqrt{\tau}$. Therefore, $\boxed{\text{NO IR}}$ divergence arise, but the result will be nevertheless $\boxed{\text{singular}}$ in the limit $m_2 \to 0$. The four-particle cut must be added.**

4. (Real) PAIR RADIATION



- **Let** q **the momentum of the internal loop, then the result is proportional to**

$$
\begin{aligned}
& \sim \frac{1}{q^4}\, I_{\mu\nu} \\
I_{\mu\nu} &= -\frac{\pi}{2\, q^2}\, \sqrt{\lambda(-q^2, m_2^2, m_2^2)}\, \theta(-q^2 - 4m_2^2)\theta(q_0)\, (q^2\delta_{\mu\nu} - q_\mu q_\nu)
\end{aligned}
$$

- **the** $q_\mu q_\nu$ **part may be dropped because of gauge invariance**

- **NO** $\boxed{\textbf{IR}}$ **divergence but the result is, again, singular for** $m_2 \to 0$.

5. If a diagram contains a subpart



- **we get**

$$\frac{1}{q^2(q^2+M_Z^2)}\,\frac{\sqrt{\lambda}}{q^2}\,q^2\delta_{\mu\nu}$$

- **which is finite in the limit $m_2 \to 0$, or $\equiv$ the interference between γ-radiated and Z-radiated secondary pairs has no mass singularity**

$\boxed{\text{A SUMMARY of the present status}}$

- **FS radiation is treated $\boxed{\text{complementary}}$ to IS radiation, $\boxed{\text{NO}}$ interference term is included which is justified**

- **as long as the (dominant) QED reactions are considered and**

- **axial vector couplings from neutral currents are ignored**

On $\boxed{\text{TOP}}$ of the Z resonance the interference is indeed $\boxed{\text{vanishing}}$ small at $\mathcal{O}(\alpha)$

- **One expects the same at $\mathcal{O}(\alpha^2)$.**

- **$\boxed{\text{However}}$ well above the resonance the interference may well contribute**

- **It is known that the interference between $\boxed{\text{primary}}$ and $\boxed{\text{secondary}}$ quark production is $\boxed{\text{NOT}}$ enhanced. But**

- **What about the I/F interference? It looks like there are mass singularities**

To be more specific, let us consider $e^+e^- \to \overline{q}q\mu^+\mu^-$, the so called *NC24*, i.e. $3 \otimes 8$ diagrams, C for Conversion, A_q for primary $\overline{q}q$ production and A_μ for secondary $\overline{q}q$ production:



C



A_q



A_μ

In general we require $M^2(\overline{q}q) \geq s_0$. The following happens:

- $|C|^2$ **is IS PP,** $\boxed{\textbf{known}}$
- $|A_q|^2$ **is FS PP,** $\boxed{\textbf{known}}$

What about $\boxed{\textbf{INTERFERENCE}}$**? Few things are obviously known:**

- $C \otimes A_q$ **when one** μ**-pair is** γ**-radiated and the other is** Z**-radiated is** $\boxed{\textbf{non-singular}}$
- $|A_\mu|^2$ **is never singular because of the cut**

It remains the interference $(C \oplus A_q) \otimes A_\mu$ **and** $C \otimes A_q$ **with QED reactions. Here we know that:**

- $A_q \otimes A_\mu$ **induced by the** $\boxed{\textbf{vector}}$ **(em or neutral) current vanishes,**
- **induced by** $\boxed{\textbf{axial vector}}$ **is different from zero but** $\boxed{\textbf{NOT}}$ **mass-singular**

NEXT level is represented by $e^+e^- \to \overline{Q}Q\overline{q}q, Q \neq q$**, with primary** Q **and secondary** q**. There are** 8 **additional annihilation diagrams with** $\boxed{\textbf{gluons}}$

Interference terms are of two types:

1. CONTAINS



- $\boxed{\textbf{vanishes}}$ **because** $\propto$ **Tr**λ^a

2. CONTAINS



- **If one of the gluons is replaced by** γ/Z **the interference is zero**

- **Since the one loop vertex** $V = \gamma/Z \to gg \propto \delta^{ab}\, a_V$ **with** $a_\gamma = 0$ **and** $a_Z = I_f^3/(2\, s_\theta c_\theta)$**, we have:**

- $V = \gamma$ **does not contribute**

- $V = Z$**, for fixed** Q**,** $q = u$ **and** $q = d$ **will cancel in the total, but for** $q = b$

Even more complicated is the case $Q = q$. For each diagram we have 4 permutations and the interference terms are obtained by cutting the following diagrams

 $+\ldots$

- **For the SINGLET part the four-parton cut is computed (the one in common with $\overline{Q}Q\overline{q}q$)**

- **The NON-SINGLET part for $\overline{Q}Q\overline{q}q$ is also computed**

- **What about the NON-SINGLET part for $\overline{q}q\overline{q}q$?**

- **It is included in the classical result for the $\mathcal{O}(\alpha_S^2)$ NS, however, it is not known how to separate the four-quark states from the associated virtual corrections**

There are $\boxed{\text{other}}$ types of mass-singularities, those associated with t-channel photons. Example is $e^+e^- \to e^+e^-\overline{q}q$, the *NC48* multi-peripheral (MP) diagrams.

- **For $q = b$ at $200\,$GeV the double-resonant approximation $e^+e^- \to ZZ \to e^+e^-\overline{b}b$ is about $10^{-2}\,$pb, while the total MP is about $1\,$pb, compared with a $\sigma_{\rm had}$ of about $80 - 90\,$pb.**

- **This is $\boxed{\text{easy-to-handle}}$ from an experimental point of view. It is cause by hadronic two-photon collisions, where the $\overline{b}b$ system is created by the two photons radiated by e^+e^-. Thus the $\boxed{\text{bulk}}$ of the cross-section is around $10\,$GeV invariant $\overline{b}b$ mass and will $\boxed{\text{Not materialize}}$ into many particles**

- **$\boxed{\text{However}}$, particle multiplicity in the detector is not directly accessible when one deals with partons.**

The $\boxed{\textbf{Final}}$ example will the the famous *CC20*: events of the type $e^+e^- \to u\overline{d}, \oplus$ one electron (in the beam pipe) and a neutrino. They $\boxed{\textbf{Cannot}}$ be excluded from the $\boxed{\textbf{hadronic lineshape}}$. $\boxed{\textit{CC20}\ \textbf{is a box full of snakes}}$

$\boxed{\textbf{How}\ \textit{CC20}\ \textbf{enters into}\ \sigma_{\rm had}\textbf{?}}$ There are two regimes:

- **Large angle $\Longrightarrow$ included with a massless MC**
- **outgoing $e^-(e^+)$ within a narrow cone around the beam axis**

- **The *CC20* process at small angles is mostly something like**



$$X = \nu + u + \overline{d}$$

$\boxed{\text{Singularities in } CC20}$

- **outgoing $e^-(e^+)$ within a narrow cone around the beam axis. Best seen within an (improved) Weizsäcker-Williams - approximation (WW)for the inside cone and for the subset of diagrams which are mass-singular in the limit of zero scattering angle, i.e. those with topology:**



The $\boxed{\textbf{WW-approximation}}$ is defined by:

$$\int d\Phi_3\, \hat{L}_{\mu\nu}\, \hat{W}_{\mu\nu} \Longrightarrow -\hat{W}_1(0,\hat{y})\, f_\gamma(\hat{Q}^2,\hat{y},x_-)$$
$$\hat{W}_1(0,\hat{y}) = -\frac{1}{2}\int d\Phi_3\, \hat{W}_{\mu\mu}(0,\hat{y})$$

- **$\hat{W}_1$ is therefore $\propto$ the cross-section for $e^+\gamma \to e^+u\overline{d}$ (everything massless)**
- **f_γ is the photon $\boxed{\textbf{flux-function}}$, which in the presence of ISR reads**

$$f_\gamma \;=\; -m^2\,(1+x_-+2\,\frac{1-x_-}{\hat{y}}) + \hat{Q}^2\,(1-\frac{2}{\hat{y}}+\frac{2}{\hat{y}^2})$$

Therefore the $\boxed{\hat{Q}^2 \textbf{ integration}}$ can be performed analytically. The integrand is the sum of two terms $\propto$

$$\frac{1}{\hat{Q}^2} \to \ln \frac{\hat{Q}_c^2}{\hat{Q}_0^2}, \qquad \frac{m^2}{\hat{Q}^4} \to \frac{m^2}{\hat{Q}_0^2} + \ldots$$

The result contains the usual $\boxed{\textbf{logarithmic}}$ enhancement but also the $\boxed{\textbf{constant}}$ term

For $M_X = 0$, i.e. massless quarks and no cut on $M(\overline{d}u)$, $\Longrightarrow q^2$ $\boxed{\text{CROSSES ZERO}}$ even for finite m_e, giving some non-integrable singularity.

- **Once a cut on $M(\overline{d}u)$ is imposed we will have two $\boxed{\text{MP diagrams}}$**



+ crossed

When $q^2 = 0 \equiv e^-$ lost in the beam pipe and the (massless) u-quark is emitted $\parallel$ to the (quasi-real) $\gamma \Longrightarrow$ the internal quark propagator will blow up.

- **massive quarks** $\Longrightarrow \ln m_e^2 \ln m_q^2$
- **massless quarks** $\Longrightarrow \ln^2 m_e^2$
- **angular cut** $\Longrightarrow \ln m_e^2 \ln \theta_c$

A SUMMARY OF QUESTIONS

- **EXPERIMENTAL guidance is needed to separate** *signal* **from** *background*, **e.g. we need to know how** *Pair Production* **is defined:**
  1. **cut on PRIMARY PAIR?**
  2. **cut on SECONDARY PAIR?**
- **Is there a** *realistic* **support for a combined 2F/4F generator? If so a detailed list of requested performances MUST be produced. The possibility to separate the contributions that are regarded as either** *signal* **or** *background* **is algorithm-dependent.**
- **IMPLEMENTATION of p_t for ISR photons. The problem is that many searches rely upon missing p_t as a signature (especially for SUSY) and so if some p_t can be carried away by ISR it can fake such a** *signal*. **Two alternatives:**
  1. *A wrong p_t is better than no p_t,*
  2. *No p_t is better than a wrong p_t.*

  If p_t is needed then, does one really need to generate n-photons? Wouldn't be enough to have
  1. **corrections beyond leading order to the showering**
  2. **smoothly merging with the higher order calculation of single,hard photon emission?**
- **Concept of** *user-friendly* **generator. Everybody knows how to generate all channels in one stroke, but to do it with a** *complete* **MC is infinitely time consuming.**

- **It takes a mother to run a generator (?)**
We have compared the CC03 predictions from three different generators: KORALW 1.21, EXCALIBUR and GRACE. The level of agreement on the cross-section predictions is reasonably good (better than 0.7 %)

- **Exact definition of theoretical accuracy needed.**

- **The FUDGE FACTOR syndrome: is** $FUDGE \otimes\ WRONG$ **numerically** $\approx$ $RIGHT$ **good enough? If so, you still need to have one example of** $RIGHT$**. Is this the fate of producing** $RIGHT$**?**

PHYSICS WISHES:

- **Reliability at zero scattering angles - massive fermions. Under investigation and not yet extended to** $e\nu_e e\nu_e$**.**

- **Improvements in ISR (photons with** p_t**) and FSR. It would require a NLO generator.**

- $\mathcal{O}(\alpha)$ **including quadruple-gauge-boson vertex (with anomalous QGCs). Again a NLO generator.**

- **Non-factorizable (non-naive) QCD corrections to** $qqqq$ **and** $ql\nu$ **final state. Not fully available, cut-dependent, problematic with the resolved component of the photon.**



Figure 1: Relative deviations between TOPAZ0 and ZFITTER for hadronic cross-section in CA3 mode as a function of the Higgs boson mass. The curves correspond to a cut $s' = 0.01\, s$.



Figure 2: Relative deviation (percent) between exact Q^2-integrated flux function and its leading logarithmic approximation, as a function of $y = p_+ \cdot Q / p_+ \cdot p_-$.