CERN / LEP2 MC workshop, October '99

RacoonWW: Radiative corrections on $e^+e^- \to \mathbf{WW} \to 4\boldsymbol{f}$

Stefan Dittmaier

Universität Bielefeld, Germany

in collaboration with

A. Denner, M. Roth and D. Wackeroth

Contents

1 Strategy for $\mathcal{O}(\alpha)$ corrections in RacoonWW

1.1 General considerations

Real corrections — $e^+e^- \to 4f + \gamma$ at tree level

Exact calculations for all final states exist.

Virtual corrections — $e^+e^- \to 4f$ at one loop

Problems:

- Enormous complexity of full one-loop calculation:
  - number of diagrams $\sim \mathcal{O}(10^3\text{–}10^4)$ per final state,
  - many non-equivalent final states,
  - numerical problems with 5- and 6-point functions.
- No convincing procedure for gauge-invariant introduction of finite widths at one loop.

Viable solution: double-pole approximation.

$\Rightarrow$ Take only diagrams for $e^+e^- \to WW \to 4f$
with two nearly resonant W bosons !

- Considerable reduction in number of Feynman graphs.
- On-shell production and decay of W's as building block.
- Generic calculation for all final states possible.
- Expected uncertainty not too close to threshold:

$$\sim \mathcal{O}\left(\frac{\Gamma_W}{M_W}\frac{\alpha}{\pi}\ln(\cdots)\right) \sim 0.1\%.$$

1.2 The RacoonWW approach

Denner, S.D., Roth, Wackeroth '99

$e^+e^- \to 4f$ at tree level

Full matrix-element calculation with massless fermions.

$\mathcal{O}(\alpha)$ radiative corrections (RCs) to $e^+e^- \to WW \to 4f$

virtual

real

$e^+e^- \to 4f + \gamma$ at tree level:
Full matrix-element calculation
$\to$ Talk of M. Roth 6/99

$e^+e^- \to WW \to 4f$ at one loop:
Double-pole approximation

Factorizable corrections



Building blocks:
– RCs to on-shell production
– RCs to on-shell decay
But: respect spin correlations!

Non-factorizable corrections



etc.

Photon exchange between production and decay subprocesses ($E_\gamma \lesssim \Gamma_W$)

Note: full off-shell phase space everywhere !

1.3 Virtual factorizable corrections



on-shell production   on-shell decays

One-loop contribution to matrix element in DPA:

$$\mathcal{M}^{e^+e^-\to WW\to 4f}_{\text{virt,fact,DPA}} = \sum_{\lambda_+,\lambda_-} \frac{1}{(k_+^2 - M_W^2 + iM_W\Gamma_W)(k_-^2 - M_W^2 + iM_W\Gamma_W)} \times \Big(\delta\mathcal{M}^{e^+e^-\to W^+W^-} \mathcal{M}^{W^+\to f_1\bar{f}_2}_{\text{Born}} \mathcal{M}^{W^-\to f_3\bar{f}_4}_{\text{Born}} + \mathcal{M}^{e^+e^-\to W^+W^-}_{\text{Born}} \delta\mathcal{M}^{W^+\to f_1\bar{f}_2} \mathcal{M}^{W^-\to f_3\bar{f}_4}_{\text{Born}} + \mathcal{M}^{e^+e^-\to W^+W^-}_{\text{Born}} \mathcal{M}^{W^+\to f_1\bar{f}_2}_{\text{Born}} \delta\mathcal{M}^{W^-\to f_3\bar{f}_4} \Big).$$

One-loop corrections to subprocesses known:

$\delta\mathcal{M}^{e^+e^-\to W^+W^-}$ — Böhm et al. '88; Fleischer, Jegerlehner, Zralek '89

$\delta\mathcal{M}^{W\to f\bar{f}'}$ — Bardin, S. Riemann, T. Riemann '86; Jegerlehner '86; Denner, Sack '90

Full result for $\mathcal{M}^{e^+e^-\to WW\to 4f}_{\text{virt,fact,DPA}}$ in two ways: — Denner, S.D., Roth, Wackeroth '99

- combination of known results (Böhm et al., Denner/Sack),
- completely new and independent calculation of all loops.

$\Rightarrow$ Perfect numerical agreement!

On-shell projection of momenta:

$\delta\mathcal{M}^{e^+e^-\to W^+W^-}$ and $\delta\mathcal{M}^{W\to f\bar{f}'}$ require on-shell kinematics.

$\Rightarrow$ Define mapping of fermion momenta: $\quad k_i \;\to\; \hat{k}_i$

with: $\quad \hat{k}_+^2 = (\hat{k}_1+\hat{k}_2)^2 = M_W^2, \quad \hat{k}_-^2 = (\hat{k}_3+\hat{k}_4)^2 = M_W^2.$

Note: actual choice of mapping involves freedom,
but ambiguity is of neglected order $\mathcal{O}\left(\frac{\alpha}{\pi}\frac{\Gamma_W}{M_W}\right)$.

Possible choice: fix directions of W^+, f_1, and f_3.

$$\hat{k}_{+0} = \frac{1}{2}\sqrt{s}, \qquad \hat{\mathbf{k}}_+ = \frac{\mathbf{k}_+}{|\mathbf{k}_+|}\beta\frac{\sqrt{s}}{2}, \qquad \hat{k}_-^\mu = p_+^\mu + p_-^\mu - \hat{k}_+^\mu,$$

$$\hat{k}_1^\mu = k_1^\mu \frac{M_W^2}{2\hat{k}_+ k_1}, \qquad \hat{k}_2^\mu = \hat{k}_+^\mu - \hat{k}_1^\mu,$$

$$\hat{k}_3^\mu = k_3^\mu \frac{M_W^2}{2\hat{k}_- k_3}, \qquad \hat{k}_4^\mu = \hat{k}_-^\mu - \hat{k}_3^\mu.$$

Some details of the new calculation

$$\mathcal{M}^{e^+e^-\to WW\to 4f,\sigma}_{\text{virt,fact,DPA}} = \sum_n F^\sigma_n(s,\hat{t})\,\mathcal{M}^\sigma_n(p_\pm,\hat{k}_i,k_+^2,k_-^2)$$

Standard matrix elements $\mathcal{M}^\sigma_n(p_\pm,\hat{k}_i,k_+^2,k_-^2)$

contain spinor chains and W polarizations:

$$\mathcal{M}^\pm_1 = \bar{v}(p_+)\not\varepsilon^*_+(\hat{\not k}_+ - \not p_+)\not\varepsilon^*_-\tfrac{1}{2}(1\pm\gamma_5)u(p_-), \quad \text{etc.}$$

with "effective W polarization vectors"

$$\varepsilon^{*,\mu}_+ = \frac{e}{\sqrt{2}s_W}\frac{\bar{u}(\hat{k}_1)\gamma^\mu\omega_- v(\hat{k}_2)}{k_+^2 - M_W^2 + iM_W\Gamma_W}, \quad \varepsilon^{*,\mu}_- = \dots$$

$\Rightarrow$ W-spin correlations respected!

Invariant functions $F^\sigma_n(s,\hat{t})$

contain coupling structures and loop integrals.

Economic numerical evaluation:

Note: $\mathcal{M}^\sigma_n$ simple, F^σ_n complicated functions!
But: $F^\sigma_n(s,\hat{t})$ depend only on single phase-space variable
$\hat{t} = (p_+ - \hat{k}_+)^2 = f(\cos\theta)$ with $\theta = \angle(\mathbf{p}_\pm, \hat{\mathbf{k}}_\pm)$.

$\Rightarrow$ Evaluate integrals before phase-space integration:

$$c^\sigma_{n,l}(s) = \tfrac{2l+1}{2}\int_{-1}^{+1} d\cos\theta\, F^\sigma_n(s,\hat{t})\, P_l(\cos\theta), \quad l = 0,1,\dots$$

Reconstruct F^σ_n algebraically during phase-space integration:

$$F^\sigma_n(s,\hat{t}) = \sum_{l=0}^{\infty} c^\sigma_{n,l}(s)\, P_l(\cos\theta),$$

$\Rightarrow$ Gain in CPU time and stability for $\cos\theta \to \pm 1$!

1.4 Virtual non-factorizable corrections

One-loop contribution to matrix element in DPA:

$$\mathcal{M}^{e^+e^-\to WW\to 4f}_{virt,nonfact,DPA} = \delta\mathcal{M}^{e^+e^-\to 4f}\Big|_{\text{doubly-resonant part}} - \mathcal{M}^{e^+e^-\to WW\to 4f}_{virt,fact,DPA}$$
$$= \mathcal{M}^{e^+e^-\to WW\to 4f}_{Born,DPA}\;\delta_{virt,nonfact,DPA}.$$

Features of the correction factor $\delta_{virt,nonfact,DPA}$:

- gauge invariance by definition,
- non-trivial dependence on off-shellness $k_\pm^2$ of W bosons
  $\Rightarrow$ no factorization of Breit–Wigner-type resonances
  (=non-factorizability),
- free of mass singularities,
- compensates IR singularities of W bosons in $\mathcal{M}^{e^+e^-\to WW\to 4f}_{virt,fact,DPA}$.

Calculation of the correction factor $\delta_{virt,nonfact,DPA}$:

- most complicated part was known from calculation of the sum $\delta_{virt+real,nonfact,DPA}$,
  Melnikov,Yakovlev '96
  Beenakker,Berends,Chapovsky '97
  Denner, S.D., Roth '97
- remainder is calculated.
  Denner, S.D., Roth, Wackeroth '99

A representative set of diagrams for non-factorizable RCs:

Genuinely non-factorizable diagrams:







Diagrams contributing to factorizable and non-factorizable RCs:









1.5 Combination of virtual and real corrections

Singularity structure of virtual and real corrections:

- IR singularities due to exchange or emission of soft photons by external charged fermions:
  corrections like $\alpha \ln(m_\gamma) |\mathcal{M}_{\mathrm{Born}}|^2$.
  $\Rightarrow$ Complete cancellation between virtual and real parts!

- Mass singularities due to exchange or emission of collinear photons by light external charged fermions:
  corrections like $\alpha \ln(m_f) |\mathcal{M}_{\mathrm{Born}}|^2$.
  $\Rightarrow$ Partial cancellation between virtual and real parts!

Note:

- Virtual corrections calculated in DPA.
  $\Rightarrow$ Singularities related to $\mathcal{M}^{\mathrm{e}^+\mathrm{e}^-\to\mathrm{WW}\to 4f}_{\mathrm{Born,DPA}}$.

- Real corrections calculated from full matrix elements.
  $\Rightarrow$ Singularities related to $\mathcal{M}^{\mathrm{e}^+\mathrm{e}^-\to 4f}_{\mathrm{Born}}$.

$\Rightarrow$ Avoid mismatch in singularity cancellation!

Strategy for treatment of singularities

1. Determine singular behaviour of $|\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f\gamma}|^2$:

$$\int d\Phi_{4f\gamma}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f\gamma}|^2 \quad \sim \quad \int d\Phi_{4f\gamma}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2\, \delta_{\text{sub}}^{4f\gamma}.$$

2. Reintroduce masses m_γ and m_f in singular factor $\delta_{\text{sub}}^{4f\gamma}$.

3. Perform integration over photon phase space analytically:
   (at least over singular region)

$$\int d\Phi_{4f\gamma}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2\, \delta_{\text{sub}}^{4f\gamma} \quad = \quad \int d\Phi_{4f}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2 \otimes \delta_{\text{sub}}^{4f}.$$

   Remarks:

   - "$\otimes$" indicates a convolution over an auxiliary parameter.
   - δ_{sub}^{4f} contains singular terms like $\alpha\ln(m_\gamma)$ and $\alpha\ln(m_f)$.

4. Split δ_{sub}^{4f} into two parts:

$$|\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2 \otimes \delta_{\text{sub}}^{4f} = |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2\, \underbrace{\delta_{\text{sub},1}^{4f}}_{\text{related to virtual corr's}} + |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2 \otimes \delta_{\text{sub},2}^{4f}$$

5. Add following contribution to phase-space integral:

$$\begin{aligned}
&- \int d\Phi_{4f\gamma}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2\, \delta_{\text{sub}}^{4f\gamma} && \text{cancels singularities of real corr's} \\
&+ \int d\Phi_{4f}\, |\mathcal{M}_{\text{Born,DPA}}^{e^+e^-\to 4f}|^2\, \delta_{\text{sub},1}^{4f} && \text{cancels singularities of virtual corr's} \\
&+ \int d\Phi_{4f}\, |\mathcal{M}_{\text{Born}}^{e^+e^-\to 4f}|^2 \otimes \delta_{\text{sub},2}^{4f} && \text{remaining (mass) singularities} \\
&\sim \quad 0 && \text{i.e. non-doubly-resonant !}
\end{aligned}$$

Two explicit procedures:

Phase-space slicing

$\delta_{\mathrm{sub}}^{4f\gamma} \neq 0$ only for $E_\gamma < \Delta E$ or $\angle(\gamma, f) < \Delta\theta$.

$\Rightarrow$ Cutoff dependences:

- Numerical cutoff dependence of $\int \mathrm{d}\Phi_{4f\gamma}$.
- Explicit cutoff dependence of $\delta_{\mathrm{sub},2}^{4f}$.

$\Rightarrow$ Numerical cancellation for $\Delta E \to 0$, $\Delta\theta \to 0$.

Subtraction procedure

$\delta_{\mathrm{sub}}^{4f\gamma} \neq 0$ over complete phase space.

Features:

- No cutoff-dependent numerical integrations.

  $\Rightarrow$ Better convergence expected!
- Non-trivial construction of $\delta_{\mathrm{sub}}^{4f}$.

Convenient general approach:

dipole subtraction formalism, formulated for

- QCD with massless, unpolarized partons, Catani, Seymour '96,'97
- photon radiation off light S.D., Roth '99
  and massive, polarized fermions. S.D. '99

Final master formula:

$$
\begin{aligned}
\int \mathrm{d}\sigma \;=\; \frac{1}{2s}\Bigg\{ \; \int \mathrm{d}\Phi_{4f} \Big[\, & |\mathcal{M}^{\mathrm{e^+e^-}\to 4f}_{\mathrm{Born}}|^2 \\
& + 2\,\mathrm{Re}\Big((\mathcal{M}^{\mathrm{e^+e^-}\to\mathrm{WW}\to 4f}_{\mathrm{Born,DPA}})^* \, \delta\mathcal{M}^{\mathrm{e^+e^-}\to\mathrm{WW}\to 4f}_{\mathrm{virt,fact,DPA}} \\
& \qquad + |\mathcal{M}^{\mathrm{e^+e^-}\to\mathrm{WW}\to 4f}_{\mathrm{Born,DPA}}|^2 \, \delta_{\mathrm{virt,nonfact,DPA}} \Big) \\
& \Big] \\
+ \int \mathrm{d}\Phi_{4f\gamma} \quad & |\mathcal{M}^{\mathrm{e^+e^-}\to 4f\gamma}_{\mathrm{Born}}|^2 \Bigg\}.
\end{aligned}
$$

$$
\text{non-singular}\ \left\{\begin{array}{l} \\ \\ \\ +\ |\mathcal{M}_{\mathrm{Born,DPA}}^{\mathrm{e^+e^-\to WW\to 4}f}|^2\,\delta_{\mathrm{sub},1}^{4f} \end{array}\right.
$$

$$
\text{explicit mass singularities}\ \left\{\ +\ |\mathcal{M}_{\mathrm{Born}}^{\mathrm{e^+e^-\to 4}f}|^2 \otimes \delta_{\mathrm{sub},2}^{4f}\right.
$$

$$
\underbrace{\left[\qquad\qquad - |\mathcal{M}_{\mathrm{Born}}^{\mathrm{e^+e^-\to 4}f}|^2\,\delta_{\mathrm{sub}}^{4f\gamma}\right]}_{\text{non-singular}}
$$

Note:

Potentially large ISR logs $\alpha\ln(m_\mathrm{e}/E)$ contained in $|\mathcal{M}_{\mathrm{Born}}^{\mathrm{e^+e^-\to 4}f}|^2\delta_{\mathrm{sub},2}^{4f}$.

$\Rightarrow$ Leading ISR corrections to background diagrams included!

(at present for CC11 class)

2 Numerical results of RacoonWW

Total cross section for $\mathrm{e}^+\mathrm{e}^- \to \mathrm{WW} \to \tau^+\nu_\tau\mu^-\bar{\nu}_\mu$:



W-production angle distribution for $\mathrm{e}^+\mathrm{e}^- \to \mathrm{WW} \to \tau^+\nu_\tau\mu^-\bar{\nu}_\mu$:



W invariant-mass distributions

Photon recombination (modified TH prescription of $4f$ proposal):

1. Discard photons (not the events!) close to beam
   if $\angle(\gamma, \mathrm{beam}) < 5°$.

2. Recombination of photons with nearest final-state fermion

   - if $M(f+\gamma) < 5\,\mathrm{GeV}$ (TH bare), or
   - if $M(f+\gamma) < 25\,\mathrm{GeV}$ (TH calo).

3. Omit events with $\angle(f, \mathrm{beam}) < 10°$ for a charged fermions.

W^+ invariant-mass distribution for $\mathrm{e}^+\mathrm{e}^- \to \mathrm{u}\bar{\mathrm{d}}\mu^-\bar{\nu}_\mu$:



Relative corrections to $\mathrm{d}\sigma/\mathrm{d}M$ for $\mathrm{e^+e^-} \to \mathrm{u\bar{d}}\mu^-\bar{\nu}_\mu$:





Internal check: subtraction method vs. phase-space slicing



2.1 Theoretical uncertainties

Estimate quite difficult !

General comments on the DPA:

- Threshold region: DPA not reliable.
  (Background diagrams important!)

  $\Rightarrow$ RacoonWW yields (improved) Born-level accuracy.

- Above threshold, $\sqrt{s} > 2M_\mathrm{W} + n\Gamma_\mathrm{W}$ with $n = \mathcal{O}(3\text{--}5)$:

  corrections to background diagrams
  $$\sim \frac{\alpha}{\pi} \frac{\Gamma_\mathrm{W}}{\sqrt{s} - 2M_\mathrm{W}} \ln(\cdots) \sim 0.1\%.$$
  (if background diagrams are not enhanced!)

Error estimates in practice: (Work in progress!)

- Estimate corrections to background diagrams
  $$\sim \frac{\alpha}{\pi} \text{(full Born c.s. – DPA Born c.s.)}$$
- Internal ambiguities in DPA:
  Compare results for different on-shell projections.
- Influence of higher-order effects:
  Compare different input or resummation schemes.

2.2 Comparison to results of other authors

Jadach et al. '99
$\mathcal{O}(\alpha)$ corr's to $e^+e^- \to WW \to 4f$
with YFSWW

Montagna et al. '99
$e^+e^- \to 4f + \gamma$ with $m_f \neq 0$
for some final states with ALPHA

others ?

$\to$ Talk of G. Passarino (?)

Jegerlehner, Kolodziej '99

$e^+e^- \to u\bar{d}\mu^-\bar{\nu}_\mu\gamma$ with $m_f \neq 0$

Full real correction to total cross section $\sigma_{\mathrm{real}}/\,\mathrm{fb}$:
(mass- and IR-singular, $m_\gamma = 10^{-6}\,\mathrm{GeV}$)

$\sqrt{s}/\,\mathrm{GeV}$	189	500	2000	10000	$\Delta E_\gamma/\,\mathrm{GeV}$
J/K '99	1285(1)	571(1)			0.001
	1285.0(8)	570.0(6)	82.5(3)	6.15(6)	0.1
	1286.3(8)	570.1(5)	83.1(3)	6.10(6)	1.0
RacoonWW	1284.6(6)	568.5(4)	82.2(1)	6.31(2)	—

$\Rightarrow$ Reasonable agreement,
but more precise comparisons desirable.